

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 March 2002 (28.03.2002)

PCT

(10) International Publication Number
WO 02/24663 A2

(51) International Patent Classification⁷: **C07D 239/00**

(21) International Application Number: **PCT/JP01/07766**

(22) International Filing Date:
7 September 2001 (07.09.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
2000-283113 19 September 2000 (19.09.2000) JP
2001-142975 14 May 2001 (14.05.2001) JP

(71) Applicant (for all designated States except US): **SUMITOMO CHEMICAL COMPANY, LIMITED** [JP/JP];
5-33, Kitahama 4-chome, Chuo-ku, Osaka-shi, Osaka 541-0041 (JP).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MIZUNO, Hajime** [JP/JP]; 4-9-17-323, Sakuragaoka, Minoo-shi, Osaka 562-0046 (JP). **SAKAMOTO, Noriyasu** [JP/JP]; 4-1-4, Kitasakurazuka, Toyonaka-shi, Osaka 560-0022 (JP). **KINOSHITA, Yoshiharu** [JP/JP]; 4-9-17-329, Sakuragaoka, Minoo-shi, Osaka 562-0046 (JP).

(74) Agents: **AOYAMA, Tamotsu et al.**; Aoyama & Partners, IMP Building, 3-7, Shiromi 1-chome, Chuo-ku, Osaka-shi, Osaka 540-0001 (JP).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 02/24663 A2

(54) Title: PYRIMIDINE COMPOUNDS AND THEIR USE



(1)

(57) Abstract: Pyrimidine compounds of formula (1), wherein R¹ is C₃-C₇ alkynyl optionally substituted with halogen; R² and R³ are independently hydrogen or the like; and R⁴ is C₃-C₇ alkynyloxy optionally substituted with halogen, optionally substituted phenyl, or the like have an excellent pesticidal effect.

DESCRIPTION
PYRIMIDINE COMPOUNDS AND THEIR USE

Technical Field

5 The present invention relates to certain pyrimidine compounds and their use.

Background Art


Various compounds have been used in the past for the purpose of pest
10 control. However, since not all these compounds have a satisfactory effect, there has been a demand for the development of compounds which are novel as pesticides and have a satisfactory effect.

It is an object of the present invention to provide novel compounds having a pesticidal effect and pesticidal compositions characterized in that
15 these compounds are contained as active ingredients.

Disclosure of the Invention

The present inventors have extensively studied in order to search for compounds having an excellent pesticidal effect. As a result, they have
20 found that the compounds of formula (1) as depicted below have pesticidal activity, thereby completing the present invention.

That is, the present invention provides pyrimidine compounds of formula (1):



25 (hereinafter referred to as the present compound(s))


wherein R^1 is C_3 - C_7 alkynyl optionally substituted with halogen;

R^2 and R^3 are independently hydrogen, halogen or C_1 - C_4 alkyl; and

R^4 is C_3 - C_7 alkynyloxy optionally substituted with halogen; C_3 - C_8 cycloalkoxy optionally substituted with halogen, hydroxy, C_1 - C_4 alkyl or C_1 - C_4

5 alkoxy; ethynyl substituted with C_3 - C_5 alkyl; or

a group of formula $-A^1R^5$ wherein A^1 is a single bond, oxygen, sulfur, carbonyl,



wherein R^8 is C_1 - C_4 alkyl; when A^1 is a single bond oxygen, or sulfur, then R^5 is optionally substituted phenyl or optionally sub-

stituted C_7 - C_9 aralkyl; or when A^1 is carbonyl, $\begin{array}{c} \diagup \\ \text{C}=\text{CH}_2 \\ \diagdown \end{array}$ or $\begin{array}{c} \diagup \\ \text{C}=\text{NOR}^8 \\ \diagdown \end{array}$

10 wherein R^8 is C_1 - C_4 alkyl, then R^5 is optionally substituted phenyl; or

a group of formula $-NR^6R^7$ wherein R^6 is C_1 - C_7 alkyl, C_1 - C_3 haloalkyl, C_2 - C_4 (alkoxymethyl), C_2 - C_4 (haloalkoxymethyl), C_3 - C_6 alkenyl, C_3 - C_6 haloalkenyl, C_3 - C_7 alkynyl, cyanomethyl, optionally substituted phenyl, or optionally substituted with C_7 - C_9 aralkyl; and R^7 is hydrogen, C_1 - C_7 alkyl, C_1 - C_3 haloalkyl, C_2 - C_4 (alkoxymethyl), C_2 - C_4 (haloalkoxymethyl), C_3 - C_6 alkenyl, C_3 - C_6 haloalkenyl, C_3 - C_7 alkynyl, cyanomethyl, optionally substituted phenyl, or optionally substituted C_7 - C_9 aralkyl;

wherein the substituent in the optionally substituted phenyl and in the optionally substituted C_7 - C_9 aralkyl is at least one selected from halogen, hydroxy, cyano, nitro, phenyl, phenoxy, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkoxy, C_1 - C_3 alkylthio, C_1 - C_4 haloalkylthio, C_3 - C_7 alkynyloxy, C_2 - C_6 (alkoxyalkoxy), C_2 - C_6 (alkylcarbonyl), and C_2 - C_6 (alkylcarbonyloxy).

The present invention further provides pesticidal compositions containing the present compounds as active ingredients.

Mode for Carrying Out the Invention

The C₃-C₇ alkynyl optionally substituted with halogen, which is represented by R¹, may include 2-propynyl, 2-butynyl, 3-butynyl, 2-pentynyl, 3-pentynyl, 2-heptynyl, 4,4-dimethyl-2-pentynyl, 3-fluoro-2-propynyl, 3-chloro-
5 2-propynyl, 3-bromo-2-propynyl, 3-iodo-2-propynyl, 3-trifluoromethyl-2-propynyl, 1-methyl-2-propynyl, 1-methyl-2-butynyl, 4-fluoro-2-butynyl, 4-chloro-2-butynyl, 4,4-difluoro-2-butynyl, and 1,1-dimethyl-2-propynyl.

The C₁-C₄ alkyl, which is represented by R² or R³, may include methyl, ethyl, propyl, 1-methylethyl, butyl, 1-methylpropyl, 2-methylpropyl, and 1,1-
10 dimethylethyl.

The C₃-C₇ alkynyloxy optionally substituted with halogen, which is represented by R⁴, may include 2-propynyloxy, 2-butynyloxy, 3-butynyloxy, 2-pentynyloxy, 3-pentynyloxy, 2-heptynyloxy, 4,4-dimethyl-2-pentynyloxy, 3-fluoro-2-propynyloxy, 3-chloro-2-propynyloxy, 3-bromo-2-propynyloxy, 3-tri-
15 fluoromethyl-2-propynyloxy, 1-methyl-2-propynyloxy, 1-methyl-2-butynyloxy, 4-fluoro-2-butynyloxy, 4-chloro-2-butynyloxy, 4,4-difluoro-2-butynyloxy, and 1,1-dimethyl-2-propynyloxy.

The C₃-C₈ cycloalkoxy optionally substituted with halogen, hydroxy, C₁-C₄ alkyl, or C₁-C₄ alkoxy, which is represented by R⁴, may include cyclo-
20 propoxy, cyclopentyloxy, 2-methylcyclopentyloxy, cyclohexyloxy, 1-methylcyclohexyloxy, 2-methylcyclohexyloxy, 3-methylcyclohexyloxy, 4-methylcyclohexyloxy, 2-fluorocyclohexyloxy, 3-fluorocyclohexyloxy, 4-fluorocyclohexyloxy, 2-chlorocyclohexyloxy, 3-chlorocyclohexyloxy, 4-chlorocyclohexyloxy, 2,3-dimethylcyclohexyloxy, 2-hydroxycyclohexyloxy, 3-hydroxycyclohexyloxy, 2-me-
25 thoxycyclohexyloxy, 3-methoxycyclohexyloxy, 2-ethylcyclohexyloxy, cycloheptyloxy, cyclooctyloxy, 4-hydroxycyclohexyloxy, 2,2-dimethylcyclopropoxy, and 4-methoxycyclohexyloxy.

The C₇-C₉ aralkyl, which is represented by R⁵, may include benzyl, 1-

phenylethyl, and 2-phenylethyl, wherein the aralkyl group may be optionally substituted on the aryl or alkyl moiety, or on both moieties.

For the substituents on the phenyl or C₇-C₉ aralkyl group, which is represented by R⁵, the C₁-C₄ alkyl may include methyl, ethyl, propyl, 1-methylethyl, and butyl; the C₁-C₄ haloalkyl may include trifluoromethyl, perfluoroethyl, and 2,2,2-trifluoroethyl; the C₁-C₄ alkoxy may include methoxy, ethoxy, propoxy, and 1-methylethoxy; the C₁-C₄ haloalkoxy may include trifluoromethoxy, perfluoroethoxy, and 2,2,2-trifluoroethoxy; the C₁-C₃ alkylthio may include methylthio and ethylthio; the C₁-C₄ haloalkylthio may include trifluoromethylthio; the C₃-C₇ alkynyloxy may include propynyloxy; the C₂-C₆ (alkoxyalkoxy) may include methoxymethoxy and ethoxymethoxy; the C₂-C₆ (alkylcarbonyl) may include acetyl and propionyl; the C₂-C₆ (alkylcarbonyloxy) may include acetoxymethoxy and isobutyryloxy.

Specific examples of the optionally substituted phenyl, which is represented by R⁵, may include phenyl, 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 2,3-difluorophenyl, 2,4-difluorophenyl, 2,5-difluorophenyl, 2,6-difluorophenyl, 3,4-difluorophenyl, 3,5-difluorophenyl, 2-fluoro-3-trifluoromethylphenyl, 2-fluoro-6-trifluoromethylphenyl, 2,3-difluoro-6-trifluoromethylphenyl, 2,3,6-trifluorophenyl, 2,4,6-trifluorophenyl, 2-chloro-3-fluorophenyl, 3-chloro-2-fluorophenyl, 2-chloro-4-fluorophenyl, 2-chloro-5-fluorophenyl, 2-chloro-6-fluorophenyl, 2-chloro-4,6-difluorophenyl, 2,3,4,6-tetrafluorophenyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 2,3-dichlorophenyl, 2,4-dichlorophenyl, 2,5-dichlorophenyl, 2,6-dichlorophenyl, 3,4-dichlorophenyl, 3,5-dichlorophenyl, 2,6-dichloro-4-fluorophenyl, 2,3,6-trichlorophenyl, 2,4,6-trichlorophenyl, 2-bromophenyl, 3-bromophenyl, 4-bromophenyl, 2-trifluoromethylphenyl, 3-trifluoromethylphenyl, 4-trifluoromethylphenyl, 3,5-bistrifluoromethylphenyl, 2-perfluoroethylphenyl, 3-(2,2,2-trifluoroethyl)phenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, 2,3-

dimethylphenyl, 2,4-dimethylphenyl, 2,5-dimethylphenyl, 2,6-dimethylphenyl, 3,4-dimethylphenyl, 3,5-dimethylphenyl, 2-ethylphenyl, 3-ethylphenyl, 4-ethylphenyl, 2-propylphenyl, 3-propylphenyl, 4-propylphenyl, 2-(1-methylethyl)phenyl, 3-(1-methylethyl)phenyl, 4-(1-methylethyl)phenyl, 4-
5 (1,1-dimethylethyl)phenyl, 2-methylthiophenyl, 3-methylthiophenyl, 4-methylthiophenyl, 2-ethylthiophenyl, 2-trifluoromethylthiophenyl, 3-trifluoromethylthiophenyl, 4-trifluoromethylthiophenyl, 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 2-ethoxyphenyl, 2-propoxyphenyl, 2-(1-methylethoxy)phenyl, 2-trifluoromethoxyphenyl, 3-trifluoromethoxyphenyl,
10 4-trifluoromethoxyphenyl, 2-(2,2,2-trifluoroethoxy)phenyl, 2-cyanophenyl, 2-nitrophenyl, 3-nitrophenyl, 4-nitrophenyl, 2-phenylphenyl, 2-phenoxyphenyl, 3-phenoxyphenyl, and 4-phenoxyphenyl.

Specific examples of the optionally substituted C₇-C₉ aralkyl, which is represented by R⁵, may include benzyl, 2-fluorobenzyl, 3-fluorobenzyl, 4-
15 fluorobenzyl, 2,3-difluorobenzyl, 2,4-difluorobenzyl, 2,5-difluorobenzyl, 2,6-difluorobenzyl, 3,4-difluorobenzyl, 3,5-difluorobenzyl, 2-fluoro-3-trifluoromethylbenzyl, 2-fluoro-6-trifluoromethylbenzyl, 2,3-difluoro-6-trifluoromethylbenzyl, 2,3,6-trifluorobenzyl, 2,4,6-trifluorobenzyl, 2-chloro-3-fluorobenzyl, 3-chloro-2-fluorobenzyl, 2-chloro-4-fluorobenzyl, 2-chloro-5-fluorobenzyl, 2-
20 chloro-6-fluorobenzyl, 2-chloro-4,6-difluorobenzyl, 2,3,4,6-tetrafluorobenzyl, 2-chlorobenzyl, 3-chlorobenzyl, 4-chlorobenzyl, 2,3-dichlorobenzyl, 2,4-dichlorobenzyl, 2,5-dichlorobenzyl, 2,6-dichlorobenzyl, 3,4-dichlorobenzyl, 3,5-dichlorobenzyl, 2,6-dichloro-4-fluorobenzyl, 2,3,6-trichlorobenzyl, 2,4,6-trichlorobenzyl, 2-bromobenzyl, 3-bromobenzyl, 4-bromobenzyl, 2-trifluoro-
25 methylbenzyl, 3-trifluoromethylbenzyl, 4-trifluoromethylbenzyl, 3,5-bis-trifluoromethylbenzyl, 2-perfluoroethylbenzyl, 3-(2,2,2-trifluoroethyl)benzyl, 2-methylbenzyl, 3-methylbenzyl, 4-methylbenzyl, 2,3-dimethylbenzyl, 2,4-dimethylbenzyl, 2,5-dimethylbenzyl, 2,6-dimethylbenzyl, 3,4-dimethylbenzyl,

3,5-dimethylbenzyl, 2-ethylbenzyl, 3-ethylbenzyl, 4-ethylbenzyl, 2-propylbenzyl, 3-propylbenzyl, 4-propylbenzyl, 2-(1-methylethyl)benzyl, 3-(1-methylethyl)benzyl, 4-(1-methylethyl)benzyl, 4-(1,1-dimethylethyl)benzyl, 2-methylthiobenzyl, 3-methylthiobenzyl, 4-methylthiobenzyl, 2-ethylthiobenzyl, 2-trifluoromethylthiobenzyl, 3-trifluoromethylthiobenzyl, 4-trifluoromethylthiobenzyl, 2-methoxybenzyl, 3-methoxybenzyl, 4-methoxybenzyl, 2-ethoxybenzyl, 2-propoxybenzyl, 2-(1-methylethoxy)benzyl, 2-trifluoromethoxybenzyl, 3-trifluoromethoxybenzyl, 4-trifluoromethoxybenzyl, 2-(2,2,2-trifluoroethoxy)benzyl, 2-cyanobenzyl, 2-nitrobenzyl, 3-nitrobenzyl, 4-nitrobenzyl, 2-phenylbenzyl, 2-phenoxybenzyl, 3-phenoxybenzyl, 4-phenoxybenzyl, 2-phenylethyl, 2-(2-fluorophenyl)ethyl, 2-(3-fluorophenyl)ethyl, 2-(4-fluorophenyl)ethyl, 2-(2,3-difluorophenyl)ethyl, 2-(2,4-difluorophenyl)ethyl, 2-(2,5-difluorophenyl)ethyl, 2-(2,6-difluorophenyl)ethyl, 2-(3,4-difluorophenyl)ethyl, 2-(3,5-difluorophenyl)ethyl, 2-(2-fluoro-3-trifluoromethylphenyl)ethyl, 2-(2-fluoro-6-trifluoromethylphenyl)ethyl, 2-(2,3-difluoro-6-fluoromethylphenyl)ethyl, 2-(2,3,6-trifluorophenyl)ethyl, 2-(2,4,6-trifluorophenyl)ethyl, 2-(2-chloro-3-fluorophenyl)ethyl, 2-(3-chloro-2-fluorophenyl)ethyl, 2-(2-chloro-4-fluorophenyl)ethyl, 2-(2-chloro-5-fluorophenyl)ethyl, 2-(2-chloro-6-fluorophenyl)ethyl, 2-(2-chloro-4,6-difluorophenyl)ethyl, 2-(2,3,4,6-tetrafluorophenyl)ethyl, 2-(2-chlorophenyl)ethyl, 2-(3-chlorophenyl)ethyl, 2-(4-chlorophenyl)ethyl, 2-(2,3-dichlorophenyl)ethyl, 2-(2,4-dichlorophenyl)ethyl, 2-(2,5-dichlorophenyl)ethyl, 2-(2,6-dichlorophenyl)ethyl, 2-(3,4-dichlorophenyl)ethyl, 2-(3,5-dichlorophenyl)ethyl, 2-(2,6-dichloro-4-fluorophenyl)ethyl, 2-(2,3,6-trichlorophenyl)ethyl, 2-(2,4,6-trichlorophenyl)ethyl, 2-(2-bromophenyl)ethyl, 2-(3-bromophenyl)ethyl, 2-(4-bromophenyl)ethyl, 2-(2-trifluoromethylphenyl)ethyl, 2-(3-trifluoromethylphenyl)ethyl, 2-(4-trifluoromethylphenyl)ethyl, 2-(3,5-bistrifluoromethylphenyl)ethyl, 2-(2-perfluoroethylphenyl)ethyl, 2-(3-(2,2,2-trifluoroethyl)phenyl)ethyl, 2-(2-methylphen-

yl)ethyl, 2-(3-methylphenyl)ethyl, 2-(4-methylphenyl)ethyl, 2-(2,3-dimethylphenyl)ethyl, 2-(2,4-dimethylphenyl)ethyl, 2-(2,5-dimethylphenyl)ethyl, 2-(2,6-dimethylphenyl)ethyl, 2-(3,4-dimethylphenyl)ethyl, 2-(3,5-dimethylphenyl)ethyl, 2-(2-ethylphenyl)ethyl, 2-(3-ethylphenyl)ethyl, 2-(4-ethylphenyl)ethyl, 2-(2-propylphenyl)ethyl, 2-(3-propylphenyl)ethyl, 2-(4-propylphenyl)ethyl, 2-(2-(1-methylethyl)phenyl)ethyl, 2-(3-(1-methylethyl)phenyl)ethyl, 2-(4-(1-methylethyl)phenyl)ethyl, 2-(4-(1,1-dimethylethyl)phenyl)ethyl, 2-(2-methylthiophenyl)ethyl, 2-(3-methylthiophenyl)ethyl, 2-(4-methylthiophenyl)ethyl, 2-(2-ethylthiophenyl)ethyl, 2-(2-trifluoromethylthiophenyl)ethyl, 2-(3-trifluoromethylthiophenyl)ethyl, 2-(4-trifluoromethylthiophenyl)ethyl, 2-(2-methoxyphenyl)ethyl, 2-(3-methoxyphenyl)ethyl, 2-(4-methoxyphenyl)ethyl, 2-(2-ethoxyphenyl)ethyl, 2-(2-propoxyphenyl)ethyl, 2-(2-(1-methylethoxy)phenyl)ethyl, 2-(2-trifluoromethoxyphenyl)ethyl, 2-(3-trifluoromethoxyphenyl)ethyl, 2-(4-trifluoromethoxyphenyl)ethyl, 2-(2-perfluoroethoxyphenyl)ethyl, 2-(2-(2,2,2-trifluoroethoxy)phenyl)ethyl, 2-(2-cyanophenyl)ethyl, 2-(2-nitrophenyl)ethyl, 2-(3-nitrophenyl)ethyl, 2-(4-nitrophenyl)ethyl, 2-(2-phenylphenyl)ethyl, 2-(2-phenoxyphenyl)ethyl, 2-(3-phenoxyphenyl)ethyl, 2-(4-phenoxyphenyl)ethyl, 1-phenylethyl, 1-(2-fluorophenyl)ethyl, 1-(3-fluorophenyl)ethyl, 1-(4-fluorophenyl)ethyl, 1-(2,3-difluorophenyl)ethyl, 1-(2,4-difluorophenyl)ethyl, 1-(2,5-difluorophenyl)ethyl, 1-(2,6-difluorophenyl)ethyl, 1-(3,4-difluorophenyl)ethyl, 1-(3,5-difluorophenyl)ethyl, 1-(2-fluoro-3-trifluoromethylphenyl)ethyl, 1-(2-fluoro-6-trifluoromethylphenyl)ethyl, 1-(2,3-difluoro-6-fluoromethylphenyl)ethyl, 1-(2,3,6-trifluorophenyl)ethyl, 1-(2,4,6-trifluorophenyl)ethyl, 1-(2-chloro-3-fluorophenyl)ethyl, 1-(3-chloro-2-fluorophenyl)ethyl, 1-(2-chloro-4-fluorophenyl)ethyl, 1-(2-chloro-5-fluorophenyl)ethyl, 1-(2-chloro-6-fluorophenyl)ethyl, 1-(2-chloro-4,6-difluorophenyl)ethyl, 1-(2,3,4,6-tetrafluorophenyl)ethyl, 1-(2-chlorophenyl)ethyl, 1-(3-chlorophenyl)ethyl, 1-(4-chlorophenyl)ethyl, 1-(2,3-dichlorophenyl)ethyl, 1-(2,4-dichloro-

phenyl)ethyl, 1-(2,5-dichlorophenyl)ethyl, 1-(2,6-dichlorophenyl)ethyl, 1-(3,4-dichlorophenyl)ethyl, 1-(3,5-dichlorophenyl)ethyl, 1-(2,6-dichloro-4-fluorophenyl)ethyl, 1-(2,3,6-trichlorophenyl)ethyl, 1-(2,4,6-trichlorophenyl)ethyl, 1-(2-bromophenyl)ethyl, 1-(3-bromophenyl)ethyl, 1-(4-bromophenyl)ethyl, 1-(2-trifluoromethylphenyl)ethyl, 1-(3-trifluoromethylphenyl)ethyl, 1-(4-trifluoromethylphenyl)ethyl, 1-(3,5-bistrifluoromethylphenyl)ethyl, 1-(2-perfluoroethylphenyl)ethyl, 1-(3-(2,2,2-trifluoroethyl)phenyl)ethyl, 1-(2-methylphenyl)ethyl, 1-(3-methylphenyl)ethyl, 1-(4-methylphenyl)ethyl, 1-(2,3-dimethylphenyl)ethyl, 1-(2,4-dimethylphenyl)ethyl, 1-(2,5-dimethylphenyl)ethyl, 1-(2,6-dimethylphenyl)ethyl, 1-(3,4-dimethylphenyl)ethyl, 1-(3,5-dimethylphenyl)ethyl, 1-(2-ethylphenyl)ethyl, 1-(3-ethylphenyl)ethyl, 1-(4-ethylphenyl)ethyl, 1-(2-propylphenyl)ethyl, 1-(3-propylphenyl)ethyl, 1-(4-propylphenyl)ethyl, 1-(2-(1-methylethyl)phenyl)ethyl, 1-(3-(1-methylethyl)phenyl)ethyl, 1-(4-(1-methylethyl)phenyl)ethyl, 1-(4-(1,1-dimethylethyl)phenyl)ethyl, 1-(2-methylthiophenyl)ethyl, 1-(3-methylthiophenyl)ethyl, 1-(4-methylthiophenyl)ethyl, 1-(2-ethylthiophenyl)ethyl, 1-(2-trifluoromethylthiophenyl)ethyl, 1-(3-trifluoromethylthiophenyl)ethyl, 1-(4-trifluoromethylthiophenyl)ethyl, 1-(2-methoxyphenyl)ethyl, 1-(3-methoxyphenyl)ethyl, 1-(4-methoxyphenyl)ethyl, 1-(2-ethoxyphenyl)ethyl, 1-(2-propoxyphenyl)ethyl, 1-(2-(1-methylethoxy)phenyl)ethyl, 1-(2-trifluoromethoxyphenyl)ethyl, 1-(3-trifluoromethoxyphenyl)ethyl, 1-(4-trifluoromethoxyphenyl)ethyl, 1-(2-perfluoroethoxyphenyl)ethyl, 1-(2-(2,2,2-trifluoroethoxy)phenyl)ethyl, 1-(2-cyanophenyl)ethyl, 1-(2-nitrophenyl)ethyl, 1-(3-nitrophenyl)ethyl, 1-(4-nitrophenyl)ethyl, 1-(2-phenylphenyl)ethyl, 1-(2-phenoxyphenyl)ethyl, 1-(3-phenoxyphenyl)ethyl, and 1-(4-phenoxyphenyl)ethyl.

The C₁-C₇ alkyl, which is represented by R⁶ or R⁷, may include methyl, ethyl, propyl, 1-methylethyl, butyl, 2-methylpropyl, 1-methylpropyl, 1,1-dimethylethyl, pentyl, 3-methylbutyl, 2,2-dimethylpropyl, 1,1-dimethylpropyl,

1-ethylpropyl, hexyl, 5-methylpentyl, 2-ethylbutyl, 3-methylpentyl, 1,3-dimethylbutyl, heptyl, and 1-ethyl-1-methylbutyl.

The C₁-C₃ haloalkyl, represented by R⁶ or R⁷, may include difluoromethyl, dibromofluoromethyl, 1-chloroethyl, 1-bromoethyl, 2,2,2-trifluoroethyl, 2-fluoroethyl, 2-chloroethyl, 2,2-difluoroethyl, 3-fluoropropyl, 3,3,3-trifluoropropyl, 2-fluoropropyl, and 2-bromopropyl.

The C₂-C₄ (alkoxymethyl), which is represented by R⁶ or R⁷, may include methoxymethyl, ethoxymethyl, propoxymethyl, and 1-methylethoxymethyl.

10 The C₂-C₄ (haloalkoxymethyl), which is represented by R⁶ or R⁷, may include chloromethoxymethyl, bromomethoxymethyl, 2-chloroethoxymethyl, 2-bromoethoxymethyl, and 2,2,2-trifluoroethoxymethyl.

15 The C₃-C₆ alkenyl, which is represented by R⁶ or R⁷, may include 2-propenyl, 2-butenyl, 3-butenyl, 1-methyl-2-propenyl, 2-methyl-2-propenyl, 2-pentenyl, 1-methyl-2-butenyl, 3-methyl-3-butenyl, 1-ethyl-2-propenyl, 2-hexenyl, 2-methyl-2-pentenyl, 3-methyl-2-pentenyl, 4-methyl-2-pentenyl, 1-methyl-3-pentenyl, 4-methyl-3-pentenyl, 1-methyl-4-pentenyl, 3-methyl-4-pentenyl, and 4-methyl-4-pentenyl.

20 The C₃-C₆ haloalkenyl, which is represented by R⁶ or R⁷, may include 3-chloro-2-propenyl, 2-chloro-2-propenyl, 2-bromo-2-propenyl, 3,3-dichloro-2-propenyl, 3,3-difluoro-2-propenyl, 2,3-dichloro-2-propenyl, 2,3-dibromo-2-propenyl, 3-chloro-4,4,4-trifluoro-2-butenyl, 3,4-dichloro-4,4-difluoro-2-butenyl, 3,4,4,4-tetrafluoro-2-butenyl, 4,4-dibromo-2-butenyl, 6,6-dichloro-5-hexenyl, and 6,6-dibromo-5-hexenyl.

25 The C₃-C₇ alkynyl, which is represented by R⁶ or R⁷, may include 2-propynyl, 2-butyne, 2-pentyne, 4,4-dimethyl-2-pentyne, 1-methyl-2-propynyl, and 1,1-dimethyl-2-propynyl.

The substituents in the optionally substituted phenyl and in the op-

tionally substituted C₇-C₉ aralkyl, which are represented by R⁶ or R⁷, may include the above substituents for R⁵, and specific examples thereof may include the above groups for R⁵.

The embodiments of the present compounds may include the following compounds.

The pyrimidine compounds of formula (1) wherein R¹ is 2-propynyl optionally substituted with halogen, 2-butynyl optionally substituted with halogen, or 2-methyl-2-butynyl optionally substituted with halogen;

The pyrimidine compounds of formula (1) wherein R² and R³ are both hydrogen;

The pyrimidine compounds of formula (1) wherein R² is hydrogen and R³ is chlorine or fluorine;

The pyrimidine compounds of formula (1) wherein R¹ is C₃-C₇ alkyl optionally substituted with halogen; R² and R³ are both hydrogen; and R⁴ is 2-halophenyl; the pyrimidine compounds of formula (1) wherein R¹ is C₃-C₇ alkyl optionally substituted with halogen; R² and R³ are both hydrogen; and R⁴ is 3-halophenyl; the pyrimidine compounds of formula (1) wherein R¹ is C₃-C₇ alkyl optionally substituted with halogen; R² and R³ are both hydrogen; and R⁴ is 2,3-dihalophenyl; the pyrimidine compounds of formula (1) wherein R¹ is C₃-C₇ alkyl optionally substituted with halogen; R² and R³ are both hydrogen; and R⁴ is 2,6-dihalophenyl;


The pyrimidine compounds of formula (1) wherein R¹ is C₃-C₇ alkyl optionally substituted with halogen; R² and R³ are both hydrogen; and R⁴ is 2-halophenoxy; the pyrimidine compounds of formula (1) wherein R¹ is C₃-C₇ alkyl optionally substituted with halogen; R² and R³ are both hydrogen; and R⁴ is 3-halophenoxy; the pyrimidine compounds of formula (1) wherein R¹ is C₃-C₇ alkyl optionally substituted with halogen; R² and R³ are both hydrogen; and R⁴ is 2,3-dihalophenoxy; the pyrimidine compounds of formula (1)

wherein R^1 is C_3 - C_7 alkyl optionally substituted with halogen; R^2 and R^3 are both hydrogen; and R^4 is 2,6-dihalophenoxy;

The pyrimidine compounds of formula (1) wherein R^1 is C_3 - C_7 alkyl optionally substituted with halogen; R^2 and R^3 are both hydrogen; and R^4 is 2-halobenzyl; the pyrimidine compounds of formula (1) wherein R^1 is C_3 - C_7 alkyl optionally substituted with halogen; R^2 and R^3 are both hydrogen; and R^4 is 3-halobenzyl; the pyrimidine compounds of formula (1) wherein R^1 is C_3 - C_7 alkyl optionally substituted with halogen; R^2 and R^3 are both hydrogen; and R^4 is 2,3-dihalobenzyl; the pyrimidine compounds of formula (1) wherein R^1 is C_3 - C_7 alkyl optionally substituted with halogen; R^2 and R^3 are both hydrogen; and R^4 is 2,6-dihalophenyl; the pyrimidine compounds of formula (1) wherein R^1 is C_3 - C_7 alkyl optionally substituted with halogen; R^2 and R^3 are both hydrogen; R^4 is a group of formula $-N(C_2H_5)R^7$; and R^7 is C_3 - C_5 2-alkynyl optionally substituted with halogen; the pyrimidine compounds of formula (1) wherein R^1 is C_3 - C_7 alkyl optionally substituted with halogen; R^2 and R^3 are both hydrogen; and R^4 is a group of formula $-N(C_2H_5)R^7$; and R^7 is C_3 - C_5 alkenyl optionally substituted with halogen; and the pyrimidine compounds of formula (1) wherein R^1 is C_3 - C_7 alkyl optionally substituted with halogen; R^2 and R^3 are both hydrogen; and R^4 is a group of formula $-N(C_2H_5)R^7$; and R^7 is C_2 - C_3 alkyl.

Specific examples of the present compounds are show below.

The compounds of formula (25):



(25)

wherein Q is oxygen, R^1 is 2-propynyl, R^2 and R^3 are both hydrogen, and the

substituent(s) (R^9)_p on the benzene ring are selected from those of Table 1.

TABLE 1

H	2-CH ₃	3-SCF ₃	2-Cl-3-F
2-Cl	3-CH ₃	4-SCF ₃	2-Cl-6-F
3-Cl	4-CH ₃	2,3-diF	2-Cl-4,6-diF
4-Cl	2-CF ₃	2,4-diF	2,3-diCl
2-F	3-CF ₃	2,5-diF	2,3-diCH ₃
3-F	4-CF ₃	2,6-diF	2,3,6-triF
4-F	2-OCF ₃	3,4-diF	2,4,6-triF
2-OCH ₃	3-OCF ₃	3,5-diF	2,3,4,6-tetraF
3-OCH ₃	4-OCF ₄	2-F-3-CF ₃	
4-OCH ₄	2-SCF ₃	2-F-6-CF ₃	

- 5 The compounds of formula (25) wherein Q is oxygen, R¹ is 2-propynyl, R² is methyl, R³ is hydrogen, and the substituent(s) (R^9)_p on the benzene ring are selected from those of Table 2.

TABLE 2

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is oxygen, R¹ is 2-propynyl, R² and R³ are both methyl, and the substituent(s) (R^9)_p on the benzene ring are those of Table 3.

TABLE 3

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is oxygen, R¹ is 2-propynyl, R² is hydrogen, R³ is methyl, and the substituent(s) (R⁹)_p on the benzene ring
5 are those of Table 4.

TABLE 4

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is oxygen, R¹ is 2-butyryl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring
10 are those of Table 5.

TABLE 5

H	3-CN	2-OPh	2,6-diCH ₃
2-Cl	4-CN	3-OPh	2-F-3-CF ₃
3-Cl	2-SCH ₃	4-OPh	2-F-6-CF ₃
4-Cl	3-SCH ₃	2-ethoxy	2-Cl-3-F
2-F	4-SCH ₃	3-ethoxy	2-Cl-4-F
3-F	2-OCF ₃	4-ethoxy	2-Cl-5-F
4-F	3-OCF ₃	2-isopropyl	2-Cl-6-F
2-Br	4-OCF ₃	3-isopropyl	3-Cl-2-F
3-Br	2-SCF ₃	4-isopropyl	2-Cl-4,6-diF
4-Br	3-SCF ₃	2,3-diF	2,3-diCl
2-I	4-SCF ₃	2,4-diF	2,4-diCl
3-I	2-CH ₂ CH ₃	2,5-diF	2,5-diCl
4-I	3-CH ₂ CH ₃	2,6-diF	2,6-diCl
2-OCH ₃	4-CH ₂ CH ₃	3,4-diF	3,4-diCl
3-OCH ₃	2-propyl	3,5-diF	3,5-diCl
4-OCH ₃	3-propyl	3,5-diCF ₃	2,3,6-triCl
2-CH ₃	4-propyl	2,3,6-triF	2,4,6-triCl
3-CH ₃	2-NO ₂	2,4,6-triF	2,6-diCl-4-F
4-CH ₃	3-NO ₂	3,4-diCH ₃	2,3-diF-6-CF ₃
2-CF ₃	4-NO ₂	3,5-diCH ₃	2,3,4,6-tetraF
3-CF ₃	2-Ph	2,3-diCH ₃	
4-CF ₃	3-Ph	2,4-diCH ₃	
2-CN	4-Ph	2,5-diCH ₃	

The compounds of formula (25) wherein Q is oxygen, R¹ is 2-butyryl, R² is methyl, R³ is hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 6.

TABLE 6

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is oxygen, R¹ is 2-butynyl, R² and R³ are both methyl, and the substituent(s) (R⁹)_p on the benzene ring
 5 are those of Table 7.

TABLE 7

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is oxygen, R¹ is 2-butynyl, R² is hydrogen, R³ is methyl, and the substituent(s) (R⁹)_p on the benzene ring
 10 are those of Table 8.

TABLE 8

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is oxygen, R¹ is 1-methyl-2-butynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 9.
 15

TABLE 9

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is oxygen, R¹ is 1-methyl-2-propynyl, R² are R³ are both hydrogen, and the substituent(s) (R⁹)_p on the
 5 benzene ring are those of Table 10.

TABLE 10

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is oxygen, R¹ is 3-pentynyl,
 10 R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 11.

TABLE 11

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

15 The compounds formula (25) wherein Q is oxygen, R¹ is 2-butynyl, R² is hydrogen, R³ is fluorine, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 12.

TABLE 12

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is CH₂, R¹ is 2-propynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring
 5 are those of Table 13.

TABLE 13

H	3-CH ₃	2-CH ₃	2-Cl-3-F
2-Cl	4-CH ₃	2-CN	2-Cl-6-F
3-Cl	2-CF ₃	4-SCF ₃	2-Cl-4,6-diF
4-Cl	3-CF ₃	2,3-diF	2,3-diCl
2-F	4-CF ₃	2,4-diF	2,3-diCH ₃
3-F	2-OCF ₃	2,5-diF	2,3,6-triF
4-F	3-OCF ₃	2,6-diF	2,4,6-triF
2-OCH ₃	4-OCF ₃	3,4-diF	2,3,4,6-tetraF
3-OCH ₃	2-SCF ₃	3,5-diF	3-CN
4-OCH ₃	3-SCF ₃	2-F-3-CF ₃	4-CN

The compounds of formula (25) wherein Q is CH₂, R¹ is 2-propynyl, R²
 10 is methyl, R³ is hydrogen, and the substituent(s) (R⁹)_p on the benzene ring
 are those of Table 14.

TABLE 14

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is CH₂, R¹ is 2-propynyl, R² and R³ are both methyl, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 15.

TABLE 15

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is CH₂, R¹ is 2-propynyl, R² is hydrogen, R³ is methyl, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 16.

TABLE 16

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is CH₂, R¹ is 2-butynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 17.

TABLE 17

H	3-CN	2-OPh	2,6-diCH ₃
2-Cl	4-CN	3-OPh	2-F-3-CF ₃
3-Cl	2-SCH ₃	4-OPh	2-F-6-CF ₃
4-Cl	3-SCH ₃	2-ethoxy	2-Cl-3-F
2-F	4-SCH ₃	3-ethoxy	2-Cl-4-F
3-F	2-OCF ₃	4-ethoxy	2-Cl-5-F
4-F	3-OCF ₃	2-isopropyl	2-Cl-6-F
2-Br	4-OCF ₃	3-isopropyl	3-Cl-2-F
3-Br	2-SCF ₃	4-isopropyl	2-Cl-4,6-diF
4-Br	3-SCF ₃	2,3-diF	2,3-diCl
2-I	4-SCF ₃	2,4-diF	2,4-diCl
3-I	2-CH ₂ CH ₃	2,5-diF	2,5-diCl
4-I	3-CH ₂ CH ₃	2,6-diF	2,6-diCl
2-OCH ₃	4-CH ₂ CH ₃	3,4-diF	3,4-diCl
3-OCH ₃	2-propyl	3,5-diF	3,5-diCl
4-OCH ₃	3-propyl	3,5-diCF ₃	2,3,6-triCl
2-CH ₃	4-propyl	2,3,6-triF	2,4,6-triCl
3-CH ₃	2-NO ₂	2,4,6-triF	2,6-diCl-4-F
4-CH ₃	3-NO ₂	3,4-diCH ₃	2,3-diF-6-CF ₃
2-CF ₃	4-NO ₂	3,5-diCH ₃	2,3,4,6-tetraF
3-CF ₃	2-Ph	2,3-diCH ₃	
4-CF ₃	3-Ph	2,4-diCH ₃	
2-CN	4-Ph	2,5-diCH ₃	

The compounds of formula (25) wherein Q is CH₂, R¹ is 2-butyryl, R² is methyl, R³ is hydrogen, and the substituent(s) (R⁹)_p on the benzene ring
5 are those of Table 18.

TABLE 18

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is CH₂, R¹ is 2-butyryl, R² and R³ are both methyl, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 19.

TABLE 19

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is CH₂, R¹ is 2-butyryl, R² is hydrogen, R³ is methyl, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 20.

TABLE 20

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is CH₂, R¹ is 2-pentyryl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 21.

TABLE 21

H	2-CH ₃	3-SCF ₃	2-Cl-3-F
2-Cl	3-CH ₃	4-SCF ₃	2-Cl-6-F
3-Cl	4-CH ₃	2,3-diF	2-Cl-4,6-diF
4-Cl	3-CF ₃	2,4-diF	2,3-diCl
2-F	3-CF ₃	2,5-diF	2,3-diCH ₃
3-F	4-CF ₃	2,6-diF	2,3,6-triF
4-F	2-OCF ₃	3,4-diF	2,4,6-triF
2-OCH ₃	3-OCF ₃	3,5-diF	2,3,4,6-tetraF
3-OCH ₃	4-OCF ₃	2-F-3-CF ₃	
4-OCH ₃	2-SCF ₃	2-F-6-CF ₃	

The compounds of formula (25) wherein Q is CH₂, R¹ is 1-methyl-2-butynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the
5 benzene ring are those of Table 22.

TABLE 22

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is CH₂, R¹ is 1-methyl-2-propynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the
10 benzene ring are those of Table 23.

TABLE 23

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds formula (25) wherein Q is CH₂, R¹ is 2-butyryl, R² is hydrogen, R³ is fluorine, and the substituent(s) (R⁹)_p on the benzene ring are
 5 those of Table 24.

TABLE 24

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is CHCH₃, R¹ is 2-propynyl,
 10 R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 25.

TABLE 25

H	2-CH ₃	3-SCF ₃	2-Cl-3-F
2-Cl	3-CH ₃	4-SCF ₃	2-Cl-6-F
3-Cl	4-CH ₃	2,3-diF	2-Cl-4,6-diF
4-Cl	3-CF ₃	2,4-diF	2,3-diCl
2-F	3-CF ₃	2,5-diF	2,3-diCH ₃
3-F	4-CF ₃	2,6-diF	2,3,6-triF
4-F	2-OCF ₃	3,4-diF	2,4,6-triF
2-OCH ₃	3-OCF ₃	3,5-diF	2,3,4,6-tetraF
3-OCH ₃	4-OCF ₃	2-F-3-CF ₃	
4-OCH ₃	2-SCF ₃	2-F-6-CF ₃	

The compounds of formula (25) wherein Q is CHCH₃, R¹ is 2-propynyl, R² is methyl, R³ is hydrogen, and the substituent(s) (R⁹)_p on the benzene ring
5 are those of Table 26.

TABLE 26

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is CHCH₃, R¹ is 2-propynyl,
10 R² and R³ are both methyl, and the substituent(s) (R⁹)_p on the benzene ring
are those of Table 27.

TABLE 27

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is CHCH₃, R¹ is 2-propynyl, R² is hydrogen, R³ is methyl, and the substituent(s) (R⁹)_p on the benzene ring
 5 are those of Table 28.

TABLE 28

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is CHCH₃, R¹ is 2-butynyl,
 10 R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring
 are those of Table 29.

TABLE 29

H	3-CN	2-OPh	2,6-diCH ₃
2-Cl	4-CN	3-OPh	2-F-3-CF ₃
3-Cl	2-SCH ₃	4-OPh	2-F-6-CF ₃
4-Cl	3-SCH ₃	2-ethoxy	2-Cl-3-F
2-F	4-SCH ₃	3-ethoxy	2-Cl-4-F
3-F	2-OCF ₃	4-ethoxy	2-Cl-5-F
4-F	3-OCF ₃	2-isopropyl	2-Cl-6-F
2-Br	4-OCF ₃	3-isopropyl	3-Cl-2-F
3-Br	2-SCF ₃	4-isopropyl	2-Cl-4,6-diF
4-Br	3-SCF ₃	2,3-diF	2,3-diCl
2-I	4-SCF ₃	2,4-diF	2,4-diCl
3-I	2-CH ₂ CH ₃	2,5-diF	2,5-diCl
4-I	3-CH ₂ CH ₃	2,6-diF	2,6-diCl
2-OCH ₃	4-CH ₂ CH ₃	3,4-diF	3,4-diCl
3-OCH ₃	2-propyl	3,5-diF	3,5-diCl
4-OCH ₃	3-propyl	3,5-diCF ₃	2,3,6-triCl
2-CH ₃	4-propyl	2,3,6-triF	2,4,6-triCl
3-CH ₃	2-NO ₂	2,4,6-triF	2,6-diCl-4-F
4-CH ₃	3-NO ₂	3,4-diCH ₃	2,3-diF-6-CF ₃
2-CF ₃	4-NO ₂	3,5-diCH ₃	2,3,4,6-tetraF
3-CF ₃	2-Ph	2,3-diCH ₃	
4-CF ₃	3-Ph	2,4-diCH ₃	
2-CN	4-Ph	2,5-diCH ₃	

The compounds of formula (25) wherein Q is CHCH₃, R¹ is 2-butyryl, R² is methyl, R³ is hydrogen, and the substituent(s) (R⁹)_p on the benzene ring
5 are those of Table 30.

TABLE 30

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is CHCH₃, R¹ is 2-butynyl, R² and R³ are both methyl, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 31.

TABLE 31

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is CHCH₃, R¹ is 2-butynyl, R² is hydrogen, R³ is methyl, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 32.

TABLE 32

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is CHCH₃, R¹ is 2-pentynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 33.

TABLE 33

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is CHCH₃, R¹ is 1-methyl-2-butynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the
5 benzene ring are those of Table 34.

TABLE 34

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is CHCH₃, R¹ is 1-methyl-2-propynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the
10 benzene ring are those of Table 35.

TABLE 35

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

15 The compounds of formula (25) wherein Q is CHCH₃, R¹ is 2-butynyl, R² is hydrogen, R³ is fluorine, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 36.

TABLE 36

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NH, R¹ is 2-propynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring
5 are those of Table 37.

TABLE 37

H	2-CH ₃	3-SCF ₃	2-Cl-3-F
2-Cl	3-CH ₃	4-SCF ₃	2-Cl-6-F
3-Cl	4-CH ₃	2,3-diF	2-Cl-4,6-diF
4-Cl	2-CF ₃	2,4-diF	2,3-diCl
2-F	3-CF ₃	2,5-diF	2,3-diCH ₃
3-F	4-CF ₃	2,6-diF	2,3,6-triF
4-F	2-OCF ₃	3,4-diF	2,4,6-triF
2-OCH ₃	3-OCF ₃	3,5-diF	2,3,4,6-tetraF
3-OCH ₃	4-OCF ₃	2-F-3-CF ₃	
4-OCH ₃	2-SCF ₃	2-F-6-CF ₃	

The compounds of formula (25) wherein Q is NH, R¹ is 2-butynyl, R²
10 and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring
are those of Table 38.

TABLE 38

H	2-CH ₃	3-SCF ₃	2-Cl-3-F
2-Cl	3-CH ₃	4-SCF ₃	2-Cl-6-F
3-Cl	4-CH ₃	2,3-diF	2-Cl-4,6-diF
4-Cl	2-CF ₃	2,4-diF	2,3-diCl
2-F	3-CF ₃	2,5-diF	2,3-diCH ₃
3-F	4-CF ₃	2,6-diF	2,3,6-triF
4-F	2-OCF ₃	3,4-diF	2,4,6-triF
2-OCH ₃	3-OCF ₃	3,5-diF	2,3,4,6-tetraF
3-OCH ₃	4-OCF ₃	2-F-3-CF ₃	
4-OCH ₃	2-SCF ₃	2-F-6-CF ₃	

The compounds of formula (25) wherein Q is NH, R¹ is 2-pentynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring
5 are those of Table 39.

TABLE 39

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₃, R¹ is 2-propynyl,
10 R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring
are those of Table 40.

TABLE 40

H	2-CH ₃	3-SCF ₃	2-Cl-3-F
2-Cl	3-CH ₃	4-SCF ₃	2-Cl-6-F
3-Cl	4-CH ₃	2,3-diF	2-Cl-4,6-diF
4-Cl	2-CF ₃	2,4-diF	2,3-diCl
2-F	3-CF ₃	2,5-diF	2,3-diCH ₃
3-F	4-CF ₃	2,6-diF	2,3,6-triF
4-F	2-OCF ₃	3,4-diF	2,4,6-triF
2-OCH ₃	3-OCF ₃	3,5-diF	2,3,4,6-tetraF
3-OCH ₃	4-OCF ₃	2-F-3-CF ₃	
4-OCH ₃	2-SCF ₃	2-F-6-CF ₃	

The compounds of formula (25) wherein Q is NCH₃, R¹ is 2-propynyl, R² is methyl, R³ is hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 41.

TABLE 41

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₃, R¹ is 2-propynyl, R² and R³ are both methyl, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 42.

TABLE 42

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₃, R¹ is 2-propynyl, R² is hydrogen, and R³ is methyl, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 43.

TABLE 43

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₃, R¹ is 2-butyryl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 44.

TABLE 44

H	2-CH ₃	3-SCF ₃	2-Cl-3-F
2-Cl	3-CH ₃	4-SCF ₃	2-Cl-6-F
3-Cl	4-CH ₃	2,3-diF	2-Cl-4,6-diF
4-Cl	2-CF ₃	2,4-diF	2,3-diCl
2-F	3-CF ₃	2,5-diF	2,3-diCH ₃
3-F	4-CF ₃	2,6-diF	2,3,6-triF
4-F	2-OCF ₃	3,4-diF	2,4,6-triF
2-OCH ₃	3-OCF ₃	3,5-diF	2,3,4,6-tetraF
3-OCH ₃	4-OCF ₃	2-F-3-CF ₃	
4-OCH ₃	2-SCF ₃	2-F-6-CF ₃	

The compounds of formula (25) wherein Q is NCH₃, R¹ is 2-butynyl, R² is methyl, R³ is hydrogen, and the substituent(s) (R⁹)_p on the benzene ring
5 are those of Table 45.

TABLE 45

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₃, R¹ is 2-butynyl,
10 R² and R³ are both methyl, and the substituent(s) (R⁹)_p on the benzene ring
are those of Table 46.

TABLE 46

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₃, R¹ is 2-butyryl, R² is hydrogen, R³ is methyl, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 47.

TABLE 47

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₃, R¹ is 2-pentyryl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 48.

TABLE 48

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₃, R¹ is 1-methyl-2-butyryl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 49.

TABLE 49

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₃, R¹ is 1-methyl-2-propynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the
 5 benzene ring are those of Table 50.

TABLE 50

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₃, R¹ is 2-butyryl,
 10 R² is hydrogen, R³ is fluorine, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 51.

TABLE 51

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

15 The compounds of formula (25) wherein Q is NCH₂CH₃, R¹ is 2-propynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 52.

TABLE 52

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₂CH₃, R¹ is 2-propynyl, R² is methyl, R³ is hydrogen, and the substituent(s) (R⁹)_p on the
5 benzene ring are those of Table 53.

TABLE 53

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₂CH₃, R¹ is 2-propynyl, R² and R³ are both methyl, and the substituent(s) (R⁹)_p on the
10 benzene ring are those of Table 54.

TABLE 54

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₂CH₃, R¹ is 2-propynyl, R² is hydrogen, R³ is methyl, and the substituent(s) (R⁹)_p on the
15 benzene ring are those of Table 55.

TABLE 55

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₂CH₃, R¹ is 2-butynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the
5 benzene ring are those of Table 56.

TABLE 56

H	3-CN	2-OPh	2,6-diCH ₃
2-Cl	4-CN	3-OPh	2-F-3-CF ₃
3-Cl	2-SCH ₃	4-OPh	2-F-6-CF ₃
4-Cl	3-SCH ₃	2-ethoxy	2-Cl-3-F
2-F	4-SCH ₃	3-ethoxy	2-Cl-4-F
3-F	2-OCF ₃	4-ethoxy	2-Cl-5-F
4-F	3-OCF ₃	2-isopropyl	2-Cl-6-F
2-Br	4-OCF ₃	3-isopropyl	3-Cl-2-F
3-Br	2-SCF ₃	4-isopropyl	2-Cl-4,6-diF
4-Br	3-SCF ₃	2,3-diF	2,3-diCl
2-I	4-SCF ₃	2,4-diF	2,4-diCl
3-I	2-CH ₂ CH ₃	2,5-diF	2,5-diCl
4-I	3-CH ₂ CH ₃	2,6-diF	2,6-diCl
2-OCH ₃	4-CH ₂ CH ₃	3,4-diF	3,4-diCl
3-OCH ₃	2-propyl	3,5-diF	3,5-diCl
4-OCH ₃	3-propyl	3,5-diCF ₃	2,3,6-triCl
2-CH ₃	4-propyl	2,3,6-triF	2,4,6-triCl
3-CH ₃	2-NO ₂	2,4,6-triF	2,6-diCl-4-F
4-CH ₃	3-NO ₂	3,4-diCH ₃	2,3-diF-6-CF ₃
2-CF ₃	4-NO ₂	3,5-diCH ₃	2,3,4,6-tetraF
3-CF ₃	2-Ph	2,3-diCH ₃	
4-CF ₃	3-Ph	2,4-diCH ₃	
2-CN	4-Ph	2,5-diCH ₃	

The compounds of formula (25) wherein Q is NCH₂CH₃, R¹ is 2-butynyl, R² is methyl, R³ is hydrogen, and the substituent(s) (R⁹)_p on the

5 benzene ring are those of Table 57.

TABLE 57

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₂CH₃, R¹ is 2-butynyl, R² and R³ are both methyl, and the substituent(s) (R⁹)_p on the
5 benzene ring are those of Table 58.

TABLE 58

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₂CH₃, R¹ is 2-butynyl, R² is hydrogen, R³ is methyl, and the substituent(s) (R⁹)_p on the
10 benzene ring are those of Table 59.

TABLE 59

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₂CH₃, R¹ is 2-pentynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the
15 benzene ring are those of Table 60.

TABLE 60

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₂CH₃, R¹ is 1-methyl-2-butyryl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 61.

TABLE 61

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₂CH₃, R¹ is 1-methyl-2-propyryl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 62.

TABLE 62

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₂CH₃, R¹ is 2-butyryl, R² is hydrogen, R³ is fluorine, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 63.

TABLE 63

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₂CH₂CH₃, R¹ is 2-propynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the
5 benzene ring are those of Table 64.

TABLE 64

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₂CH₂CH₃, R¹ is 2-propynyl, R² is methyl, and R³ is hydrogen, and the substituent(s) (R⁹)_p on
10 the benzene ring are those of Table 65.

TABLE 65

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₂CH₂CH₃, R¹ is 2-propynyl, R² and R³ are both methyl, and the substituent(s) (R⁹)_p on the
15 benzene ring are those of Table 66.

TABLE 66

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₂CH₂CH₃, R¹ is 2-propynyl, R² is hydrogen, R³ is methyl, and the substituent(s) (R⁹)_p on the
5 benzene ring are those of Table 67.

TABLE 67

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₂CH₂CH₃, R¹ is 2-butynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the
10 benzene ring are those of Table 68.

TABLE 68

H	3-CN	2-OPh	2,6-diCH ₃
2-Cl	4-CN	3-OPh	2-F-3-CF ₃
3-Cl	2-SCH ₃	4-OPh	2-F-6-CF ₃
4-Cl	3-SCH ₃	2-ethoxy	2-Cl-3-F
2-F	4-SCH ₃	3-ethoxy	2-Cl-4-F
3-F	2-OCF ₃	4-ethoxy	2-Cl-5-F
4-F	3-OCF ₃	2-isopropyl	2-Cl-6-F
2-Br	4-OCF ₃	3-isopropyl	3-Cl-2-F
3-Br	2-SCF ₃	4-isopropyl	2-Cl-4,6-diF
4-Br	3-SCF ₃	2,3-diF	2,3-diCl
2-I	4-SCF ₃	2,4-diF	2,4-diCl
3-I	2-CH ₂ CH ₃	2,5-diF	2,5-diCl
4-I	3-CH ₂ CH ₃	2,6-diF	2,6-diCl
2-OCH ₃	4-CH ₂ CH ₃	3,4-diF	3,4-diCl
3-OCH ₃	2-propyl	3,5-diF	3,5-diCl
4-OCH ₃	3-propyl	3,5-diCF ₃	2,3,6-triCl
2-CH ₃	4-propyl	2,3,6-triF	2,4,6-triCl
3-CH ₃	2-NO ₂	2,4,6-triF	2,6-diCl-4-F
4-CH ₃	3-NO ₂	3,4-diCH ₃	2,3-diF-6-CF ₃
2-CF ₃	4-NO ₂	3,5-diCH ₃	2,3,4,6-tetraF
3-CF ₃	2-Ph	2,3-diCH ₃	
4-CF ₃	3-Ph	2,4-diCH ₃	
2-CN	4-Ph	2,5-diCH ₃	

The compounds of formula (25) wherein Q is NCH₂CH₂CH₃, R¹ is 2-butynyl, R² is methyl, R³ is hydrogen, and the substituent(s) (R⁹)_p on the

5 benzene ring are those of Table 69.

TABLE 69

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₂CH₂CH₃, R¹ is 2-butynyl, R² and R³ are both methyl, and the substituent(s) (R⁹)_p on the
5 benzene ring are those of Table 70.

TABLE 70

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₂CH₂CH₃, R¹ is 2-butynyl, R² is hydrogen, R³ is methyl, and the substituent(s) (R⁹)_p on the
10 benzene ring are those of Table 71.

TABLE 71

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₂CH₂CH₃, R¹ is 2-pentynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the
15 benzene ring are those of Table 72.

TABLE 72

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₂CH₂CH₃, R¹ is 1-methyl-2-butynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 73.

TABLE 73

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₂CH₂CH₃, R¹ is 1-methyl-2-propynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 74.

TABLE 74

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is NCH₂CH₂CH₃, R¹ is 2-butynyl, R² is hydrogen, R³ is fluorine, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 75.

TABLE 75

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is sulfur, R¹ is 2-propynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring
 5 are those of Table 76.

TABLE 76

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is sulfur, R¹ is 2-propynyl, R² is methyl, R³ is hydrogen, and the substituent(s) (R⁹)_p on the benzene ring
 10 are those of Table 77.

TABLE 77

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is sulfur, R¹ is 2-propynyl, R² and R³ are both methyl, and the substituent(s) (R⁹)_p on the benzene ring
 15 are those of Table 78.

TABLE 78

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is sulfur, R¹ is 2-propynyl, R² is hydrogen, R³ is methyl, and the substituent(s) (R⁹)_p on the benzene ring
5 are those of Table 79.

TABLE 79

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is sulfur, R¹ is 2-butynyl,
10 R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 80.

TABLE 80

H	2-CH ₃	3-SCF ₃	2-Cl-3-F
2-Cl	3-CH ₃	4-SCF ₃	2-Cl-6-F
3-Cl	4-CH ₃	2,3-diF	2-Cl-4,6-diF
4-Cl	2-CF ₃	2,4-diF	2,3-diCl
2-F	3-CF ₃	2,5-diF	2,3-diCH ₃
3-F	4-CF ₃	2,6-diF	2,3,6-triF
4-F	2-OCF ₃	3,4-diF	2,4,6-triF
2-OCH ₃	3-OCF ₃	3,5-diF	2,3,4,6-tetraF
3-OCH ₃	4-OCF ₄	2-F-3-CF ₃	
4-OCH ₄	2-SCF ₃	2-F-6-CF ₃	

The compounds of formula (25) wherein Q is sulfur, R¹ is 2-butynyl, R² is methyl, R³ is hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 81.

TABLE 81

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is sulfur, R¹ is 2-butynyl, R² and R³ are both methyl, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 82.

TABLE 82

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is sulfur, R¹ is 2-butynyl, R² is hydrogen, R³ is methyl, and the substituent(s) (R⁹)_p on the benzene ring
 5 are those of Table 83.

TABLE 83

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is sulfur, R¹ is 2-pentynyl,
 10 R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring
 are those of Table 84.

TABLE 84

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

15 The compounds of formula (25) wherein Q is sulfur, R¹ is 1-methyl-2-butynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 85.

TABLE 85

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is sulfur, R¹ is 1-methyl-2-propynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the
 5 benzene ring are those of Table 86.

TABLE 86

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is sulfur, R¹ is 2-butyryl,
 10 R² is hydrogen, R³ is fluorine, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 87.

TABLE 87

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (25) wherein Q is CHCN, R¹ is 2-propynyl,
 15 R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 88.

TABLE 88

H	2-Cl	2,3-diF	2,6-diF
2-F	3-F	2-Cl-6-F	2,3-diCl

The compounds of formula (25) wherein Q is CHCN, R¹ is 2-butynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring
 5 are those of Table 89.

TABLE 89

H	2-Cl	2,3-diF	2,6-diF
2-F	3-F	2-Cl-6-F	2,3-diCl

The compounds of formula (25) wherein Q is carbonyl, R¹ is 2-
 10 propynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 90.

TABLE 90

H	2-Cl	2,3-diF	2,6-diF
2-F	3-F	2-Cl-6-F	2,3-diCl


15 The compounds of formula (25) wherein Q is carbonyl, R¹ is 2-butynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 91.

TABLE 91

H	2-Cl	2,3-diF	2,6-diF
2-F	3-F	2-Cl-6-F	2,3-diCl

51

The compounds of formula (26):



wherein R^1 is 2-propynyl, R^2 and R^3 are both hydrogen, and the substituent(s) $(R^9)_p$ on the benzene ring are selected from those of Table 92.

5

TABLE 92

H	2-CH ₃	3-SCF ₃	2-Cl-3-F
2-Cl	3-CH ₃	4-SCF ₃	2-Cl-6-F
3-Cl	4-CH ₃	2,3-diF	2-Cl-4,6-diF
4-Cl	2-CF ₃	2,4-diF	2,3-diCl
2-F	3-CF ₃	2,5-diF	2,3-diCH ₃
3-F	4-CF ₃	2,6-diF	2,3,6-triF
4-F	2-OCF ₃	3,4-diF	2,4,6-triF
2-OCH ₃	3-OCF ₃	3,5-diF	2,3,4,6-tetraF
3-OCH ₃	4-OCF ₄	2-F-3-CF ₃	
4-OCH ₄	2-SCF ₃	2-F-6-CF ₃	

The compounds of formula (26) wherein R^1 is 2-propynyl, R^2 is methyl, R^3 is hydrogen, and the substituent(s) $(R^9)_p$ on the benzene ring are selected

10 from those of Table 93.

TABLE 93

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (26) wherein R^1 is 2-propynyl, R^2 and R^3 are both methyl, and the substituent(s) $(R^9)_p$ on the benzene ring are selected from those of Table 94.

5

TABLE 94

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (26) wherein R^1 is 2-propynyl, R^2 is hydrogen, R^3 is methyl, and the substituent(s) $(R^9)_p$ on the benzene ring are selected from those of Table 95.

10

TABLE 95

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (26) wherein R^1 is 2-butynyl, R^2 and R^3 are both hydrogen, and the substituent(s) $(R^9)_p$ on the benzene ring are selected

15 from those of Table 96.

TABLE 96

H	3-CN	2-OPh	2,6-diCH ₃
2-Cl	4-CN	3-OPh	2-F-3-CF ₃
3-Cl	2-SCH ₃	4-OPh	2-F-6-CF ₃
4-Cl	3-SCH ₃	2-ethoxy	2-Cl-3-F
2-F	4-SCH ₃	3-ethoxy	2-Cl-4-F
3-F	2-OCF ₃	4-ethoxy	2-Cl-5-F
4-F	3-OCF ₃	2-isopropyl	2-Cl-6-F
2-Br	4-OCF ₃	3-isopropyl	3-Cl-2-F
3-Br	2-SCF ₃	4-isopropyl	2-Cl-4,6-diF
4-Br	3-SCF ₃	2,3-diF	2,3-diCl
2-I	4-SCF ₃	2,4-diF	2,4-diCl
3-I	2-CH ₂ CH ₃	2,5-diF	2,5-diCl
4-I	3-CH ₂ CH ₃	2,6-diF	2,6-diCl
2-OCH ₃	4-CH ₂ CH ₃	3,4-diF	3,4-diCl
3-OCH ₃	2-propyl	3,5-diF	3,5-diCl
4-OCH ₃	3-propyl	3,5-diCF ₃	2,3,6-triCl
2-CH ₃	4-propyl	2,3,6-triF	2,4,6-triCl
3-CH ₃	2-NO ₂	2,4,6-triF	2,6-diCl-4-F
4-CH ₃	3-NO ₂	3,4-diCH ₃	2,3-diF-6-CF ₃
2-CF ₃	4-NO ₂	3,5-diCH ₃	2,3,4,6-tetraF
3-CF ₃	2-Ph	2,3-diCH ₃	
4-CF ₃	3-Ph	2,4-diCH ₃	
2-CN	4-Ph	2,5-diCH ₃	

The compounds of formula (26) wherein R¹ is 2-butyryl, R² is methyl, R³ is hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of

5 Table 97.

TABLE 97

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (26) wherein R¹ is 2-butynyl, R² and R³ are both methyl, and the substituent(s) (R⁹)_p on the benzene ring are those of

5 Table 98.

TABLE 98

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (26) wherein R¹ is 2-butynyl, R² is

10 hydrogen, R³ is methyl, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 99.

TABLE 99

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

15 The compounds of formula (26) wherein R¹ is 2-pentynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 100.

TABLE 100

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (26) wherein R¹ is 3-butynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of

5 Table 101.

TABLE 101

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (26) wherein R¹ is 1-methyl-2-butynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 102.

10

TABLE 102

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (26) wherein R¹ is 1-methyl-2-propynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 103.

15

TABLE 103

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (26) wherein R¹ is 3-pentynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those
5 of Table 104.

TABLE 104

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (26) wherein R¹ is 3-chloro-2-propynyl, R²
10 and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 105.

TABLE 105

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (26) wherein R¹ is 4-fluoro-2-butyryl, R²
15 and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 106.

TABLE 106

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

The compounds of formula (26) wherein R¹ is 2-heptynyl, R² and R³ are both hydrogen, and the substituent(s) (R⁹)_p on the benzene ring are those
5 of Table 107.

TABLE 107

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	


The compounds of formula (26) wherein R¹ is 2-butyryl, R² is
10 hydrogen, R³ is fluorine, and the substituent(s) (R⁹)_p on the benzene ring are those of Table 108.

TABLE 108

H	3-Cl	2,3-diCH ₃	2,6-diF
2-Cl	3-Br	2-Cl-6-F	2,3-diF
2-Br	3-F	2-Cl-3-F	2-Cl-3,6-diF
2-F	2,3-diCl	3-Cl-2-F	

58

The compounds of formula (27):



(27)

wherein R¹ is 2-propynyl, R², R³, and R⁷ are all hydrogen, and R⁶ is selected from those of Table 109.

5

TABLE 109

H	CH ₃	1-CH ₃ -2-propenyl	2-CH ₃ -2-propenyl
CH ₂ CH ₃	(CH ₂) ₂ CH ₃	3-Cl-2-propenyl	2-Cl-2-propenyl
CH(CH ₃) ₂	(CH ₂) ₃ CH ₃	3,3-diCl-2-propenyl	3,3-diF-2-propenyl
CH ₂ CH(CH ₃) ₂	C(CH ₃) ₃	2-propynyl	2-butyryl
allyl	2-butenyl	2-pentyryl	2,2,2-trifluoroethyl

The compounds of formula (27) wherein R¹ is 2-butyryl, R², R³, and R⁷ are all hydrogen, and R⁶ is selected from those of Table 110.

10

TABLE 110

H	CH ₃	1-CH ₃ -2-propenyl	2-CH ₃ -2-propenyl
CH ₂ CH ₃	(CH ₂) ₂ CH ₃	3-Cl-2-propenyl	2-Cl-2-propenyl
CH(CH ₃) ₂	(CH ₂) ₃ CH ₃	3,3-diCl-2-propenyl	3,3-diF-2-propenyl
CH ₂ CH(CH ₃) ₂	C(CH ₃) ₃	2-propynyl	2-butyryl
allyl	2-butenyl	2-pentyryl	2,2,2-trifluoroethyl

The compounds of formula (27) wherein R¹ is 2-pentyryl, R², R³, and R⁷ are all hydrogen, and R⁶ is selected from those of Table 111.

15

TABLE 111

H	CH ₃	1-CH ₃ -2-propenyl	2-CH ₃ -2-propenyl
CH ₂ CH ₃	(CH ₂) ₂ CH ₃	3-Cl-2-propenyl	2-Cl-2-propenyl
CH(CH ₃) ₂	(CH ₂) ₃ CH ₃	3,3-diCl-2-propenyl	3,3-diF-2-propenyl
CH ₂ CH(CH ₃) ₂	C(CH ₃) ₃	2-propynyl	2-butylnyl
allyl	2-butenyl	2-pentynyl	2,2,2-trifluoroethyl

The compounds of formula (27) wherein R¹ is 2-propynyl, R² and R³ are both hydrogen, R⁷ is methyl, and R⁶ is selected from those of Table 112.

5

TABLE 112

CH ₂ CH ₃	CH ₃	1-CH ₃ -2-propenyl	2-CH ₃ -2-propenyl
CH(CH ₃) ₂	(CH ₂) ₂ CH ₃	3-Cl-2-propenyl	2-Cl-2-propenyl
CH ₂ CH(CH ₃) ₂	(CH ₂) ₃ CH ₃	3,3-diCl-2-propenyl	3,3-diF-2-propenyl
allyl	C(CH ₃) ₃	2-propynyl	2-butylnyl
2-butenyl	2-pentynyl	2,2,2-trifluoroethyl	

The compounds of formula (27) wherein R¹ is 2-butylnyl, R² and R³ are both hydrogen, R⁷ is methyl, and R⁶ is selected from those of Table 113.

10

TABLE 113

CH ₂ CH ₃	CH ₃	1-CH ₃ -2-propenyl	2-CH ₃ -2-propenyl
CH(CH ₃) ₂	(CH ₂) ₂ CH ₃	3-Cl-2-propenyl	2-Cl-2-propenyl
CH ₂ CH(CH ₃) ₂	(CH ₂) ₃ CH ₃	3,3-diCl-2-propenyl	3,3-diF-2-propenyl
allyl	C(CH ₃) ₃	2-propynyl	2-butylnyl
2-butenyl	2-pentynyl	2,2,2-trifluoroethyl	

The compounds of formula (27) wherein R¹ is 2-pentynyl, R² and R³ are both hydrogen, R⁷ is methyl, and R⁶ is selected from those of Table 114.

15

TABLE 114

CH_2CH_3	CH_3	1- CH_3 -2-propenyl	2- CH_3 -2-propenyl
$\text{CH}(\text{CH}_3)_2$	$(\text{CH}_2)_2\text{CH}_3$	3-Cl-2-propenyl	2-Cl-2-propenyl
$\text{CH}_2\text{CH}(\text{CH}_3)_2$	$(\text{CH}_2)_3\text{CH}_3$	3,3-diCl-2-propenyl	3,3-diF-2-propenyl
allyl	$\text{C}(\text{CH}_3)_3$	2-propynyl	2-butynyl
2-butenyl	2-pentynyl	2,2,2-trifluoroethyl	

The compounds of formula (27) wherein R^1 is 2-propynyl, R^2 and R^3 are both hydrogen, R^7 is ethyl, and R^6 is selected from those of Table 115.

5

TABLE 115

CH_2CH_3	$(\text{CH}_2)_2\text{CH}_3$	1- CH_3 -2-propenyl	2- CH_3 -2-propenyl
$\text{CH}(\text{CH}_3)_2$	$(\text{CH}_2)_3\text{CH}_3$	3-Cl-2-propenyl	2-Cl-2-propenyl
$\text{CH}_2\text{CH}(\text{CH}_3)_2$	$\text{C}(\text{CH}_3)_3$	3,3-diCl-2-propenyl	3,3-diF-2-propenyl
allyl	2-pentynyl	2-propynyl	2,2,2-trifluoroethyl
2-butenyl	2-butynyl		

The compounds of formula (27) wherein R^1 is 2-butynyl, R^2 and R^3 are both hydrogen, R^7 is ethyl, and R^6 is selected from those of Table 116.

10

TABLE 116

CH_2CH_3	$(\text{CH}_2)_2\text{CH}_3$	1- CH_3 -2-propenyl	2- CH_3 -2-propenyl
$\text{CH}(\text{CH}_3)_2$	$(\text{CH}_2)_3\text{CH}_3$	3-Cl-2-propenyl	2-Cl-2-propenyl
$\text{CH}_2\text{CH}(\text{CH}_3)_2$	$\text{C}(\text{CH}_3)_3$	3,3-diCl-2-propenyl	3,3-diF-2-propenyl
allyl	2-pentynyl	2-propynyl	2,2,2-trifluoroethyl
2-butenyl	2-butynyl		

The compounds of formula (27) wherein R^1 is 2-pentynyl, R^2 and R^3 are both hydrogen, R^7 is ethyl, and R^6 is selected from those of Table 117.

15

TABLE 117

CH ₂ CH ₃	(CH ₂) ₂ CH ₃	1-CH ₃ -2-propenyl	2-CH ₃ -2-propenyl
CH(CH ₃) ₂	(CH ₂) ₃ CH ₃	3-Cl-2-propenyl	2-Cl-2-propenyl
CH ₂ CH(CH ₃) ₂	C(CH ₃) ₃	3,3-diCl-2-propenyl	3,3-diF-2-propenyl
allyl	2-pentynyl	2-propynyl	2,2,2-trifluoroethyl
2-butenyl	2-butynyl		

The compounds of formula (27) wherein R¹ is 2-pentynyl, R² and R³ are both hydrogen, R⁷ is propyl, and R⁶ is selected from those of Table 118.

5

TABLE 118

CH(CH ₃) ₂	(CH ₂) ₂ CH ₃	1-CH ₃ -2-propenyl	2-CH ₃ -2-propenyl
CH ₂ CH(CH ₃) ₂	(CH ₂) ₃ CH ₃	3-Cl-2-propenyl	2-Cl-2-propenyl
allyl	C(CH ₃) ₃	3,3-diCl-2-propenyl	3,3-diF-2-propenyl
2-butenyl	2-pentynyl	2-propynyl	2,2,2-trifluoroethyl
2-butynyl			

The compounds of formula (27) wherein R¹ is 2-butynyl, R² and R³ are both hydrogen, R⁷ is propyl, and R⁶ is selected from those of Table 119.

10

TABLE 119

CH(CH ₃) ₂	(CH ₂) ₂ CH ₃	1-CH ₃ -2-propenyl	2-CH ₃ -2-propenyl
CH ₂ CH(CH ₃) ₂	(CH ₂) ₃ CH ₃	3-Cl-2-propenyl	2-Cl-2-propenyl
allyl	C(CH ₃) ₃	3,3-diCl-2-propenyl	3,3-diF-2-propenyl
2-butenyl	2-pentynyl	2-propynyl	2,2,2-trifluoroethyl
2-butynyl			

The compounds of formula (27) wherein R¹ is 2-pentynyl, R² and R³ are both hydrogen, R⁷ is propyl, and R⁶ is selected from those of Table 120.

15

TABLE 120

$\text{CH}(\text{CH}_3)_2$	$(\text{CH}_2)_2\text{CH}_3$	1- CH_3 -2-propenyl	2- CH_3 -2-propenyl
$\text{CH}_2\text{CH}(\text{CH}_3)_2$	$(\text{CH}_2)_3\text{CH}_3$	3-Cl-2-propenyl	2-Cl-2-propenyl
allyl	$\text{C}(\text{CH}_3)_3$	3,3-diCl-2-propenyl	3,3-diF-2-propenyl
2-butenyl	2-pentynyl	2-propynyl	2,2,2-trifluoroethyl
2-butyryl			

The compounds of formula (27) wherein R^1 is 2-propynyl, R^2 and R^3 are both hydrogen, R^7 is propyl, and R^6 is selected from those of Table 121.

5

TABLE 121

$\text{CH}_2\text{CH}(\text{CH}_3)_2$	$(\text{CH}_2)_2\text{CH}_3$	1- CH_3 -2-propenyl	2- CH_3 -2-propenyl
allyl	$(\text{CH}_2)_3\text{CH}_3$	3-Cl-2-propenyl	2-Cl-2-propenyl
2-butenyl	$\text{C}(\text{CH}_3)_3$	3,3-diCl-2-propenyl	3,3-diF-2-propenyl
2-butyryl	2-pentynyl	2-propynyl	2,2,2-trifluoroethyl

The compounds of formula (27) wherein R^1 is 2-butyryl, R^2 and R^3 are both hydrogen, R^7 is 1-methylethyl, and R^6 is selected from those of Table

10 122.

TABLE 122

$\text{CH}_2\text{CH}(\text{CH}_3)_2$	$(\text{CH}_2)_2\text{CH}_3$	1- CH_3 -2-propenyl	2- CH_3 -2-propenyl
allyl	$(\text{CH}_2)_3\text{CH}_3$	3-Cl-2-propenyl	2-Cl-2-propenyl
2-butenyl	$\text{C}(\text{CH}_3)_3$	3,3-diCl-2-propenyl	3,3-diF-2-propenyl
2-butyryl	2-pentynyl	2-propynyl	2,2,2-trifluoroethyl

The compounds of formula (27) wherein R^1 is 2-pentynyl, R^2 and R^3 are both hydrogen, R^7 is 1-methylethyl, and R^6 is selected from those of Table

15 123.

TABLE 123

$\text{CH}_2\text{CH}(\text{CH}_3)_2$	$(\text{CH}_2)_2\text{CH}_3$	1- CH_3 -2-propenyl	2- CH_3 -2-propenyl
allyl	$(\text{CH}_2)_3\text{CH}_3$	3-Cl-2-propenyl	2-Cl-2-propenyl
2-butenyl	$\text{C}(\text{CH}_3)_3$	3,3-diCl-2-propenyl	3,3-diF-2-propenyl
2-butyngyl	2-pentyngyl	2-propynyl	2,2,2-trifluoroethyl

The compounds of formula (27) wherein R^1 is 2-propynyl, R^2 and R^3 are both hydrogen, R^7 is 2,2,2-trifluoroethyl, and R^6 is selected from those of

5 Table 124.

TABLE 124

$\text{CH}_2\text{CH}(\text{CH}_3)_2$	$(\text{CH}_2)_2\text{CH}_3$	1- CH_3 -2-propenyl	2- CH_3 -2-propenyl
allyl	$(\text{CH}_2)_3\text{CH}_3$	3-Cl-2-propenyl	2-Cl-2-propenyl
2-butenyl	$\text{C}(\text{CH}_3)_3$	3,3-diCl-2-propenyl	3,3-diF-2-propenyl
2-butyngyl	2-pentyngyl	2-propynyl	

The compounds of formula (27) wherein R^1 is 2-butyngyl, R^2 and R^3 are

10 both hydrogen, R^7 is 2,2,2-trifluoroethyl, and R^6 is selected from those of

Table 125.

TABLE 125

$\text{CH}_2\text{CH}(\text{CH}_3)_2$	$(\text{CH}_2)_2\text{CH}_3$	1- CH_3 -2-propenyl	2- CH_3 -2-propenyl
allyl	$(\text{CH}_2)_3\text{CH}_3$	3-Cl-2-propenyl	2-Cl-2-propenyl
2-butenyl	$\text{C}(\text{CH}_3)_3$	3,3-diCl-2-propenyl	3,3-diF-2-propenyl
2-butyngyl	2-pentyngyl	2-propynyl	


The compounds of formula (27) wherein R^1 is 2-pentyngyl, R^2 and R^3 are both hydrogen, R^7 is 2,2,2-trifluoroethyl, and R^6 is selected from those of

15 Table 126.

TABLE 126

$\text{CH}_2\text{CH}(\text{CH}_3)_2$	$(\text{CH}_2)_2\text{CH}_3$	1- CH_3 -2-propenyl	2- CH_3 -2-propenyl
allyl	$(\text{CH}_2)_3\text{CH}_3$	3-Cl-2-propenyl	2-Cl-2-propenyl
2-butenyl	$\text{C}(\text{CH}_3)_3$	3,3-diCl-2-propenyl	3,3-diF-2-propenyl
2-butyryl	2-pentyryl	2-propynyl	

The compounds of formula (28):



- 5 wherein R^1 is 2-propynyl, R^2 and R^3 are both hydrogen, and the substituent(s) $(\text{R}^9)_p$ on the benzene ring are selected from those of Table 127.

TABLE 127

H	2-Cl	2,3-diF	2,6-diF
2-F	3-F	2-Cl-6-F	2,3-diCl


- 10 The compounds of formula (28) wherein R^1 is 2-butyryl, R^2 and R^3 are both hydrogen, and the substituent(s) $(\text{R}^9)_p$ on the benzene ring are selected from those of Table 128.

TABLE 128

H	2-Cl	2,3-diF	2,6-diF
2-F	3-F	2-Cl-6-F	2,3-diCl

65

The compounds of formula (29):



(29)


wherein R^1 is 2-butynyl, R^2 and R^3 are both hydrogen, and R^{10} is selected from those of Table 129.

5

TABLE 129

F	OCH_3	OC_2H_5
---	---------	-----------

The compounds of formula (30):



(30)


10 wherein R^1 is 2-butynyl, R^2 and R^3 are both hydrogen, and R^{11} is selected from those of Table 130.

TABLE 130

CH_3	C_2H_5	$CH(CH_3)_2$
--------	----------	--------------

66

The compounds of formula (31):



(31)

wherein R^2 and R^3 are both hydrogen and R^1 and R^{13} make a combination as defined in Table 131.

5

TABLE 131


R^1	R^{13}
2-propynyl	2-propynyl
2-propynyl	2-butynyl
2-propynyl	2-pentynyl
2-butynyl	2-butynyl
2-propynyl	3-butynyl
2-butynyl	2-pentynyl
2-propynyl	1-methyl-2-propynyl
2-butynyl	1-methyl-2-propynyl
2-butynyl	4,4-dimethyl-2-pentynyl
4,4-dimethyl-2-pentynyl	4,4-dimethyl-2-pentynyl

Production Process 1

A production process for the present compounds wherein R^4 is C_3 - C_7 alkynyloxy optionally substituted with halogen or a group of formula $-A^{1-1}R^5$ wherein A^{1-1} is oxygen or sulfur; and R^5 is optionally substituted phenyl or optionally substituted C_7 - C_9 aralkyl.

The present compounds of formula (4) can be produced from the 4,6-dichloropyrimidine compounds of formula (2) through step (1-1) and step (1-2) according to the following scheme.

67



wherein R¹ is C₃-C₇ alkynyl optionally substituted with halogen; R² and R³ are the same or different and are independently hydrogen, halogen or C₁-C₄ alkyl; and R⁴⁻¹ is C₃-C₇ alkynyloxy optionally substituted with halogen or a group of formula A¹⁻¹R⁵⁻¹ wherein A¹⁻¹ is oxygen or sulfur; and R⁵⁻¹ is optionally substituted phenyl or optionally substituted C₇-C₉ aralkyl.

Step (1-1)

The compounds of formula (3) can be produced by reacting the 4,6-dichloropyrimidine compounds of formula (2) with the compounds of formula R¹OH wherein R¹ is as defined above in the presence of a base.

The reaction is usually carried out in a solvent. The solvent used in the reaction may include ethers such as tetrahydrofuran, diethyl ether, and methyl t-butyl ether; acid amides such as N,N-dimethylformamide; dimethylsulfoxide; and mixtures thereof.

The base used in the reaction may include inorganic bases such as sodium hydride. The amount of the base used in the reaction is usually in the range of 1 to 1.5 moles, relative to 1 mole of the 4,6-dichloropyrimidine compound of formula (2).

The amount of the alcohol compound of formula R¹OH used in the reaction is usually in the range of 1 to 1.2 moles, relative to 1 mole of the 4,6-dichloropyrimidine compound of formula (2).

The reaction temperature is usually in the range of 0°C to 80°C.

The reaction time is usually in the range of 0.1 to 12 hours.

After completion of the reaction, the reaction mixture is subjected to

the ordinary post-treatment including extraction with an organic solvent and concentration for isolation of the compounds of formula (3). The compounds of formula (3) thus isolated may be purified by chromatography or other techniques.

5 Step (1-2)

The present compounds of formula (4) can be produced by reacting the compounds of formula (3) with the compounds of formula $R^{4-1}H$ wherein R^{4-1} is as defined above in the presence of a base.

10 The reaction is usually carried out in a solvent. The solvent used in the reaction may include ethers such as tetrahydrofuran, diethyl ether, and methyl t-butyl ether; acid amides such as N,N-dimethylformamide; dimethylsulfoxide; acetonitrile; and mixtures thereof.

15 The base used in the reaction may include inorganic bases such as sodium hydride and potassium carbonate; tertiary amines such as triethylamine and diisopropylethylamine; and nitrogen-containing aromatic compounds such as pyridine. The amount of the base used in the reaction is usually in the range of 1 to 1.5 moles, relative to 1 mole of the compound of formula (3).

20 The amount of the compound of formula $R^{4-1}H$ used in the reaction is usually in the range of 1 to 1.2 moles, relative to 1 mole of the compound of formula (3).

The reaction temperature is usually in the range of 0°C to 80°C.


The reaction time is usually in the range of 0.1 to 12 hours.

25 After completion of the reaction, the reaction mixture is subjected to the ordinary post-treatment including extraction with an organic solvent and concentration for isolation of the compounds of formula (4). The compounds of formula (4) thus isolated may be purified by chromatography or other techniques.

Production Process 2

A production process for the present compounds wherein R^4 is a group of formula $-A^1R^5$; A^1 is oxygen; and R^5 is optionally substituted C_7 - C_9 aralkyl.

- 5 The present compounds of formula (8) can be produced from the 4,6-dichloropyrimidine compounds of formula (2) through step (2-1) to step (2-4) according to the following scheme.



- wherein R^1 , R^2 , and R^3 are as defined above and R^{4-2} is a group of formula $-OR^{5-2}$ wherein R^{5-2} is optionally substituted C_7 - C_9 aralkyl.
- 10

Step (2-1)

The compounds of formula (5) can be produced by reacting the compounds of formula (2) with sodium thiomethoxide.

- The reaction is usually carried out in a solvent. The solvent used in the reaction may include ethers such as tetrahydrofuran, diethyl ether, and methyl t-butyl ether; acid amides such as N,N-dimethylformamide; nitriles such as acetonitrile; alcohols such as methanol and ethanol; dimethylsulfoxide; acetonitrile; and mixtures thereof.
- 15

The amount of methanethiol sodium salt used in the reaction is usually in the range of 1 to 1.5 moles, relative to 1 mole of the compound of formula (2).

The reaction temperature is usually in the range of 0°C to 80°C.

5 The reaction time is usually in the range of 1 to 12 hours.

After completion of the reaction, the reaction mixture is subjected to the ordinary post-treatment including extraction with an organic solvent and concentration for isolation of the compounds of formula (5). The compounds of formula (5) thus isolated may be purified by chromatography or other
10 techniques.

Step (2-2)

The compounds of formula (6) can be produced by reacting the compounds of formula (5) with the compounds of formula R^1OH wherein R^1 is as defined above in the presence of a base.

15 The reaction is usually carried out in a solvent. The solvent used in the reaction may include ethers such as tetrahydrofuran, diethyl ether, and methyl t-butyl ether; acid amides such as N,N-dimethylformamide; dimethylsulfoxide; and mixtures thereof.

The base used in the reaction may include inorganic bases such as
20 sodium hydride. The amount of the base used in the reaction is usually in the range of 1 to 1.5 moles, relative to 1 mole of the compound of formula (5).

The amount of the alcohol compound of formula R^1OH used in the reaction is usually in the range of 1 to 1.2 moles, relative to 1 mole of the compound of formula (5).

25 The reaction temperature is usually in the range of 0°C to 80°C.

The reaction time is usually in the range of 1 to 12 hours.

After completion of the reaction, the reaction mixture is subjected to the ordinary post-treatment including extraction with an organic solvent and

concentration for isolation of the compounds of formula (6). The compounds of formula (6) thus isolated may be purified by chromatography or other techniques.

Step (2-3)

5 The compounds of formula (7) can be produced by oxidation reaction of the compounds of formula (6).

 The oxidation reaction is usually carried out in a solvent. The solvent used in the reaction may include halogenated hydrocarbons such as chloroform and methylene chloride; nitriles such as acetonitrile; aromatic
10 hydrocarbons such as benzene and toluene; and mixtures thereof.

 The oxidizing agent used in the reaction may include peracids such as 3-chloroperbenzoic acid and peracetic acid. The amount of the oxidizing agent is usually in the range of 2 to 2.5 moles, relative to 1 mole of the compound of formula (6).

15 The reaction temperature is usually in the range of 0°C to 80°C.

 The reaction time is usually in the range of 1 to 12 hours.

 After completion of the reaction, the reaction mixture is treated with a reducing agent such as an aqueous sodium thiosulfate solution and then subjected to the ordinary post-treatment including extraction with an organic
20 solvent and concentration for isolation of the compounds of formula (7). The compounds of formula (7) thus isolated may be purified by chromatography or other techniques.

Step (2-4)

 The compounds of formula (8) can be produced by reacting the compounds of formula (7) with the compounds of formula $R^{5-2}OH$ wherein R^{5-2} is
25 as defined above in the presence of a base.

 The reaction is usually carried out in a solvent. The solvent used in the reaction may include ethers such as tetrahydrofuran, diethyl ether, and

methyl t-butyl ether; acid amides such as N,N-dimethylformamide; nitriles such as acetonitrile; dimethylsulfoxide; and mixtures thereof.

The base used in the reaction may include inorganic bases such as sodium carbonate and potassium carbonate. The amount of the base used
5 in the reaction is usually in the range of 1 to 1.5 moles, relative to 1 mole of the compound of formula (7).

The amount of the compound of formula $R^{5,2}OH$ used in the reaction is usually in the range of 1 to 1.2 moles, relative to 1 mole of the compound of formula (7).

10 The reaction temperature is usually in the range of 0°C to 80°C.


The reaction time is usually in the range of 1 to 12 hours.

After completion of the reaction, the reaction mixture is subjected to the ordinary post-treatment including extraction with an organic solvent and concentration for isolation of the compounds of formula (8). The compounds
15 of formula (8) thus isolated may be purified by chromatography or other techniques.

Production Process 3

A production process for the present compounds wherein R^4 is a group of formula A^1R^5 ; A^1 is oxygen; and R^5 is optionally substituted C_7-C_9
20 aralkyl, or R^4 is C_3-C_8 cycloalkoxy optionally substituted with halogen, hydroxy, C_1-C_4 alkyl or C_1-C_4 alkoxy.

The present compounds of formula (10) can be produced from the 4,6-dichloropyrimidine compounds of formula (2) through step (3-1) and step (3-2) according to the following scheme.



wherein R^1 , R^2 , and R^3 are as defined above and R^{4-3} is a group of formula A^1R^{5-3} ; A^1 is oxygen; and R^{5-3} is optionally substituted C_7-C_9 aralkyl, or R^{4-3} is C_3-C_8 cycloalkoxy optionally substituted with halogen, hydroxy, C_1-C_4 alkyl or C_1-C_4 alkoxy.

Step (3-1)

The compounds of formula (9) can be produced by reacting the 4,6-dichloropyrimidine compounds of formula (2) with the alcohol compounds of formula $R^{4-3}H$ wherein R^{4-3} is as defined above in the presence of a base.

The reaction is usually carried out in a solvent. The solvent used in the reaction may include ethers such as tetrahydrofuran, diethyl ether, and methyl t-butyl ether; acid amides such as N,N-dimethylformamide; dimethylsulfoxide; and mixtures thereof.

The base used in the reaction may include inorganic bases such as sodium hydride. The amount of the base used in the reaction is usually in the range of 1 to 1.5 moles, relative to 1 mole of the 4,6-dichloropyrimidine compound of formula (2).

The amount of the alcohol compound of formula $R^{4-3}H$ used in the reaction is usually in the range of 1 to 1.2 moles, relative to 1 mole of the 4,6-dichloropyrimidine compound of formula (2).

The reaction temperature is usually in the range of 0°C to 80°C .

The reaction time is usually in the range of 1 to 12 hours.

After completion of the reaction, the reaction mixture is subjected to the ordinary post-treatment including extraction with an organic solvent and

concentration for isolation of the compounds of formula (9). The compounds of formula (9) thus isolated may be purified by chromatography or other techniques.

Step (3-2)

5 The compounds of formula (10) can be produced by reacting the compounds of formula (9) with the alcohol compounds of formula R^1OH wherein R^1 is as defined above in the presence of a base.

 The reaction is usually carried out in a solvent. The solvent used in the reaction may include ethers such as tetrahydrofuran, diethyl ether, and
10 methyl t-butyl ether; acid amides such as N,N-dimethylformamide; dimethylsulfoxide; and mixtures thereof.

 The base used in the reaction may include inorganic bases such as sodium hydride. The amount of the base used in the reaction is usually in the range of 1 to 1.5 moles, relative to 1 mole of the compound of formula (9).

15 The amount of the alcohol compound of formula R^1OH used in the reaction is usually in the range of 1 to 1.2 moles, relative to 1 mole of the compound of formula (9).

 The reaction temperature is usually in the range of 0°C to 80°C.

 The reaction time is usually in the range of 1 to 12 hours.


20 After completion of the reaction, the reaction mixture is subjected to the ordinary post-treatment including extraction with an organic solvent and concentration for isolation of the compounds of formula (10). The compounds of formula (10) thus isolated may be purified by chromatography or other techniques.

25 Production Process 4

 A production process for the present compounds wherein R^4 is optionally substituted phenyl.

 The present compounds of formula (12) can be produced through step

(41-) and step (4-2) according to the following scheme.



wherein R¹, R², and R³ are as defined above and R⁴⁻⁴ is optionally substituted phenyl.

5 Step (4-1)

The compounds of formula (11) can be produced by reacting the 4,6-dichloropyrimidine compounds of formula (2) with the phenylboronic acid compounds of formula R⁴⁻⁴B(OH)₂ wherein R⁴⁻⁴ is as defined above in the presence of a transition metal compound under an atmosphere of a gas inert to the reaction, such as argon.

The reaction is usually carried out in a solvent. The solvent used in the reaction may include alcohols such as methanol, ethanol, and 2-propanol; ethers such as 1,4-dioxane, tetrahydrofuran, 1,2-dimethoxyethane, and methyl t-butyl ether; aliphatic hydrocarbons such as hexane and heptane; acid amides such as N,N-dimethylformamide; water; and mixtures thereof.

The transition metal compound used in the reaction may include palladium compounds, specific examples of which are palladium acetate, tetrakis(triphenylphosphine)palladium, {1,1'-bis(diphenylphosphino)ferrocene}-dichloropalladium(II) methylene chloride complex, and bis(triphenylphosphine)palladium (II) chloride. The amount of the transition metal compound used in the reaction, although it may be altered within the range to attain the purpose, is usually in the range of 0.01 to 0.1 mole, relative to 1 mole of the 4,6-dichloropyrimidine compound of formula (2).

The amount of the phenylboronic acid of formula R⁴⁻⁴B(OH)₂ used in

the reaction is usually in the range of 0.9 to 1.2 moles, relative to 1 mole of the 4,6-dichlorophenylpyrimidine compound of formula (2).

The reaction temperature is usually in the range of 0°C to 80°C.

The reaction may also be carried out in the presence of a base and a
5 phase transfer catalyst, if necessary. The base which can be used in the reaction may include inorganic bases such as barium hydroxide, potassium carbonate, sodium hydrogencarbonate, and tripotassium phosphate; and alkali metal salts such as sodium acetate and potassium acetate. The phase transfer catalyst may include quaternary ammonium salts such as tetra-
10 butylammonium bromide and benzyltrimethyl ammonium bromide.

After completion of the reaction, the reaction mixture is subjected to the ordinary post-treatment including extraction with an organic solvent and concentration for isolation of the compounds of formula (11). The compounds of formula (11) thus isolated may be purified by chromatography or
15 other techniques.

The phenylboronic acid compounds of formula $R^{4,4}B(OH)_2$ can be produced, for example, by reacting Grignard compounds such as $R^{4,4}MgBr$ or organic lithium compounds such as $R^{4,4}Li$ with boronic acid esters such as trimethoxyborane or triethoxyborane. The Grignard compounds of formula
20 $R^{4,4}MgBr$ can be produced by reacting the corresponding halides, i.e., $R^{4,4}Br$ with magnesium. The organic lithium compounds of formula $R^{4,4}Li$ can be produced by reacting $R^{4,4}Br$ with n-butyl lithium.

Step (4-2)

The compounds of formula (12) can be produced by reacting the com-
25 pounds of formula (11) with the alcohol compounds of formula R^1OH wherein R^1 is as defined above in the presence of a base.

The reaction is usually carried out in a solvent. The solvent used in the reaction may include ethers such as tetrahydrofuran, diethyl ether, and

methyl t-butyl ether; acid amides such as N,N-dimethylformamide; nitriles such as acetonitrile; dimethylsulfoxide; and mixtures thereof.

The base used in the reaction may include inorganic bases such as sodium hydride. The amount of the base used in the reaction is usually in the range of 1 to 1.5 moles, relative to 1 mole of the compound of formula (11).

The amount of the alcohol compound of formula R^1OH used in the reaction is usually in the range of 1 to 1.2 moles, relative to 1 mole of the compound of formula (11).

The reaction temperature is usually in the range of $0^{\circ}C$ to $80^{\circ}C$.

The reaction time is usually in the range of 1 to 12 hours.


After completion of the reaction, the reaction mixture is subjected to the ordinary post-treatment including extraction with an organic solvent and concentration for isolation of the compounds of formula (12). The compounds of formula (12) thus isolated may be purified by chromatography or other techniques.

Production Process 5

A production process for the present compounds wherein R^4 is a group of formula NR^6R^7 ; R^6 is C_1-C_7 alkyl, C_1-C_3 haloalkyl, C_2-C_4 (alkoxymethyl), C_2-C_4 (haloalkoxymethyl), C_3-C_6 alkenyl, C_3-C_6 haloalkenyl, C_3-C_7 alkynyl, cyanomethyl, optionally substituted phenyl, or optionally substituted C_7-C_9 aralkyl; and R^7 is hydrogen, C_1-C_7 alkyl, C_1-C_3 haloalkyl, C_2-C_4 (alkoxymethyl), C_2-C_4 (haloalkoxymethyl), C_3-C_6 alkenyl, C_3-C_6 haloalkenyl, C_3-C_7 alkynyl, cyanomethyl, optionally substituted phenyl, or optionally substituted C_7-C_9 aralkyl.

The compounds of formula (15) can be produced from the 4,6-dichloropyrimidine compounds of formula (2) through step (5-1) and step (5-2) or through step (5-3) and step (5-4) according to the following scheme.

78



wherein R^1 , R^2 , and R^3 are as defined above; R^{6-5} is C_1 - C_7 alkyl, C_1 - C_8 haloalkyl, C_2 - C_4 (alkoxymethyl), C_2 - C_4 (haloalkoxymethyl), C_3 - C_6 alkenyl, C_3 - C_6 haloalkenyl, C_3 - C_7 alkynyl, cyanomethyl, optionally substituted phenyl, or optionally substituted C_7 - C_9 aralkyl; and R^{7-5} is hydrogen, C_1 - C_7 alkyl, C_1 - C_3 haloalkyl, C_2 - C_4 (alkoxymethyl), C_2 - C_4 (haloalkoxymethyl), C_3 - C_6 alkenyl, C_3 - C_6 haloalkenyl, C_3 - C_7 alkynyl, cyanomethyl, optionally substituted phenyl, or optionally substituted C_7 - C_9 aralkyl.

Step (5-1)

The compounds of formula (13) can be produced by reacting the 4,6-dichloropyrimidine compounds of formula (2) with the amine compounds of formula $R^{6-5}R^{7-5}NH$ wherein R^{6-5} and R^{7-5} are as defined above.

The reaction is usually carried out in the presence or absence of a base in a solvent.

The solvent used in the reaction may include ethers such as tetrahydrofuran, diethyl ether, and methyl *t*-butyl ether; acid amides such as *N,N*-dimethylformamide; alcohols such as methanol and ethanol; and mixtures thereof.

The base used in the reaction may include inorganic bases such as sodium hydride; and organic bases such as triethylamine, and the base may suitable be selected depending upon the kind of solvent used in the reaction. When a base is used in the reaction, the amount of the base used in the
5 reaction is usually in the range of 1 to 2.5 moles, relative to 1 mole of the 4,6-dichloropyrimidine compound of formula (2).

The amine of the amine compound of formula $R^{6-5}R^{7-5}NH$ used in the reaction is usually in the range of 1 to 1.2 moles, relative to 1 mole of the 4,6-dichloropyrimidine compound of formula (2).

10 The reaction temperature is usually in the range of 0°C to 80°C.

The reaction time is usually in the range of 1 to 12 hours.

After completion of the reaction, the reaction mixture is subjected to the ordinary post-treatment including extraction with an organic solvent and concentration for isolation of the compounds of formula (13). The com-
15 pounds of formula (13) thus isolated may be purified by chromatography or other techniques.

Step (5-2)

The compounds of formula (15) can be produced by reacting the compounds of formula (13) with the alcohol compounds of formula R^1OH wherein
20 R^1 is as defined above in the presence of a base.

The reaction is usually carried out in a solvent. The solvent used in the reaction may include ethers such as tetrahydrofuran, diethyl ether, and methyl t-butyl ether; acid amides such as N,N-dimethylformamide; nitriles such as acetonitrile; dimethylsulfoxide; and mixtures thereof.

25 The base used in the reaction may include inorganic bases such as sodium hydride. The amount of the base used in the reaction is usually in the range of 1 to 1.5 moles, relative to 1 mole of the compound of formula (13).

The amount of the alcohol compound of formula R^1OH used in the reaction is usually in the range of 1 to 1.2 moles, relative to 1 mole of the compound of formula (13).

The reaction temperature is usually in the range of 0°C to 80°C.

5 The reaction time is usually in the range of 1 to 12 hours.

After completion of the reaction, the reaction mixture is subjected to the ordinary post-treatment including extraction with an organic solvent and concentration for isolation of the compounds of formula (15). The compounds of formula (15) thus isolated may be purified by chromatography or
10 other techniques.

Step (5-3)

The compounds of formula (14) can be produced by reacting the 4,6-dichloropyrimidine compounds of formula (2) with the alcohol compounds of formula R^1OH wherein R^1 is as defined above in the presence of a base.

15 The reaction is usually carried out in a solvent. The solvent used in the reaction may include ethers such as tetrahydrofuran, diethyl ether, and methyl t-butyl ether; acid amides such as N,N-dimethylformamide; nitriles such as acetonitrile; dimethylsulfoxide; and mixtures thereof.

The base used in the reaction may include inorganic bases such as
20 sodium hydride. The amount of the base used in the reaction is usually in the range of 1 to 1.5 moles, relative to 1 mole of the 4,6-dichloropyrimidine compound of formula (2).

The amount of the alcohol compound of formula R^1OH used in the reaction is usually in the range of 1 to 1.2 moles, relative to 1 mole of the
25 4,6-dichloropyrimidine compound of formula (2).

The reaction temperature is usually in the range of 0°C to 80°C.

The reaction time is usually in the range of 1 to 12 hours.

After completion of the reaction, the reaction mixture is subjected to

the ordinary post-treatment including extraction with an organic solvent and concentration for isolation of the compounds of formula (14). The compounds of formula (14) thus isolated may be purified by chromatography or other techniques.

5 Step (5-4)

The compounds of formula (15) can be produced by reacting the compounds of formula (14) with the amine compounds of formula $R^{6-5}R^{7-5}NH$ wherein R^{6-5} and R^{7-5} are as defined above.

10 The reaction is usually carried out in the presence or absence of a base in a solvent.

The solvent used in the reaction may include ethers such as tetrahydrofuran, diethyl ether, and methyl t-butyl ether; acid amides such as N,N-dimethylformamide; alcohols such as methanol and ethanol; and mixtures thereof.

15 The base used in the reaction may include inorganic bases such as sodium hydride; and organic bases such as triethylamine, and the base may suitable be selected depending upon the kind of solvent used in the reaction. When a base is used in the reaction, the amount of the base used in the reaction is usually in the range of 1 to 5 moles, relative to 1 mole of the compound of formula (14).

20 The amine of the amine compound of formula $R^{6-5}R^{7-5}NH$ used in the reaction is usually in the range of 1 to 4 moles, relative to 1 mole of the compound of formula (14).

The reaction temperature is usually in the range of 0°C to 80°C.

25 The reaction time is usually in the range of 1 to 48 hours.

After completion of the reaction, the reaction mixture is treated, for example, by the following method for isolation of the desired product.

1) Method involving extraction of the reaction mixture with an

organic solvent and the subsequent concentration; or


2) Method involving direct concentration of the reaction mixture without any treatment.

The compounds of formula (15) thus isolated may be purified by
5 chromatography or other techniques.

Production Process 6

A production process for the present compounds wherein R^4 is a group of formula NR^6R^7 ; R^6 is C_1 - C_7 alkyl, C_1 - C_3 haloalkyl, C_2 - C_4 (alkoxymethyl), C_2 - C_4 (haloalkoxymethyl), C_3 - C_6 alkenyl, C_3 - C_6 haloalkenyl, C_3 - C_7
10 alkynyl, cyanomethyl, optionally substituted phenyl, or optionally substituted C_7 - C_9 aralkyl; and R^7 is C_1 - C_7 alkyl, C_1 - C_3 haloalkyl, C_2 - C_4 (alkoxymethyl), C_2 - C_4 (haloalkoxymethyl), C_3 - C_6 alkenyl, C_3 - C_6 haloalkenyl, C_3 - C_7 alkynyl, cyanomethyl, or optionally substituted C_7 - C_9 aralkyl.

The compounds of formula (19) can be produced from the 4,6-dichloropyrimidine compounds of formula (2) according to the following scheme.
15



wherein R^1 , R^2 , and R^3 are as defined above; R^{6-6} is C_1 - C_7 alkyl, C_1 - C_3 haloalkyl, C_2 - C_4 (alkoxymethyl), C_2 - C_4 (haloalkoxymethyl), C_3 - C_6 alkenyl, C_3 - C_6 haloalkenyl, C_3 - C_7 alkynyl, cyanomethyl, optionally substituted phenyl, or optionally substituted C_7 - C_9 aralkyl; and R^{7-6} is C_1 - C_7 alkyl, C_1 - C_3 haloalkyl, C_2 - C_4 (alkoxymethyl), C_2 - C_4 (haloalkoxymethyl), C_3 - C_6 alkenyl, C_3 - C_6 haloalkenyl, C_3 - C_7 alkynyl, cyanomethyl, or optionally substituted C_7 - C_9 aralkyl.

Step (6-1)

The compounds of formula (16) can be produced by reacting the 4,6-dichloropyrimidine compounds of formula (2) with the amine compounds of formula $R^{6-6}NH_2$ wherein R^{6-6} is as defined above.

The reaction is usually carried out in the presence or absence of a base in a solvent.

The solvent used in the reaction may include ethers such as tetra-

hydrofuran, diethyl ether, and methyl t-butyl ether; acid amides such as N,N-dimethylformamide; alcohols such as methanol and ethanol; and mixtures thereof.

The base used in the reaction may include inorganic bases such as sodium hydride; and organic bases such as triethylamine, and the base may
5 suitable be selected depending upon the kind of solvent used in the reaction. When a base is used in the reaction, the amount of the base used in the reaction is usually in the range of 1 to 2 moles, relative to 1 mole of the 4,6-dichloropyrimidine compound of formula (2).

10 The amine compound of formula $R^{6-6}NH_2$ used in the reaction is usually in the range of 1 to 1.2 moles, relative to 1 mole of the 4,6-dichloropyrimidine compound of formula (2).

The reaction temperature is usually in the range of 0°C to 80°C.

The reaction time is usually in the range of 1 to 12 hours.

15 After completion of the reaction, the reaction mixture is subjected to the ordinary post-treatment including extraction with an organic solvent and concentration for isolation of the compounds of formula (16). The compounds of formula (16) thus isolated may be purified by chromatography or other techniques.

20 Step (6-2)

The compounds of formula (17) can be produced by reacting the compounds of formula (16) with the compounds of formula $R^{7-6}L$ wherein R^{7-6} is as defined above and L is chlorine, bromine, iodine, methanesulfonyloxy, 4-toluenesulfonyloxy, or trifluoromethanesulfonyloxy in the presence of a base.

25 The reaction is usually carried out in a solvent. The solvent used in the reaction may include ethers such as tetrahydrofuran, diethyl ether, and methyl t-butyl ether; acid amides such as N,N-dimethylformamide; nitriles such as acetonitrile; dimethylsulfoxide; and mixtures thereof.

The base used in the reaction may include inorganic bases such as sodium hydride and potassium hydride; and tertiary amines such as triethylamine and diisopropylethylamine. The amount of the base used in the reaction is usually in the range of 1 to 1.5 moles, relative to 1 mole of the compound of formula (16).

The amount of the compound of formula $R^{7-6}L$ used in the reaction is usually in the range of 1 to 1.2 moles, relative to 1 mole of the compound of formula (16).

The reaction temperature is usually in the range of 0°C to 80°C.

The reaction time is usually in the range of 1 to 12 hours.

After completion of the reaction, the reaction mixture is subjected to the ordinary post-treatment including extraction with an organic solvent and concentration for isolation of the compounds of formula (17). The compounds of formula (17) thus isolated may be purified by chromatography or other techniques.

Step (6-3)

The compounds of formula (19) can be produced by reacting the compounds of formula (17) with the alcohol compounds of formula R^1OH wherein R^1 is as defined above in the presence of a base.

The reaction is usually carried out in a solvent. The solvent used in the reaction may include ethers such as tetrahydrofuran, diethyl ether, and methyl t-butyl ether; acid amides such as N,N-dimethylformamide; nitriles such as acetonitrile; dimethylsulfoxide; and mixtures thereof.

The base used in the reaction may include inorganic bases such as sodium hydride. The amount of the base used in the reaction is usually in the range of 1 to 1.5 moles, relative to 1 mole of the compound of formula (17).

The amount of the alcohol compound of formula R^1OH used in the

reaction is usually in the range of 1 to 1.2 moles, relative to 1 mole of the compound of formula (17).

The reaction temperature is usually in the range of 0°C to 80°C.

The reaction time is usually in the range of 1 to 12 hours.

5 After completion of the reaction, the reaction mixture is subjected to the ordinary post-treatment including extraction with an organic solvent and concentration for isolation of the compounds of formula (19). The compounds of formula (19) thus isolated may be purified by chromatography or other techniques.

10 Step (6-4)

The compounds of formula (18) can be produced by reacting the compounds of formula (16) with the alcohol compounds of formula R^1OH wherein R^1 is as defined above in the presence of a base.

15 The reaction is usually carried out in a solvent. The solvent used in the reaction may include ethers such as tetrahydrofuran, diethyl ether, and methyl t-butyl ether; acid amides such as N,N-dimethylformamide; nitriles such as acetonitrile; dimethylsulfoxide; and mixtures thereof.

20 The base used in the reaction may include inorganic bases such as sodium hydride. The amount of the base used in the reaction is usually in the range of 1 to 1.5 moles, relative to 1 mole of the compound of formula (16).

The amount of the alcohol compound of formula R^1OH used in the reaction is usually in the range of 1 to 1.2 moles, relative to 1 mole of the compound of formula (16).

25 The reaction temperature is usually in the range of 0°C to 80°C.

The reaction time is usually in the range of 1 to 12 hours.

After completion of the reaction, the reaction mixture is subjected to the ordinary post-treatment including extraction with an organic solvent and

concentration for isolation of the compounds of formula (18). The compounds of formula (18) thus isolated may be purified by chromatography or other techniques.

Step (6-5)

5 The compounds of formula (19) can be produced by reacting the compounds of formula (18) with the compounds of formula $R^{7-6}L$ wherein R^{7-6} is as defined above and L is chlorine, bromine, iodine, methanesulfonyloxy, 4-toluenesulfonyloxy, or trifluoromethanesulfonyloxy in the presence of a base.

10 The reaction is usually carried out in a solvent. The solvent used in the reaction may include ethers such as tetrahydrofuran, diethyl ether, and methyl t-butyl ether; acid amides such as N,N-dimethylformamide; nitriles such as acetonitrile; dimethylsulfoxide; and mixtures thereof.

15 The base used in the reaction may include inorganic bases such as sodium hydride and potassium hydride. The amount of the base used in the reaction is usually in the range of 1 to 1.5 moles, relative to 1 mole of the compound of formula (18).

 The amount of the compound of formula $R^{7-6}L$ used in the reaction is usually in the range of 1 to 1.2 moles, relative to 1 mole of the compound of formula (18).

20 The reaction temperature is usually in the range of 0°C to 80°C.

 The reaction time is usually in the range of 1 to 12 hours.

After completion of the reaction, the reaction mixture is subjected to the ordinary post-treatment including extraction with an organic solvent and concentration for isolation of the compounds of formula (18). The compounds of formula (18) thus isolated may be purified by chromatography or other techniques.


25

Production Process 7

A production process for the present compounds wherein R^4 is op-

tionally substituted C₇-C₉ aralkyl.

The compounds for formula (21) can be produced from the 4,6-dichloropyrimidine compounds of formula (2) through step (7-1) and step (7-2) according to the following scheme.



wherein R¹, R², and R³ are as defined above and R⁴⁻⁷ is optionally substituted C₇-C₉ aralkyl.

Step (7-1)

The compounds of formula (20) can be produced by reacting the 4,6-dichloropyrimidine compounds of formula (2) with zinc compounds of formula R⁴⁻⁷ZnX wherein X is chlorine, bromine, or iodine in the presence of a transition metal compound.

The reaction is usually carried out in a solvent. The solvent used in the reaction may include ethers such as tetrahydrofuran and diethyl ether; acid amides such as N,N-dimethylformamide; nitriles such as acetonitrile; and mixtures thereof.

The transition metal compound used in the reaction may include palladium compounds, specific examples of which are tetrakis(triphenylphosphine)palladium and bis(triphenylphosphine)palladium chloride. The amount of the transition metal compound used in the reaction is usually in the range of 0.01 to 0.1 mole, relative to 1 mole of the 4,6-dichloropyrimidine compound of formula (2).

The amount of the zinc compound of formula R⁴⁻⁷ZnX used in the reaction is usually in the range of 1 to 1.2 moles, relative to 1 mole of the

4,6-dichlorophenylpyrimidine compound of formula (2).

The reaction temperature is usually in the range of 0°C to 100°C (or the boiling point of a solvent used in the reaction, when it is 80°C or lower).

After completion of the reaction, the reaction mixture is subjected to
5 the ordinary post-treatment including extraction with an organic solvent and concentration for isolation of the compounds of formula (20). The compounds of formula (20) thus isolated may be purified by chromatography or other techniques.

The zinc compounds of formula $R^{4-7}ZnX$ are usually formed in the
10 system from the compounds of formula $R^{4-7}X$, zinc, trimethylsilane chloride, and dibromomethane, and they are used in the reaction.

Step (7-2)

The compounds of formula (21) can be produced by reacting the compounds of formula (20) with the compounds of formula R^1OH wherein R^1 is as
15 defined above in the presence of a base.

The reaction is usually carried out in a solvent. The solvent used in the reaction may include ethers such as tetrahydrofuran, diethyl ether, and methyl t-butyl ether; acid amides such as N,N-dimethylformamide; nitriles such as acetonitrile; dimethylsulfoxide; and mixtures thereof.

20 The base used in the reaction may include inorganic bases such as sodium hydride. The amount of the base used in the reaction is usually in the range of 1 to 1.5 moles, relative to 1 mole of the compound of formula (20).

The amount of the alcohol compound of formula R^1OH used in the
25 reaction is usually in the range of 1 to 1.2 moles, relative to 1 mole of the compound of formula (20).

The reaction temperature is usually in the range of 0°C to 80°C.


The reaction time is usually in the range of 0.5 to 12 hours.

After completion of the reaction, the reaction mixture is subjected to the ordinary post-treatment including extraction with an organic solvent and concentration for isolation of the compounds of formula (21). The compounds of formula (21) thus isolated may be purified by chromatography or other techniques.

Production Process 8

A production process for the present compounds wherein R^4 is a group of formula $-(C=O)-R^{5-8}$ and R^{5-8} is optionally substituted phenyl as well as for the present compounds wherein R^4 is a group of formula $-(C=NOR^{8-8})-R^{5-8}$; R^{5-8} is optionally substituted phenyl; and R^{8-8} is C_1-C_4 alkyl.

The compounds of formula (23) can be produced from the compounds of formula (3) through step (8-1) and step (8-2) according to the following scheme, and the compounds of formula (24) can further be produced through step (8-3).



wherein R^1 , R^2 , and R^3 are as defined above; R^{5-8} is optionally substituted

phenyl; and R^{8-8} is C_1-C_4 alkyl.

Step (8-1)

The compounds of formula (22) can be produced by reacting the compounds of formula (3) with the nitrile compounds of formula $R^{5-8}CH_2CN$ wherein R^{5-8} is as defined above in the presence of a base.

The reaction is usually carried out in a solvent. The solvent used in the reaction may include ethers such as tetrahydrofuran, diethyl ether, and methyl t-butyl ether.

The base used in the reaction may include inorganic bases such as sodium hydride and potassium hydride; lithium amides such as lithium diisopropylamide; alkali metal carbonates such as potassium carbonate and sodium carbonate; and alkali metal alkoxides such as potassium t-butoxide. The amount of the base used in the reaction is usually in the range of 1 to 2 moles, relative to 1 mole of the compound of formula (3).

The amount of the nitrile compound of formula $R^{5-8}CH_2CN$ used in the reaction is usually in the range of 1 to 1.2 moles, relative to 1 mole of the compound of formula (3).

The reaction temperature is usually in the range of $0^\circ C$ to $80^\circ C$.

The reaction time is usually in the range of 1 to 12 hours.

After completion of the reaction, the reaction mixture is subjected to the ordinary post-treatment including extraction with an organic solvent and concentration for isolation of the compounds of formula (22). The compounds of formula (22) thus isolated may be purified by chromatography or other techniques.

Step (8-2)

The compounds of formula (23) can be produced by oxidation reaction of the compounds of formula (22) with an oxygen gas in the presence of a base.

The oxidation reaction is usually carried out in a solvent. The solvent used in the reaction may include ethers such as tetrahydrofuran and diethyl ether.

The base used in the reaction may include alkali metal hydrides such as sodium hydride and potassium hydride; alkali metal carbonates such as potassium carbonate; and alkali metal alkoxides such as potassium t-butoxide. The amount of the base used in the reaction is usually in the range of 1 to 2 moles, relative to 1 mole of the compound of formula (22).

The reaction temperature is usually in the range of 0°C to 80°C.

The reaction time is usually in the range of 12 to 48 hours.

After completion of the reaction, the reaction mixture is subjected to the ordinary post-treatment including extraction with an organic solvent and concentration for isolation of the compounds of formula (23). The compounds of formula (23) thus isolated may be purified by chromatography or other techniques.

Step (8-3)

The compounds of formula (24) can be produced by reacting the compounds of formula (23) with the hydroxylamine compounds of formula $R^{6-8}O-NH_2$ wherein R^{6-8} is as defined above or salts thereof.

The reaction is usually carried out in a solvent or without any solvent, and if necessary, in the presence of a base. The solvent used in the reaction may include alcohols such as ethanol, ethers such as tetrahydrofuran, diethyl ether, and methyl t-butyl ether; pyridine; water; and mixtures thereof.

The base used in the reaction may include tertiary amines such as triethylamine; nitrogen-containing aromatic compounds such as pyridine; and carboxylic acid alkali metal salts such as sodium acetate. When a base is used in the reaction, the amount of the base is usually 1 mole to excess, relative to 1 mole of the compound of formula (23).

The amount of the compound of formula $R^{8-8}ONH_2$ used in the reaction is usually in the range of 1 to 1.2 moles, relative to 1 mole of the compound of formula (23).

The reaction temperature is usually in the range of 0°C to 150°C.

5 The reaction time is usually in the range of 1 to 12 hours.

After completion of the reaction, the reaction mixture is subjected to the ordinary post-treatment including extraction with an organic solvent and concentration for isolation of the compounds of formula (24). The compounds of formula (24) thus isolated may be purified by chromatography or
10 other techniques.

The pests against which the present compounds have an effect may include arthropods (*e.g.*, insects, acarines) and nemathelminthes, specific examples of which are as follows:

15 *Hemiptera:*

Delphacidae such as *Laodelphax striatellus*, *Nilaparvata lugens*, and *Sogatella furcifera*, *Deltocephalidae* such as *Nephotettix cincticeps* and *Empoasca onukii*, *Aphididae* such as *Aphis gossypii* and *Myzus persicae*, *Pentatomidae*, *Aleyrodidae* such as *Trialeurodes vaporariorum*, *Bemisia tabaci*, and *Bemisia argentifolii*, *Coccidae*, *Tingidae*, *Psyllidae*, etc.
20

Lepidoptera:

Pyralidae such as *Chilo suppressalis*, *Cnaphalocrocis medinalis*, *Ostrinia nubilalis*, and *Parapediasia teterrella*, *Noctuidae* such as *Spodoptera litura*, *Spodoptera exigua*, *Pseudaletia separata*, *Mamestra brassicae*,
25 *Agrotis ipsilon*, *Trichoplusia* spp., *Heliothis* spp., *Helicoverpa* spp., and *Earias* spp., *Pieridae* such as *Pieris rapae crucivora*, *Tortricidae* such as *Adoxophyes orana fasciata*, *Grapholita molesta*, and *Cydia pomonella*, *Carposinidae* such as *Carposina niponensis*, *Lyonetiidae* such as *Lyonetia*

clerkella, *Gracillariidae* such as *Phyllonorycter ringoniella*, *Phyllocnistidae* such as *Phyllocnistis citrella*, *Yponomeutidae* such as *Plutella xylostella*, *Gelechiidae* such as *Pectinophora gossypiella*, *Arctiidae*, *Tineidae*, etc.

Diptera:

- 5 *Calicidae* such as *Culex pipiens pallens*, *Culex tritaeniorhynchus*, and *Culex quinquefasciatus*, *Aedes* spp. such as *Aedes aegypti* and *Aedes albopictus*, *Anopheles* spp. such as *Anopheles sinensis*, *Chironomidae*, *Muscidae* such as *Musca domestica* and *Muscina stabulans*, *Calliphoridae*, *Sarcophagidae*, *Anthomyiidae*, *Cecidomyiidae* such as *Delia platura* and *Delia*
10 *antiqua*, *Tephritidae*, *Drosophilidae*, *Psychodidae*, *Tabanidae*, *Simuliidae*, *Stomoxysidae*, *Agromyzidae*, etc.

Coleoptera:

- Diabrotica* spp. such as *Diabrotica virgifera virgifera* and *Diabrotica undecimpunctata howardi*, *Scarabaeidae* such as *Anomala cuprea* and *Ano-*
15 *mala rufocuprea*, *Curculionidae* such as *Sitophilus zeamais*, *Lissorhoptrus oryzophilus*, and *Callosobruchus chienensis*, *Tenebrionidae* such as *Tenebrio molitor* and *Tribolium castaneum*, *Chrysomelidae* such as *Oulema oryzae*, *Aulacophora femoralis*, *Phyllotreta striolata*, and *Leptinotarsa decem-*
20 *lineata*, *Anobiidae*, *Epilachna* spp. such as *Epilachna vigintioctopunctata*, *Lyctidae*, *Bostrychidae*, *Cerambycidae*, *Paederus fuscipes*, etc.

Thysanoptera:

Thripidae such as *Thrips* spp., e.g., *Thrips palmi*, *Frankliniella* spp., e.g., *Frankliniella occidentalis*, and *Sciltothrips* spp., e.g., *Sciltothrips dorsalis*, and *Phlaeotheripidae*, etc.

- 25 *Hymenoptera:*

Tenthredinidae, *Formicidae*, *Vespididae*, etc.

Dictyoptera:

Blattidae, *Blattellidae*, etc.

*Orthoptera:**Acrididae, Gryllotalpidae, etc.**Aphaniptera:**Purex irritans etc.*5 *Anoplura:**Pediculus humanus capitis etc.**Isoptera:**Termitidae etc.**Acarina:*

- 10 *Tetranychidae* such as *Tetranychus urticae*, *Tetranychus kanzawai*,
Panonychus citri, *Panonychus ulmi*, and *Oligonychus*, *Eriophyidae* such as
Aculops pelekassi and *Aculus schlechtendali*, *Tarsonemidae* such as *Poly-*
phagotarsonemus latus, *Tenuipalpidae*, *Tuckerellidae*, *Ixodidae* such as
Haemaphysalis longicornis, *Haemaphysalis flava*, *Dermacentor taiwanicus*,
15 *Ixodes ovatus*, *Ixodes persulcatus*, and *Boophilus microplus*, *Acaridae* such
as *Tyrophagus putrescentiae*, *Epidermoptidae* such as *Dermatophagoides*
farinae and *Dermatophagoides ptrehyssnus*, *Cheyletidae* such as *Cheyletus*
eruditus and *Cheyletus malaccensis*, *Dermanyssus* spp., etc.

Nematodes.

- 20 *Pratylenchus coffeae*, *Pratylenchus fallax*, *Heterodera glycines*,
Globadera rostochiensis, *Meloidogyne hapla*, *Meloidogyne incognita*, etc.

When the present compounds are used as pesticides, they may be
used as such; however, they are usually used after formulation into oil sprays,
emulsifiable concentrates, flowables, granules, dusts, poison baits, micro-
25 capsules, or application forms by mixing with solid carriers, liquid carriers,
gaseous carriers, or baits, and if necessary, by addition of surfactants or
other auxiliaries and processing.

These formulations may usually contain the present compounds in

0.01% to 95% by weight.

The solid carrier used in the formulation may include fine powder or granules of clay materials such as kaolin clay, diatomaceous earth, synthetic hydrated silicon oxide, bentonite, Fubasami clay, and acid clay; various kinds
5 of talc, ceramics, and other inorganic minerals such as sericite, quartz, sulfur, activated charcoal, calcium carbonate, and hydrated silica; and chemical fertilizers such as ammonium sulfate, ammonium phosphate, ammonium nitrate, urea, and ammonium chloride.

The liquid carrier may include water; alcohols such as methanol and
10 ethanol; ketones such as acetone and methyl ethyl ketone; aromatic hydrocarbons such as benzene, toluene, xylene, ethylbenzene, and methylnaphthalene; aliphatic hydrocarbons such as hexane, cyclohexane, kerosene, and light oil; esters such as ethyl acetate and butyl acetate; nitriles such as acetonitrile and isobutyronitrile; ethers such as diisopropyl ether and dioxane;
15 ane; acid amides such as N,N-dimethylformamide and N,N-dimethylacetamide; halogenated hydrocarbons such as dichloromethane, trichloroethane, and carbon tetrachloride; dimethyl sulfoxide; and vegetable oils such as soybean oil and cottonseed oil.

The gaseous carrier or propellant may include Freon gas, butane gas,
20 LPG (liquefied petroleum gas), dimethyl ether, and carbon dioxide.

The surfactant may include alkyl sulfates, alkyl sulfonates, alkyl arylsulfonates, alkyl aryl ethers and their polyoxyethylene derivatives, polyethylene glycol ethers, polyhydric alcohol esters, and sugar alcohol derivatives.

25 The auxiliaries may include fixing agents, dispersing agents, and stabilizers, specific examples of which are casein, gelatin, polysaccharides such as starch, gum arabic, cellulose derivatives, and alginic acid; lignin derivatives, bentonite, sugars, synthetic water-soluble polymers such as

polyvinyl alcohol, polyvinyl pyrrolidone, and polyacrylic acid; PAP (isopropyl acid phosphate), BHT (2,6-di-tert-butyl-4-methylphenol), BHA (mixtures of 2-tert-butyl-4-methoxyphenol and 3-tert-butyl-4-methoxyphenol), vegetable oils, mineral oils, and fatty acids and their esters.

5 The base material for poison baits may include bait materials such as grain powder, vegetable oils, sugars, and crystalline cellulose; antioxidants such as dibutylhydroxytoluene and nordihydroguaiaretic acid; preservatives such as dehydroacetic acid; substances for preventing erroneous eating, such as red pepper powder; pest attractant flavors such as cheese flavor, onion
10 flavor, and peanut oil.

 When the present compounds are used as pesticides, their application amounts are usually 0.1 to 1000 g in amounts of the present compounds per 1000 m². For emulsifiable concentrates, wettable powders, flowables, or microcapsules, these formulations are usually applied after water dilution so
15 that the concentrations of active ingredients come to 10 to 10000 ppm, and for granules or dusts, these formulations are usually applied as such.

 These formulations or their water dilutions may be used in the foliar treatment of plants such as crop plants to be protected against pests, or may be applied to the nursery beds before planting of crop plant seedlings or to
20 the planting holes or the bases of plants at the time of planting. For the purpose of controlling pests inhabiting the soil of a cultivated land, they may be applied to the soil. In addition, resin formulations processed into a sheet, string or other shapes may be applied by directly winding around crop plants, extending in the neighborhood of crop plants, or laying on the soil surface at
25 the bases of plants.

 Furthermore, they may be used in admixture with or separately but simultaneously with other insecticides, nematocides, acaricides, bactericides, fungicides, herbicides, plant growth regulators, synergists, fertilizers, soil

conditioners and/or animal feeds.

The insecticide and/or nematocide and/or acaricide which can be used may include organophosphorus compounds such as Fenitrothion, Fenthion, Pyridaphenthion, Diazinon, Chlorpyrifos, Chlorpyrifos-methyl, Acephate, 5 Methidathion, Disulfoton, DDVP, Sulprofos, Profenofos, Cyanophos, Dioxabenzofos, Dimethoate, Phenthoate, Malathion, Trichlorfon, Azinphos-methyl, Monocrotophos, Dicrotophos, Ethion and Fosthiazate; carbamate compounds such as BPMC, Benfuracarb, Propoxur, Carbosulfan, Carbaryl, Methomyl, Ethiofencarb, Aldicarb, Oxamyl, Fenothiocarb, Thiodicarb, and Alanycarb; 10 pyrethroid compounds such as Etofenprox, Fenvalerate, Esfenvalerate, Fenpropathrin, Cypermethrin, α -Cypermethrin, Z-Cypermethrin, Permethrin, Cyhalothrin, λ -Cyhalothrin, Cyfluthrin, β -Cyfluthrin, Deltamethrin, Cycloprothrin, τ -Fluvalinate, Flucythrinate, Bifenthrin, Acrinathrin, Tralomethrin, Silafluofen, and Halfenprox; neonicotinoid compounds such as Acetamiprid, Clothianidin, Nitenpyram, Thiamethoxam, Dinotefuran, Imidacloprid, 15 and Thiacloprid; benzoylphenylurea compounds such as Chlorfluazuron, Teflubenzuron, Fulfenoxuron, and Lufenuron; benzoylhydrazide compounds such as Tebufenozide, Halofenozide, Methoxyfenozide, and Chromafenozide; thiadiazine derivatives such as Buprofezin; Nereistoxin derivatives such as 20 Cartap, Thiocyclam, and Bensultap; chlorinated hydrocarbon compounds such as Endosulfan, γ -BHC, and 1,1-bis(chlorophenyl)-2,2,2-trichloroethanol; formamidine derivatives such as Amitraz and Chlordimeform; thiourea derivatives such as Diafenthiuron; phenylpyrazole compounds; Chlorfenapyr, Pymetrozine, Spinosad, Indoxacarb, Pyridalyl, Pyriproxyfen, Fenoxycarb, 25 Diofenolan, Cyromazine, Bromopropylate, Tetradifon, Quinomethionate, Propargate, Fenbutatin oxide, Hexythiazox, Etoxazole, Chlofentezine, Pyridaben, Fenpyroximate, Tebfenpyrad, Pyrimidifen, Fenazaquin, Acequinocyl, Bifenazate, Fluacrypyrim, Spirodiclofen, Milbemectin, Avermectin, Ema-

mectin benzoate, Azadilactin [AZAD], and polynactin complexes [*e.g.*, tetra-nactin, dinactin, trinactin].

The present invention will hereinafter be further illustrated by many production examples, formulation examples, and test examples; however, the
5 present invention is not limited to these examples.

In the production examples and reference production examples, ¹H-NMR shows, unless otherwise indicated, data measured using tetramethylsilane as an internal standard in a deuterated chloroform solvent.

In the production examples, the present compound numbers stand
10 for the numbers shown in Tables 124 to 129.

First, the production examples for the present compounds will be described below.

Production Example 1

In 4 ml of tetrahydrofuran was suspended 0.27 g of sodium hydride
15 (60% in oil), to which 0.7 ml of a tetrahydrofuran solution containing 0.41 g of 2-butyne-1-ol was slowly added dropwise with stirring at room temperature. The mixture was then stirred at room temperature for 20 minutes, to which 0.7 ml of a tetrahydrofuran solution containing 0.4 g of 4,6-dichloropyrimidine was slowly added dropwise. After completion of the dropwise addition,
20 the mixture was stirred at room temperature for 40 minutes. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chroma-
25 tography to give 0.39 g of 4,6-bis(2-butyneoxy)pyrimidine (the present compound (1)), m.p.: 82.9°C.

Production Example 2

In 4 ml of tetrahydrofuran was suspended 0.12 g of sodium hydride

(60% in oil), to which 0.7 ml of a tetrahydrofuran solution containing 0.18 g of 2-butyne-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes and then cooled to 0°C, to which 0.7 ml of a tetrahydrofuran solution containing 0.4 g of 4-chloro-6-(2-propynyloxy)pyrimidine was slowly added dropwise. The mixture was further stirred at 0°C for 4.5 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.9 g of 4-(2-butynyloxy)-6-(2-propynyloxy)pyrimidine (the present compound (2)).

¹H-NMR: 1.87 (t, 3H), 2.54 (t, 1H), 4.95 (q, 2H), 5.00 (d, 2H), 6.19 (s, 1H), 8.48 (s, 1H)

15 Production Example 3

In 5 ml of tetrahydrofuran was suspended 0.34 g of sodium hydride (60% in oil), to which 0.9 ml of a tetrahydrofuran solution containing 0.62 g of 2-pentyne-1-ol was slowly added dropwise with stirring at room temperature. The mixture was then stirred at room temperature for 20 minutes, to which 0.9 ml of a tetrahydrofuran solution containing 0.5 g of 4,6-dichloropyrimidine was slowly added dropwise. After completion of the dropwise addition, the mixture was stirred at room temperature for 40 minutes. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.81 g of 4,6-bis(2-pentynyloxy)pyrimidine (the present compound (3)).

¹H-NMR: 1.15 (t, 6H), 2.16-2.35 (m, 4H), 4.97 (t, 4H), 6.18 (s, 1H), 8.46 (s, 1H)

Production Example 4

In 14 ml of tetrahydrofuran was suspended 0.57 g of sodium hydride
5 (60% in oil), to which 2.5 ml of a tetrahydrofuran solution containing 0.8 g of
2-pentyn-1-ol was slowly added dropwise with stirring at room temperature.
The mixture was stirred at room temperature for 20 minutes and then cooled
to 0°C, to which 2.5 ml of a tetrahydrofuran solution containing 1.6 g of 4-
chloro-6-(2-propynyloxy)pyrimidine was slowly added dropwise. The mix-
10 ture was further stirred at 0°C for 3.5 hours. The reaction mixture was
then poured into a saturated aqueous ammonium chloride solution and
extracted three times with chloroform. The chloroform layers were com-
bined, washed with water, dried over anhydrous magnesium sulfate, and
then concentrated. The residue was subjected to silica gel column chroma-
15 tography to give 0.98 g of 4-(2-pentynyloxy)-6-(2-propynyloxy)pyrimidine
(the present compound (5)).

¹H-NMR: 1.15 (t, 3H), 2.04-2.29 (m, 2H), 2.53 (t, 1H), 4.97-5.01 (m,
4H), 6.19 (s, 1H), 8.47 (s, 1H)

Production Example 5

20 In 2 ml of tetrahydrofuran was suspended 0.1 g of sodium hydride
(60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.15 g
of 2-pentyn-1-ol was slowly added dropwise with stirring at room tempera-
ture. The mixture was stirred at room temperature for 20 minutes and
then cooled to 0°C, to which 0.6 ml of a tetrahydrofuran solution containing
25 0.3 g of 4-chloro-6-(2-butynyloxy)pyrimidine was slowly added dropwise.
The mixture was further stirred at 0°C for 3.5 hours. The reaction mixture
was then poured into a saturated aqueous ammonium chloride solution and
extracted three times with chloroform. The chloroform layers were com-

bined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.33 g of 4-(2-butynyloxy)-6-(2-pentynyloxy)pyrimidine (the present compound (6)), m.p.: 67.4°C.

5 Production Example 6

In 2 ml of tetrahydrofuran was suspended 0.1 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.2 g of 4,4-dimethyl-2-pentyn-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes
10 and then cooled to 0°C, to which 0.6 ml of a tetrahydrofuran solution containing 0.3 g of 4-chloro-6-(2-butynyloxy)pyrimidine was slowly added dropwise. The mixture was further stirred at 0°C for 3.5 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were
15 combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.15 g of 4-(2-butynyloxy)-6-(4,4-dimethyl-2-pentynyloxy)-pyrimidine (the present compound (7)) and 0.074 g of 4,6-bis(4,4-dimethyl-2-pentynyloxy)pyrimidine (the present compound (8)).

20 The melting point of 4-(2-butynyloxy)-6-(4,4-dimethyl-2-pentynyloxy)pyrimidine: 113°C

 The melting point of 4,6-bis(4,4-dimethyl-2-pentynyloxy)pyrimidine: 83.5°C

 Production Example 7

25 To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.38 g of potassium carbonate, and 0.1 g of phenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammo-

nium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water. The organic layer was dried over anhydrous magnesium sulfate and concentrated. The residue was subjected to silica gel column chromatography to give 0.16 g of 4-(2-butynyloxy)-6-phenoxy-
5 pyrimidine (the present compound (9)).

¹H-NMR: 1.86 (t, 3H), 4.97 (q, 2H), 6.17 (s, 1H), 7.14 (d, 2H), 7.25 (t, 1H), 7.42 (t, 2H), 8.47 (s, 1H)

Production Example 8

10 In 2 ml of tetrahydrofuran was suspended 0.1 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.12 g of 3-butyn-2-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes and then cooled to 0°C, to which 0.6 ml of a tetrahydrofuran solution containing 0.4 g of 4-
15 chloro-6-(2-propynyloxy)pyrimidine was slowly added dropwise. The mixture was further stirred at 0°C for 4.5 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and
20 then concentrated. The residue was subjected to silica gel column chromatography to give 0.14 g of 4-(1-methyl-2-propynyloxy)-6-(2-propynyloxy)pyrimidine (the present compound (11)).

¹H-NMR: 1.63 (d, 3H), 2.47 (d, 1H), 2.51 (t, 1H), 5.00 (d, 2H), 5.72-5.81 (m, 1H), 6.17 (s, 1H), 8.49 (s, 1H)

25 Production Example 9

In 2 ml of tetrahydrofuran was suspended 0.1 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.13 g of 3-butyn-1-ol was slowly added dropwise with stirring at room temperature.

The mixture was stirred at room temperature for 20 minutes and then cooled to 0°C, to which 0.6 ml of a tetrahydrofuran solution containing 0.3 g of 4-chloro-6-(2-propynyloxy)pyrimidine was slowly added dropwise. The mixture was further stirred at 0°C for 4.5 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.14 g of 4-(3-butynyloxy)-6-(2-propynyloxy)pyrimidine (the present compound (12)).

¹H-NMR: 2.02 (t, 1H), 2.50 (t, 1H), 2.64-2.70 (m, 2H), 4.46 (t, 2H), 5.00 (d, 2H), 6.15 (s, 1H), 8.44 (s, 1H)

Production Example 10

In 4 ml of tetrahydrofuran was suspended 0.13 g of sodium hydride (60% in oil), to which 0.7 ml of a tetrahydrofuran solution containing 0.12 g of 2-propyn-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes and then cooled to 0°C, to which 0.7 ml of a tetrahydrofuran solution containing 0.4 g of 4-chloro-6-benzyloxy pyrimidine was slowly added dropwise. The mixture was further stirred at 0°C for 4.5 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.17 g of 4-benzyloxy-6-(2-propynyloxy)pyrimidine (the present compound (13)).

¹H-NMR: 2.49 (t, 1H), 4.96 (d, 2H), 5.38 (s, 2H), 6.17 (s, 1H), 7.30-7.41 (m, 5H), 8.46 (s, 1H)

Production Example 11

To 5 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-propynyloxy)pyrimidine, 0.23 g of potassium carbonate and 0.4 g of 4-chlorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.1 g of 4-(4-chlorophenoxy)-6-(2-propynyloxy)-pyrimidine (the present compound (14)), m.p.: 100.3°C.

Production Example 12

To 5 ml of N,N-dimethylformamide were added 0.19 g of 4-chloro-6-(2-propynyloxy)pyrimidine, 0.22 g of potassium carbonate, and 0.13 g of 3-chlorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.26 g of 4-(3-chlorophenoxy)-6-(2-propynyloxy)pyrimidine (the present compound (15)), m.p.: 71.6°C.

Production Example 13

To 5 ml of N,N-dimethylformamide were added 0.3 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.34 g of potassium carbonate, and 0.24 g of 2-chloro-4-fluorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three

times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.53 g of 4-(2-chloro-4-fluorophenoxy)-6-(2-butyloxy)pyrimidine (the present compound (16)).

¹H-NMR: 1.87 (t, 3H), 4.98 (q, 2H), 6.28 (s, 1H), 7.00-7.23 (m, 3H), 8.41 (s, 1H)

Production Example 14

To 5 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-propynyloxy)pyrimidine, 0.25 g of potassium carbonate, and 0.19 g of 3-trifluoromethylphenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.26 g of 4-(3-trifluoromethylphenoxy)-6-(2-propynyloxy)pyrimidine (the present compound (17)).

¹H-NMR: 2.52 (t, 1H), 5.04 (d, 2H), 6.30 (s, 1H), 7.28-7.54 (m, 4H), 8.47 (s, 1H)

Production Example 15

To 5 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-propynyloxy)pyrimidine, 0.25 g of potassium carbonate, and 0.19 g of 2-trifluoromethylphenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous

magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.14 g of 4-(2-trifluoromethylphenoxy)-6-(2-propynyloxy)pyrimidine (the present compound (18)).

¹H-NMR: 2.54 (t, 1H), 5.04 (d, 2H), 6.35 (s, 1H), 7.24 (d, 1H), 7.34 (t, 1H), 7.61 (t, 1H), 7.72 (d, 1H), 8.45 (s, 1H)

Production Example 16

To 5 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-propynyloxy)pyrimidine, 0.25 g of potassium carbonate, and 0.15 g of 2-chlorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.14 g of 4-(2-chlorophenoxy)-6-(2-propynyloxy)pyrimidine (the present compound (19)), m.p.: 76.2°C.

Production Example 17

To 5 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-propynyloxy)pyrimidine, 0.25 g of potassium carbonate, and 0.19 g of 4-trifluoromethylphenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.21 g of 4-(4-trifluoromethylphenoxy)-6-(2-propynyloxy)pyrimidine (the present compound (20)).

¹H-NMR: 2.53 (t, 1H), 5.04 (d, 2H), 6.32 (s, 1H), 7.26 (d, 2H), 7.78 (d,

2H), 8.47 (s, 1H)

Production Example 18

To 5 ml of N,N-dimethylformamide were added 0.43 g of 4-chloro-6-(2-propynyloxy)pyrimidine, 0.52 g of potassium carbonate, and 0.4 g of 2,6-difluorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.53 g of 4-(2,6-difluorophenoxy)-6-(2-propynyloxy)pyrimidine (the present compound (21)), m.p.: 67.2°C.

Production Example 19

To 5 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-propynyloxy)pyrimidine, 0.25 g of potassium carbonate, and 0.21 g of 2,4-dichlorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.24 g of 4-(2,4-dichlorophenoxy)-6-(2-propynyloxy)pyrimidine (the present compound (22)), m.p.: 106.7°C.

Production Example 20

To 5 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-propynyloxy)pyrimidine, 0.25 g of potassium carbonate, and 0.21 g of 3,4-dichlorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a satu-

rated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel
5 column chromatography to give 0.14 g of 4-(3,4-dichlorophenoxy)-6-(2-propynyloxy)pyrimidine (the present compound (23)), m.p.: 109.2°C.

Production Example 21

To 5 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-propynyloxy)pyrimidine, 0.25 g of potassium carbonate, and 0.21 g of 3,5-
10 dichlorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magne-
15 sium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.24 g of 4-(3,5-dichlorophenoxy)-6-(2-propynyloxy)pyrimidine (the present compound (24)), m.p.: 136.5°C.

Production Example 22

To 5 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-
20 propynyloxy)pyrimidine, 0.25 g of potassium carbonate, and 0.21 g of 2,5-dichlorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with dilut-
25 ed hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.19 g of 4-(2,5-dichlorophenoxy)-6-(2-propynyloxy)pyrimidine (the present compound (25)), m.p.: 87.7°C.

Production Example 23

To 5 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-propynyloxy)pyrimidine, 0.25 g of potassium carbonate, and 0.21 g of 2,3-dichlorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.25 g of 4-(2,3-dichlorophenoxy)-6-(2-propynyloxy)pyrimidine (the present compound (26)), m.p.: 91.9°C.

Production Example 24

To 5 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-propynyloxy)pyrimidine, 0.25 g of potassium carbonate, and 0.14 g of 2-methylphenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.15 g of 4-(2-methylphenoxy)-6-(2-propynyloxy)pyrimidine (the present compound (27)), m.p.: 64.8°C.

Production Example 25

To 5 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-propynyloxy)pyrimidine, 0.25 g of potassium carbonate, and 0.14 g of 4-methylphenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times

with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.28 g of 4-(4-methylphenoxy)-6-(2-propynyloxy)pyrimidine (the present compound (28)).

¹H-NMR: 2.37 (s, 3H), 2.51 (t, 1H), 5.00 (d, 2H), 6.18 (s, 1H), 7.01 (d, 2H), 7.21 (d, 2H), 8.47 (s, 1H)

Production Example 26

To 5 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-propynyloxy)pyrimidine, 0.25 g of potassium carbonate, and 0.14 g of 3-methylphenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.27 g of 4-(3-methylphenoxy)-6-(2-propynyloxy)pyrimidine (the present compound (29)).

¹H-NMR: 2.36 (s, 3H), 2.52 (t, 1H), 5.01 (d, 2H), 6.17 (s, 1H), 6.93-6.94 (m, 2H), 7.07 (d, 1H), 7.29 (t, 1H), 8.47 (s, 1H)

Production Example 27

To 5 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-propynyloxy)pyrimidine, 0.25 g of potassium carbonate, and 0.18 g of 3-methoxyphenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magne-

sium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.23 g of 4-(3-methoxyphenoxy)-6-(2-propynyloxy)pyrimidine (the present compound (30)).

¹H-NMR: 2.51 (t, 1H), 3.80 (s, 3H), 5.02 (d, 2H), 6.19 (s, 1H), 6.67-6.83
5 (m, 3H), 7.32 (t, 1H), 8.49 (s, 1H)

Production Example 28

To 5 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-propynyloxy)pyrimidine, 0.25 g of potassium carbonate, and 0.18 g of 4-methoxyphenol, followed by stirring at 60°C for 7 hours. The reaction mix-
10 ture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel
15 column chromatography to give 0.14 g of 4-(4-methoxyphenoxy)-6-(2-propynyloxy)pyrimidine (the present compound (31)), m.p.: 72.0°C.

Production Example 29

To 5 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-propynyloxy)pyrimidine, 0.25 g of potassium carbonate, and 0.18 g of 2-methoxyphenol, followed by stirring at 60°C for 7 hours. The reaction mix-
20 ture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel
25 column chromatography to give 0.3 g of 4-(2-methoxyphenoxy)-6-(2-propynyloxy)pyrimidine (the present compound (32)).

¹H-NMR: 2.52 (t, 1H), 3.77 (s, 3H), 6.20 (s, 1H), 6.96-7.26 (m, 4H),

8.44 (s, 1H)

Production Example 30

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.23 g of potassium carbonate, and 0.17 g of 2,6-difluorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.15 g of 4-(2-butynyloxy)-6-(2,6-difluorophenoxy)pyrimidine (the present compound (33)), m.p.: 79.8°C.

Production Example 31

To 5 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-propynyloxy)pyrimidine, 0.25 g of potassium carbonate, and 0.16 g of 2-fluorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.21 g of 4-(2-fluorophenoxy)-6-(2-propynyloxy)pyrimidine (the present compound (34)).

¹H-NMR: 2.53 (t, 1H), 5.02 (d, 2H), 6.32 (s, 1H), 7.16-7.29 (m, 4H), 8.44 (s, 1H)

Production Example 32

To 5 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-propynyloxy)pyrimidine, 0.25 g of potassium carbonate, and 0.16 g of 4-fluo-

rophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted
5 hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.15 g of 4-(4-fluorophenoxy)-6-(2-propynyl-
oxy)pyrimidine (the present compound (35)), m.p.: 81.4°C.

Production Example 33

10 To 5 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-propynyloxy)pyrimidine, 0.25 g of potassium carbonate, and 0.16 g of 3-fluorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with
15 chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.26 g of 4-(3-fluorophenoxy)-6-(2-propynyl-
oxy)pyrimidine (the present compound (36)).

20 ¹H-NMR: 2.53 (t, 1H), 5.03 (d, 2H), 6.25 (s, 1H), 6.87-6.98 (m, 3H), 7.34-7.42 (m, 1H), 8.48 (s, 1H)

Production Example 34

In 5 ml of tetrahydrofuran were dissolved 194 mg of 4-chloro-6-phenylpyrimidine and 68 mg of 2-propyn-1-ol, to which 50 mg of sodium hydride
25 (60% in oil) was added with stirring at room temperature, followed by further stirring for 3 hours. The reaction mixture was then poured into water and extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magne-

sium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 190 mg of 4-phenyl-6-(2-propynyloxy)pyrimidine (the present compound (37)), m.p.: 65.1°C.

Production Example 35

5 In 5 ml of tetrahydrofuran were dissolved 186 mg of 4-chloro-6-phenylpyrimidine and 82 mg of 2-butyne-1-ol, to which 47 mg of sodium hydride (60% in oil) was added with stirring at room temperature, followed by further stirring for 3 hours. The reaction mixture was then poured into water and extracted with ethyl acetate. The organic layer was washed with a
10 saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 190 mg of 4-(2-butynyloxy)-6-phenylpyrimidine (the present compound (38)), m.p.: 59.6°C.

Production Example 36

15 To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.23 g of potassium carbonate, and 0.17 g of 2,3-difluorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with
20 chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.17 g of 4-(2-butynyloxy)-6-(2,3-difluorophenoxy)pyrimidine (the present compound (39)).

25 ¹H-NMR: 1.89 (t, 3H), 5.00 (q, 2H), 6.35 (s, 1H), 6.96-7.14 (m, 3H), 8.43 (s, 1H)

Production Example 37

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-

butynyloxy)pyrimidine, 0.23 g of potassium carbonate, and 0.16 g of 3-cyanophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with
5 chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.2 g of 4-(2-butynyloxy)-6-(3-cyanophenoxy)-pyrimidine (the present compound (40)), m.p.: 121.2°C.

10 Production Example 38

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.23 g of potassium carbonate and 0.16 g of 4-cyanophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with
15 chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.2 g of 4-(2-butynyloxy)-6-(4-cyanophenoxy)-pyrimidine (the present compound (41)), m.p.: 162.0°C.
20

 Production Example 39

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.23 g of potassium carbonate and 0.16 g of 2-cyanophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was
25 then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium

sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.2 g of 4-(2-butynyloxy)-6-(2-cyanophenoxy)-pyrimidine (the present compound (42)).

¹H-NMR: 1.89 (t, 3H), 5.01 (q, 2H), 6.43 (s, 1H), 7.23-7.39 (m, 2H),
5 7.63-7.74 (m, 2H), 8.44 (s, 1H)

Production Example 40

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.23 g of potassium carbonate, and 0.17 g of 2,5-difluorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture
10 was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel
15 column chromatography to give 0.26 g of 4-(2-butynyloxy)-6-(2,5-difluorophenoxy)pyrimidine (the present compound (43)).

¹H-NMR: 1.88 (t, 3H), 5.00 (q, 2H), 6.35 (s, 1H), 6.89-7.02 (m, 2H),
7.10-7.20 (m, 1H), 8.43 (s, 1H)

Production Example 41

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.23 g of potassium carbonate, and 0.17 g of 2,4-difluorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture
20 was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with
25 chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.27 g of 4-(2-butynyloxy)-6-(2,4-difluoro-

phenyl)pyrimidine (the present compound (44)), m.p.: 63.9°C.

Production Example 42

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.23 g of potassium carbonate, and 0.19 g of 2,4,6-trifluorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.3 g of 4-(2-butynyloxy)-6-(2,4,6-trifluorophenoxy)pyrimidine (the present compound (45)), m.p.: 60.3°C.

Production Example 43

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.23 g of potassium carbonate, and 0.19 g of 2,3,6-trifluorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.21 g of 4-(2-butynyloxy)-6-(2,3,6-trifluorophenoxy)pyrimidine (the present compound (46)).

¹H-NMR: 1.88 (t, 3H), 5.01 (q, 2H), 6.46 (s, 1H), 6.91-7.11 (m, 2H), 8.41 (s, 1H)

Production Example 44

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.23 g of potassium carbonate, and 0.22 g of 2-chlo-

ro-4,6-difluorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with
5 diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.34 g of 4-(2-butynyloxy)-6-(2-chloro-4,6-difluorophenoxy)pyrimidine (the present compound (47)).

¹H-NMR: 1.88 (t, 3H), 5.01 (q, 2H), 6.43 (s, 1H), 6.87-6.95 (m, 1H),
10 7.03-7.08 (m, 1H), 8.40 (s, 1H)

Production Example 45

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.23 g of potassium carbonate and 0.24 g of 4-fluoro-3-trifluoromethylphenol, followed by stirring at 60°C for 7 hours. The
15 reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was
20 subjected to silica gel column chromatography to give 0.35 g of 4-(2-butynyloxy)-6-(4-fluoro-3-trifluoromethylphenoxy)pyrimidine (the present compound (48)), m.p.: 90.1°C.

Production Example 46

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.23 g of potassium carbonate, and 0.23 g of 3-tri-
25 fluoromethoxyphenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three

times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.33 g of 4-(2-butynyloxy)-6-(3-tri-
5 fluoromethoxy)pyrimidine (the present compound (49)), m.p.: 63.1°C.

Production Example 47

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.23 g of potassium carbonate, and 0.23 g of 4-tri-
10 fluoromethoxyphenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to
15 silica gel column chromatography to give 0.32 g of 4-(2-butynyloxy)-6-(4-tri-fluoromethoxy)pyrimidine (the present compound (50)), m.p.: 87.7°C.

Production Example 48

To 6 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-propynyloxy)pyrimidine, 0.25 g of potassium carbonate, and 0.13 g of phenol,
20 followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then
25 concentrated. The residue was subjected to silica gel column chromatography to give 0.15 g of 4-phenoxy-6-(2-propynyloxy)pyrimidine (the present compound (51)), m.p.: 71.1°C.

Production Example 49

In 4 ml of tetrahydrofuran was suspended 0.27 g of sodium hydride (60% in oil), to which 0.7 ml of a tetrahydrofuran solution containing 0.33 g of 2-propyn-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes and
5 then cooled to 0°C, to which 0.7 ml of a tetrahydrofuran solution containing 0.4 g of 4,6-dichloropyrimidine was slowly added dropwise. After completion of the dropwise addition, the mixture was stirred at room temperature for 40 minutes. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloro-
10 form. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.35 g of 4,6-bis(2-propynyloxy)pyrimidine (the present compound (52)), m.p.: 74.0°C.

Production Example 50

15 To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.23 g of potassium carbonate and 0.19 g of 4-chloro-2-fluorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times
20 with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.29 g of 4-(2-butynyloxy)-6-(4-chloro-2-fluorophenoxy)pyrimidine (the present compound (4)), m.p.: 117.7°C.

25 Production Example 51

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.23 g of potassium carbonate, and 0.17 g of 3,4-difluorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture

was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.2 g of 4-(2-butynyloxy)-6-(3,4-difluorophen-
5 oxy)pyrimidine (the present compound (10)), m.p.: 109.7°C.

Production Example 52

In 1.1 ml of ethanol were dissolved 0.2 g of 4-chloro-6-(2-butynyloxy)-
10 pyrimidine and 0.15 g of aniline, followed by heating under reflux for 7 hours. The reaction mixture was then left for cooling to room temperature and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography to give 0.13 g of 4-anilino-6-(2-butynyloxy)pyrimidine (the present compound (53)), m.p.: 159.3°C.

15 Production Example 53

In 5 ml of N,N-dimethylformamide were dissolved 212 mg of 4-chloro-6-(2-fluorophenyl)pyrimidine and 68 mg of 2-propyn-1-ol, to which 49 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 8 hours. The reaction mixture was then poured into water and
20 extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 170 mg of 4-(2-fluorophenyl)-6-(2-propynyloxy)pyrimidine (the present compound (54)).

25 ¹H-NMR: 2.53 (t, 1H), 5.08 (d, 2H), 7.14-7.24 (m, 1H), 7.26-7.36 (m, 2H), 7.28 (s, 1H), 8.02 (dt, 1H), 8.88 (s, 1H)

Production Example 54

In 5 ml of N,N-dimethylformamide were dissolved 207 mg of 4-chlo-

ro-6-(2-fluorophenyl)pyrimidine and 83 mg of 2-butyne-1-ol, to which 48 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 8 hours. The reaction mixture was then poured into water and extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 160 mg of 4-(2-fluorophenyl)-6-(2-butyne-1-oxo)pyrimidine (the present compound (55)).

¹H-NMR: 1.89 (t, 3H), 5.03 (q, 2H), 7.13-7.36 (m, 2H), 7.31 (s, 1H), 7.38-7.50 (m, 1H), 8.02 (dt, 1H), 8.87 (s, 1H)

Production Example 55

In 7 ml of N,N-dimethylformamide were dissolved 207 mg of 4-chloro-6-(3-fluorophenyl)pyrimidine and 67 mg of 2-propyne-1-ol, to which 48 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 6 hours. The reaction mixture was then poured into water and extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 170 mg of 4-(3-fluorophenyl)-6-(2-propynyloxy)pyrimidine (the present compound (56)).

¹H-NMR: 2.53 (t, 1H), 5.09 (d, 2H), 7.15 (s, 1H), 7.15-7.25 (m, 1H), 7.49-7.51 (m, 1H), 7.73-7.83 (m, 2H), 8.86 (s, 1H)

Production Example 56

In 7 ml of N,N-dimethylformamide were dissolved 204 mg of 4-chloro-6-(3-fluorophenyl)pyrimidine and 82 mg of 2-butyne-1-ol, to which 47 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 6 hours. The reaction mixture was then poured into water and extracted with ethyl acetate. The organic layer was washed with a saturated

ed aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 151 mg of 4-(3-fluorophenyl)-6-(2-butyloxy)pyrimidine (the present compound (57)).

5 ¹H-NMR: 1.89 (t, 3H), 5.04 (q, 2H), 7.13-7.26 (m, 1H), 7.13 (s, 1H), 7.40-7.52 (m, 1H), 7.70-7.83 (m, 2H), 8.85 (s, 1H)

Production Example 57

In 7 ml of N,N-dimethylformamide were dissolved 199 mg of 4-chloro-6-(4-fluorophenyl)pyrimidine and 64 mg of 2-propyn-1-ol, to which 46 mg
10 of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 9 hours. The reaction mixture was then poured into water and extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica
15 gel column chromatography to give 150 mg of 4-(4-fluorophenyl)-6-(2-propynyloxy)pyrimidine (the present compound (58)).

¹H-NMR: 2.53 (t, 1H), 5.08 (d, 2H), 7.12 (s, 1H), 7.14-7.20 (m, 2H), 8.01-8.05 (m, 2H), 8.84 (s, 1H)

Production Example 58

20 In 8 ml of N,N-dimethylformamide were dissolved 207 mg of 4-chloro-6-(4-fluorophenyl)pyrimidine and 83 mg of 2-butyne-1-ol, to which 48 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 9 hours. The reaction mixture was then poured into water and extracted with ethyl acetate. The organic layer was washed with a saturated
25 aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 138 mg of 4-(4-fluorophenyl)-6-(2-butyloxy)pyrimidine (the present compound (59)).

¹H-NMR: 1.89 (t, 3H), 5.04 (q, 2H), 7.10 (s, 1H), 7.15-7.23 (m, 2H), 7.99-8.09 (m, 2H), 8.83 (s, 1H)

Production Example 59

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-pentynyloxy)pyrimidine, 0.21 g of potassium carbonate, and 0.16 g of 2,3-difluorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.26 g of 4-(2,3-difluorophenoxy)-6-(2-pentynyloxy)pyrimidine (the present compound (60)).

¹H-NMR: 1.57 (t, 3H), 2.04-2.30 (qt, 2H), 5.02 (t, 2H), 6.35 (s, 1H), 6.97-7.13 (m, 3H), 8.42 (s, 1H)

Production Example 60

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.23 g of potassium carbonate, and 0.15 g of 3-fluorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.23 g of 4-(3-fluorophenoxy)-6-(2-butynyloxy)pyrimidine (the present compound (61)), m.p.: 60.1°C.

Production Example 61

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-

butynyloxy)pyrimidine, 0.23 g of potassium carbonate, and 0.15 g of 4-fluorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with
5 chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.19 g of 4-(4-fluorophenoxy)-6-(2-butynyloxy)-pyrimidine (the present compound (62)), m.p.: 115.8°C.

10 Production Example 62

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.23 g of potassium carbonate, and 0.15 g of 2-fluorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated
15 aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.26 g of 4-(2-fluorophenoxy)-6-(2-butyloxy)pyrimidine (the present compound (63)).
20

¹H-NMR: 1.88 (t, 3H), 4.99 (q, 2H), 6.31 (s, 1H), 7.16-7.27 (m, 4H), 8.44 (s, 1H)

 Production Example 63

To 2 ml of N,N-dimethylformamide were added 183 mg of 4-chloro-6-(2-butynyloxy)pyrimidine, 166 mg of potassium carbonate, and 138 mg of
25 2,3-methylenedioxyphenol, followed by stirring at 80°C for 7 hours and then at 120°C for 3 hours. The reaction mixture was then left for cooling to room temperature and subjected to phase separation three times between ethyl

acetate and an aqueous sodium chloride solution. The organic layers were dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 193 mg of 4-(2-butynyloxy)-6-(2,3-methylenedioxyphenoxy)pyrimidine (the present compound (64)).

¹H-NMR: 1.88 (t, 3H), 4.98 (q, 2H), 5.97 (s, 2H), 6.28 (s, 1H), 6.70 (d, 1H), 6.77 (d, 1H), 6.87 (t, 1H), 8.47 (s, 1H)

Production Example 64

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.23 g of potassium carbonate, and 0.21 g of 2-fluoro-4-nitrophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.07 g of 4-(2-fluoro-4-nitrophenoxy)-6-(2-butynyloxy)pyrimidine (the present compound (65)), m.p.: 132.1°C.

Production Example 65

In 2 ml of tetrahydrofuran was suspended 0.04 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.05 g of 2-butyne-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes and then cooled to 0°C, to which 0.6 ml of a tetrahydrofuran solution containing 0.16 g of 4-chloro-6-(N-methyl-N-(2,3-difluorophenyl)amino)pyrimidine was slowly added dropwise, followed by stirring for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were

combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.14 g of 4-(N-methyl-N-(2,3-difluorophenyl)amino)-6-(2-butyloxy)pyrimidine (the present compound (66)), m.p.: 77.5°C.

5 Production Example 66

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butyloxy)pyrimidine, 0.23 g of potassium carbonate, and 0.16 g of 2,3-dimethylphenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water. The organic layer was dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.26 g of 4-(2,3-dimethylphenoxy)-6-(2-butyloxy)pyrimidine (the present compound (67)).

¹H-NMR: 1.87 (t, 3H), 2.06 (s, 3H), 2.31 (s, 3H), 4.96 (q, 2H), 6.07 (s, 1H), 6.87-7.16 (m, 3H), 8.46 (s, 1H)

 Production Example 67

To 2 ml of N,N-dimethylformamide were added 0.15 g of 4-(2-butyloxy)-6-methanesulfonylpyrimidine, 0.14 g of potassium carbonate, and 0.16 g of 2,6-difluorobenzylalcohol, followed by stirring at 50°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.14 g of 4-(2,6-difluorobenzoyloxy)-6-(2-butyloxy)pyrimidine (the present compound (68)).

¹H-NMR: 1.87 (t, 3H), 4.95 (q, 2H), 5.47 (s, 2H), 6.13 (s, 1H), 6.91-6.98 (m, 2H), 7.28-7.40 (m, 1H), 8.50 (s, 1H)

Production Example 68

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.23 g of potassium carbonate, and 0.22 g of 3-phenylphenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water. The organic layer was dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.25 g of 4-(3-phenylphenoxy)-6-(2-butynyloxy)pyrimidine (the present compound (69)).

¹H-NMR: 1.86 (t, 3H), 4.97 (q, 2H), 6.22 (s, 1H), 7.11-7.14 (m, 1H), 7.34-7.59 (m, 8H), 8.49 (s, 1H)

Production Example 69

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.23 g of potassium carbonate, and 0.24 g of 3-phenoxyphenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water. The organic layer was dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.33 g of 4-(2-butynyl-oxy)-6-(3-phenoxyphenoxy)pyrimidine (the present compound (70)).

¹H-NMR: 1.86 (t, 3H), 4.97 (q, 2H), 6.19 (s, 1H), 6.77-6.90 (m, 3H), 7.04-7.15 (m, 3H), 7.31-7.38 (m, 3H), 8.47 (s, 1H)

Production Example 70

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.23 g of potassium carbonate, and 0.18 g of 3'-hydroxyacetophenone, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.25 g of 4-(2-butynyloxy)-6-(3-acetylphenoxy)pyrimidine (the present compound (71)), m.p.: 94.0°C.

Production Example 71

In 5 ml of tetrahydrofuran was suspended 0.41 g of potassium t-butoxide, to which 0.56 g of (2,3-difluorophenyl)acetonitrile and 0.5 g of 4-chloro-6-(2-butynyloxy)pyrimidine were added, followed by stirring at room temperature for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined and washed with a saturated aqueous sodium chloride solution. The organic layer was dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.26 g of 4-(α -cyano-2,3-difluorobenzyl)-6-(2-butynyloxy)pyrimidine (the present compound (72)).

¹H-NMR: 1.87 (t, 3H), 5.00 (q, 2H), 5.46 (s, 1H), 6.88 (s, 1H), 6.88-7.35 (m, 3H), 8.77 (s, 1H)

25 Production Example 72

In 4 ml of tetrahydrofuran was suspended 0.33 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.42 g of 2,3-difluoroaniline were slowly added dropwise with stirring at room tem-

perature. The mixture was stirred at room temperature for 20 minutes, to which 1 ml of a tetrahydrofuran solution containing 0.5 g of 4-chloro-6-(2-butynyloxy)pyrimidine was slowly added dropwise, followed by stirring at room temperature for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.25 g of 4-(N-(2,3-difluorophenyl)amino)-6-(2-butynyloxy)pyrimidine (the present compound (73)), m.p.: 147.6°C.

Production Example 73

In 5 ml of tetrahydrofuran was suspended 0.46 g of potassium t-butoxide, to which 0.39 g of phenylacetonitrile and 0.5 g of 4-chloro-6-(2-butynyloxy)pyrimidine were added, followed by stirring at room temperature for 8 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined and washed with a saturated aqueous sodium chloride solution. The organic layer was dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.16 g of 4-(α -cyanobenzyl)-6-(2-butynyloxy)pyrimidine (the present compound (74)) and 0.19 g of 4-benzoyl-6-(2-butynyloxy)pyrimidine (the present compound (75)).

The present compound (74):

¹H-NMR: 1.86 (t, 3H), 4.98 (q, 2H), 5.17 (s, 1H), 6.84 (s, 1H), 7.35-7.46 (m, 5H), 8.77 (s, 1H)

The present compound (75):

¹H-NMR: 1.88 (t, 3H), 5.06 (q, 2H), 7.32 (s, 1H), 7.48 (t, 2H), 7.61 (t, 1H), 8.06 (d, 2H), 8.93 (s, 1H)

Production Example 74

To 2 ml of N,N-dimethylformamide were added 0.15 g of 4-(2-butyn-
yloxy)-6-methanesulfonylpyrimidine, 0.14 g of potassium carbonate, and 0.16
g of 2,3-difluorobenzylalcohol, followed by stirring at 50°C for 7 hours. The
5 reaction mixture was then left for cooling to room temperature and poured
into a saturated aqueous ammonium chloride solution, which was extracted
three times with chloroform. The chloroform layers were combined, washed
with diluted hydrochloric acid and then with water, and dried over anhy-
drous magnesium sulfate and then concentrated. The residue was sub-
10 jected to silica gel column chromatography to give 0.13 g of 4-(2,3-difluoro-
benzyloxy)-6-(2-butynyloxy)pyrimidine (the present compound (76)), m.p.:
84.9°C.

Production Example 75

In 10 ml of N,N-dimethylformamide were dissolved 304 mg of 4-chlo-
15 ro-2-methyl-6-phenylpyrimidine and 92 mg of 2-propyn-1-ol, to which 66 mg
of sodium hydride (60% in oil) was added, followed by stirring at room tem-
perature for 12 hours. The reaction mixture was then poured into water
and extracted with ethyl acetate. The organic layer was washed with a
saturated aqueous sodium chloride solution, dried over anhydrous magne-
20 sium sulfate, and then concentrated. The resulting residue was subjected
to silica gel column chromatography to give 256 mg of 2-methyl-4-phenyl-6-
(2-propynyloxy)pyrimidine (the present compound (77)).

¹H-NMR: 2.53 (t, 1H), 2.68 (s, 3H), 5.07 (d, 2H), 6.97 (s, 1H), 7.46 (m,
3H), 8.00 (m, 2H)

25 Production Example 76

In 10 ml of N,N-dimethylformamide were dissolved 313 mg of 4-chlo-
ro-2-methyl-6-phenylpyrimidine and 118 mg of 2-butyn-1-ol, to which 67 mg
of sodium hydride (60% in oil) was added, followed by stirring at room tem-

perature for 12 hours. The reaction mixture was then poured into water and extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected
5 to silica gel column chromatography to give 260 mg of 2-methyl-4-phenyl-6-(2-butynyloxy)pyrimidine (the present compound (78)).

¹H-NMR: 1.90 (t, 3H), 2.68 (s, 3H), 5.03 (q, 2H), 6.98 (s, 1H), 7.46 (m, 3H), 8.00 (m, 2H)

Production Example 77

10 In 2 ml of tetrahydrofuran was suspended 0.03 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.04 g of 2-butyne-1-ol were slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 0.6 ml of a tetrahydrofuran solution containing 0.16 g of 4-chloro-6-
15 (N-methoxymethyl-N-(2,3-difluorophenyl)amino)pyrimidine was slowly added dropwise at room temperature, followed by stirring for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous mag-
20 nesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.14 g of 4-(N-methoxymethyl-N-(2,3-difluorophenyl)amino)-6-(2-butynyloxy)pyrimidine (the present compound (79)).

¹H-NMR: 1.85 (t, 3H), 3.42 (s, 3H), 4.91 (q, 2H), 5.28 (s, 2H), 5.72 (s, 25 1H), 7.09-7.25 (m, 3H), 8.43 (s, 1H)

Production Example 78

In 2 ml of tetrahydrofuran was suspended 0.05 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.07 g

of 2-butyne-1-ol were slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 0.6 ml of a tetrahydrofuran solution containing 0.2 g of 4-chloro-6-(2,6-difluorobenzyl)pyrimidine was slowly added dropwise at room temperature, followed by stirring for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.24 g of 4-(2,6-difluorobenzyl)-6-(2-butyneoxy)pyrimidine (the present compound (80)), m.p.: 57.6°C.

Production Example 79

In 0.8 ml of tetrahydrofuran was suspended 0.04 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.15 g of 4-anilino-6-(2-butyneoxy)pyrimidine were slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 0.4 ml of a tetrahydrofuran solution containing 0.12 g of iodoethane was slowly added dropwise at room temperature, followed by stirring for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.12 g of 4-(N-ethyl-N-phenylamino)-6-(2-butyneoxy)pyrimidine (the present compound (81)).

¹H-NMR: 1.20 (t, 3H), 1.83 (t, 3H), 3.98 (q, 2H), 4.85 (q, 2H), 5.54 (s, 1H), 7.19 (d, 2H), 7.32 (t, 1H), 7.44 (t, 2H), 8.38 (s, 1H)

Production Example 80

In 6 ml of N,N-dimethylformamide were dissolved 96 mg of 4-chloro-6-(2,3-difluorophenyl)pyrimidine and 26 mg of 2-propyn-1-ol, to which 19 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 8 hours. The reaction mixture was then poured into water and
5 extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 42 mg of 4-(2,3-difluorophenyl)-6-(2-propynyloxy)pyrimidine (the present compound (82)).

10 ¹H-NMR: 2.54 (t, 1H), 5.10 (d, 2H), 7.15-7.34 (m, 2H), 7.32 (s, 1H), 7.89 (s, 1H), 8.90 (s, 1H)


Production Example 81

In 10 ml of N,N-dimethylformamide were dissolved 280 mg of 4-chloro-6-(2,3-difluorophenyl)pyrimidine and 96 mg of 2-butyn-1-ol, to which 55
15 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 8 hours. The reaction mixture was then poured into water and extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected
20 to silica gel column chromatography to give 212 mg of 4-(2,3-difluorophenyl)-6-(2-butyloxy)pyrimidine (the present compound (83)).

¹H-NMR: 1.90 (t, 3H), 5.06 (q, 2H), 7.15-7.34 (m, 2H), 7.30 (s, 1H), 7.88 (t, 1H), 8.89 (s, 1H)

Production Example 82

25 In 5 ml of tetrahydrofuran was suspended 0.46 g of potassium t-butoxide, to which 0.56 g of 2-chloro-6-fluorobenzylcyanide of formula



and 0.5 g of 4-chloro-6-(2-butynyloxy)pyrimidine were added, followed by stirring at room temperature 8 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined and washed with a saturated aqueous sodium chloride solution. The organic layer was dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.25 g of 4-(α -cyano-2-chloro-6-fluorobenzyl)-6-(2-butynyloxy)pyrimidine (the present compound (84)).

¹H-NMR: 1.88 (t, 3H), 5.00 (q, 2H), 5.82 (s, 1H), 6.97 (s, 1H), 7.07-7.14 (m, 1H), 7.28-7.42 (m, 2H), 8.75 (s, 1H)

Production Example 83

In 2 ml of tetrahydrofuran was suspended 0.05 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.07 g of 2-butyne-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 0.6 ml of a tetrahydrofuran solution containing 0.24 g of 4-chloro-6-(N-cyanomethyl-N-(2,3-difluorophenyl)amino)pyrimidine was slowly added dropwise, followed by stirring at room temperature for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.22 g of 4-(2-butynyloxy)-6-(N-cyanomethyl-N-(2,3-difluorophenyl)amino)pyrimidine (the present

compound (85)).

¹H-NMR: 1.85 (t, 3H), 4.84 (s, 2H), 4.91 (q, 2H), 5.66 (s, 1H), 7.16-7.30 (m, 3H), 8.51 (s, 1H)

Production Example 84

5 In 10 ml of N,N-dimethylformamide were dissolved 522 mg of 4-chloro-6-(2-fluorophenyl)pyrimidine and 231 mg of 2-pentyn-1-ol, to which 110 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 6 hours. The reaction mixture was then poured into water and extracted with ethyl acetate. The organic layer was washed with a
10 saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 470 mg of 4-(2-fluorophenyl)-6-(2-pentynyloxy)pyrimidine (the present compound (86)).

¹H-NMR: 1.19 (t, 3H), 2.22-2.36 (m, 2H), 5.05 (q, 2H), 7.11-7.30 (m, 2H), 7.34 (s, 1H), 7.40-7.50 (m, 1H), 8.02 (dt, 1H), 8.89 (s, 1H)
15

Production Example 85

In 10 ml of N,N-dimethylformamide were dissolved 226 mg of 4-chloro-6-(2,6-difluorophenyl)pyrimidine and 84 mg of 2-butyne-1-ol, to which 48 mg of sodium hydride (60% in oil) was added, followed by stirring at room
20 temperature for 9 hours. The reaction mixture was then poured into water and extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 114 mg of 4-(2,6-difluorophenyl)-
25 6-(2-butynyloxy)pyrimidine (the present compound (87)).

¹H-NMR: 1.89 (t, 3H), 5.05 (q, 2H), 6.98-7.10 (m, 2H), 7.29 (s, 1H), 7.33-7.46 (m, 1H), 8.92 (s, 1H)

Production Example 86

In 2 ml of tetrahydrofuran was suspended 0.05 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.08 g of 2-butyne-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 0.6 ml
5 of a tetrahydrofuran solution containing 0.2 g of 4-chloro-6-(2-fluorobenzyl)-pyrimidine was slowly added dropwise at room temperature, followed by stirring for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over
10 anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.19 g of 4-(2-fluorobenzyl)-6-(2-butyneoxy)pyrimidine (the present compound (88)).

¹H-NMR: 1.85 (t, 3H), 4.07 (s, 2H), 4.95 (q, 2H), 6.54 (s, 1H), 7.02-7.12 (m, 2H), 7.21-7.30 (m, 2H), 8.72 (s, 1H)

15 Production Example 87

In 2 ml of tetrahydrofuran was suspended 0.05 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.07 g of 2-butyne-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 0.6 ml
20 of a tetrahydrofuran solution containing 0.2 g of 4-chloro-6-(2-chlorobenzyl)-pyrimidine was slowly added dropwise at room temperature, followed by stirring for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over
25 anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.19 g of 4-(2-chlorobenzyl)-6-(2-butyneoxy)pyrimidine (the present compound (89)).

¹H-NMR: 1.85 (t, 3H), 4.18 (s, 2H), 4.94 (q, 2H), 6.48 (s, 1H), 7.21-7.31

(m, 3H), 7.37-7.40 (m, 1H), 8.73 (s, 1H)

Production Example 88

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.23 g of potassium carbonate, and 0.22 g of 2,3,5,6-tetrafluorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined and washed with diluted hydrochloric acid and then with water. The organic layer was dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.10 g of 4-(2,3,5,6-tetrafluorophenoxy)-6-(2-butynyloxy)pyrimidine (the present compound (90)).

¹H-NMR: 1.89 (t, 3H), 5.02 (q, 2H), 6.48 (s, 1H), 6.97-7.06 (m, 1H), 8.40 (s, 1H)

Production Example 89

In 2 ml of tetrahydrofuran was suspended 0.06 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.08 g of 2-butyne-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 0.6 ml of a tetrahydrofuran solution containing 0.2 g of 4-chloro-6-benzylpyrimidine was slowly added dropwise at room temperature, followed by stirring for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.21 g of 4-(2-butynyloxy)-6-benzylpyrimidine (the present compound (91)).

¹H-NMR: 1.84 (t, 3H), 4.02 (s, 2H), 4.94 (q, 2H), 6.52 (s, 1H), 7.23-7.34 (m, 5H), 8.72 (s, 1H)

Production Example 90

In 2 ml of tetrahydrofuran was suspended 0.06 g of sodium hydride
5 (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.08 g
of 2-butyne-1-ol was slowly added dropwise with stirring at room temperature.
The mixture was stirred at room temperature for 20 minutes, to which 0.6 ml
of a tetrahydrofuran solution containing 0.2 g of 4-chloro-6-(2-methylbenzyl)pyrimidine was slowly added dropwise at room temperature, followed by
10 stirring for 4 hours. The reaction mixture was then poured into a saturated
aqueous ammonium chloride solution and extracted three times with chloroform.
The chloroform layers were combined, washed with water, dried over
anhydrous magnesium sulfate, and then concentrated. The residue was
subjected to silica gel column chromatography to give 0.25 g of 4-(2-butyne-1-
15 oxy)-6-(2-methylbenzyl)pyrimidine (the present compound (92)).

¹H-NMR: 1.84 (t, 3H), 2.23 (s, 3H), 4.05 (s, 2H), 4.93 (q, 2H), 6.36 (s, 1H), 7.16-7.19 (m, 4H), 8.72 (s, 1H)

Production Example 91

In 3 ml of tetrahydrofuran was suspended 0.10 g of sodium hydride
20 (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.14 g
of 2-butyne-1-ol was slowly added dropwise with stirring at room temperature.
The mixture was stirred at room temperature for 20 minutes, to which 0.6 ml
of a tetrahydrofuran solution containing 0.36 g of 4-chloro-6-(N-methyl-N-phenylamino)pyrimidine was slowly added dropwise at room temperature,
25 followed by stirring for 6 hours. The reaction mixture was then poured into
a saturated aqueous ammonium chloride solution and extracted three times
with chloroform. The chloroform layers were combined, washed with water,
dried over anhydrous magnesium sulfate, and then concentrated. The resi-

due was subjected to silica gel column chromatography to give 0.18 g of 4-(N-methyl-N-phenylamino)-6-(2-butynyloxy)pyrimidine (the present compound (93)), m.p.: 57.6°C.

Production Example 92

5 In 2 ml of tetrahydrofuran was suspended 0.06 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.08 g of 2-butyne-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 0.6 ml of a tetrahydrofuran solution containing 0.2 g of 4-chloro-6-(α -methylbenzyl)pyrimidine was slowly added dropwise at room temperature, followed by
10 stirring for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was
15 subjected to silica gel column chromatography to give 0.16 g of 4-(2-butyne-1-ol)-6-(α -methylbenzyl)pyrimidine (the present compound (94)).

$^1\text{H-NMR}$: 1.66 (d, 3H), 1.84 (t, 3H), 4.14 (q, 1H), 4.94 (q, 2H), 6.57 (s, 1H), 7.18-7.33 (m, 5H), 8.73 (s, 1H)

Production Example 93

20 In 2 ml of tetrahydrofuran was suspended 0.04 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.06 g of 2-butyne-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 0.6 ml of a tetrahydrofuran solution containing 0.2 g of 4-chloro-6-(2-trifluoromethylbenzyl)pyrimidine was slowly added dropwise at room temperature, fol-
25 lowed by stirring for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water,

dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.24 g of 4-(2-butynyloxy)-6-(2-trifluoromethylbenzyl)pyrimidine (the present compound (95)).

5 ¹H-NMR: 1.84 (t, 3H), 4.25 (s, 2H), 4.94 (q, 2H), 6.38 (s, 1H), 7.38 (t, 2H), 7.49 (t, 1H), 7.69 (d, 1H), 8.73 (s, 1H)

Production Example 94

In 2 ml of tetrahydrofuran was suspended 0.05 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.07 g
10 of 2-butyne-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 0.6 ml of a tetrahydrofuran solution containing 0.2 g of 4-chloro-6-(2,3-difluorobenzyl)pyrimidine was slowly added dropwise at room temperature, followed by stirring for 4 hours. The reaction mixture was then poured into a saturated
15 aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.22 g of 4-(2-butynyloxy)-6-(2,3-difluorobenzyl)pyrimidine (the present compound (96)).

20 ¹H-NMR: 1.85 (t, 3H), 4.09 (s, 2H), 4.96 (q, 2H), 6.56 (s, 1H), 7.01-7.10 (m, 3H), 8.72 (s, 1H)

Production Example 95

To 5 ml of chloroform were added 0.57 ml of triethylamine and 0.5 g of 4-chloro-6-(2-butynyloxy)pyrimidine, to which 0.6 ml of a chloroform solution containing 0.33 g of thiophenol was slowly added dropwise, followed by
25 stirring at room temperature for 8 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed

with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.22 g of 4-(2-butynyloxy)-6-(thiophenoxy)pyrimidine (the present compound (97)).

¹H-NMR: 1.84 (t, 3H), 4.90 (q, 2H), 6.14 (s, 1H), 7.45-7.47 (m, 3H),
5 7.57-7.60 (m, 2H), 8.54 (s, 1H)

Production Example 96

In 1.5 ml of tetrahydrofuran was suspended 0.05 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.2 g of 4-(2-butynyloxy)-6-anilinopyrimidine was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 0.3 ml of a tetrahydrofuran solution containing 0.17 g of iodopropane was slowly added dropwise at room temperature, followed by stirring for 8 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with
15 chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.12 g of 4-(N-propyl-N-phenylamino)-6-(2-butynyloxy)pyrimidine (the present compound (98)).

20 ¹H-NMR: 0.91 (t, 3H), 1.63 (dt, 3H), 1.84 (t, 3H), 3.88 (t, 3H), 4.85 (q, 2H), 5.53 (s, 1H), 7.19 (d, 2H), 7.32 (t, 1H), 7.44 (t, 2H), 8.37 (s, 1H)

Production Example 97

In 2 ml of tetrahydrofuran was suspended 0.05 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.08 g of 2-butyne-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 0.6 ml of a tetrahydrofuran solution containing 0.2 g of 4-chloro-6-(2,4-difluorobenzyl)pyrimidine was slowly added dropwise at room temperature, followed by

stirring for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was
5 subjected to silica gel column chromatography to give 0.21 g of 4-(2-butynyl-oxy)-6-(2,4-difluorobenzyl)pyrimidine (the present compound (99)).

¹H-NMR: 1.86 (t, 3H), 4.02 (s, 2H), 4.96 (q, 2H), 6.54 (s, 1H), 6.78-6.89 (m, 2H), 7.20-7.31 (m, 1H), 8.72 (s, 1H)

Production Example 98

10 In 2 ml of tetrahydrofuran was suspended 0.05 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.08 g of 2-butyn-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 0.6 ml of a tetrahydrofuran solution containing 0.2 g of 4-chloro-6-(3-fluorobenzyl)-
15 pyrimidine was slowly added dropwise at room temperature, followed by stirring for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was
20 subjected to silica gel column chromatography to give 0.21 g of 4-(2-butynyl-oxy)-6-(3-fluorobenzyl)pyrimidine (the present compound (100)), m.p. 51.5°C.

Production Example 99

In 2 ml of tetrahydrofuran was suspended 0.05 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.07 g
25 of 2-butyn-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 0.6 ml of a tetrahydrofuran solution containing 0.2 g of 4-chloro-6-(2-chloro-6-fluorobenzyl)pyrimidine was slowly added dropwise at room temperature, fol-

lowed by stirring for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.26 g of 4-(2-butynyloxy)-6-(2-chloro-6-fluorobenzyl)pyrimidine (the present compound (101)).

¹H-NMR: 1.85 (t, 3H), 4.25 (s, 2H), 4.94 (q, 2H), 6.43 (s, 1H), 7.00-7.07 (m, 1H), 7.21-7.24 (m, 2H), 8.72 (s, 1H)

10 Production Example 100

In 2 ml of tetrahydrofuran was suspended 0.05 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.07 g of 2-butyn-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 0.6 ml of a tetrahydrofuran solution containing 0.2 g of 4-chloro-6-(3-chloro-2-fluorobenzyl)pyrimidine was slowly added dropwise at room temperature, followed by stirring for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.23 g of 4-(2-butynyloxy)-6-(3-chloro-2-fluorobenzyl)pyrimidine (the present compound (102)).

¹H-NMR: 1.86 (t, 3H), 4.07 (s, 2H), 4.95 (q, 2H), 6.56 (s, 1H), 7.02 (t, 1H), 7.15-7.35 (m, 2H), 8.72 (s, 1H)

25 Production Example 101

In 2 ml of tetrahydrofuran was suspended 0.04 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.06 g

of 2-butyne-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 0.6 ml of a tetrahydrofuran solution containing 0.2 g of 4-chloro-6-(2-bromobenzyl)-pyrimidine was slowly added dropwise at room temperature, followed by
5 stirring for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.22 g of 4-(2-butyne-
10 oxy)-6-(2-bromobenzyl)pyrimidine (the present compound (103)).

¹H-NMR: 1.86 (t, 3H), 4.20 (s, 2H), 4.94 (q, 2H), 6.50 (s, 1H), 7.11-7.17 (m, 1H), 7.28-7.30 (m, 2H), 7.57-7.60 (m, 1H), 8.73 (s, 1H)

Production Example 102

To 2 ml of N,N-dimethylformamide were added 183 mg of 4-chloro-6-(2-butyneoxy)pyrimidine, 166 mg of potassium carbonate, and 87 mg of N-ethylpropylamine, followed by stirring at 60°C for 7 hours. Then, 166 mg of potassium carbonate and 87 mg of N-ethylpropylamine were added, and the mixture was stirred at a bath temperature of 60°C for 6 hours. The reaction mixture was then left for cooling to room temperature and subjected to phase
20 separation three times between ethyl acetate and an aqueous sodium chloride solution. The organic layer was dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel thin layer chromatography to give 136 mg of 4-(2-butyneoxy)-6-(N-ethyl-N-propylamino)pyrimidine (the present compound (104)).

25 ¹H-NMR: 0.92 (t, 3H), 1.16 (t, 3H), 1.63 (m, 2H), 1.87 (t, 3H), 3.35 (t, 2H), 3.48 (q, 2H), 4.91 (q, 2H), 5.74 (s, 1H), 8.29 (s, 1H)

Production Example 103

To 2 ml of N,N-dimethylformamide were added 183 mg of 4-chloro-6-

(2-butynyloxy)pyrimidine, 166 mg of potassium carbonate, and 87 mg of N-ethylisopropylamine, followed by stirring at 60°C for 7 hours. Then, 166 mg of potassium carbonate and 435 mg of N-ethylisopropylamine were added, and the mixture was stirred at 80°C for 8 hours. Then, 166 mg of potassium carbonate and 435 mg of N-ethylisopropylamine were added, and the mixture was stirred at 120°C for 5 hours. The reaction mixture was then left for cooling to room temperature and subjected to phase separation three times between ethyl acetate and an aqueous sodium chloride solution. The organic layer was dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to preparative silica gel thin layer chromatography to give 79 mg of 4-(2-butynyloxy)-6-(N-ethyl-N-isopropylamino)pyrimidine (the present compound (105)).

¹H-NMR: 1.17 (t, 3H), 1.19 (d, 6H), 1.88 (t, 3H), 3.33 (q, 2H), 4.80 (br, 1H), 4.91 (q, 2H), 5.76 (s, 1H), 8.31 (s, 1H)

15 Production Example 104

To 4 ml of dimethylsulfoxide were added 365 mg of 4-chloro-6-(2-butynyloxy)pyrimidine, 332 mg of potassium carbonate, and 591 mg of isopropylamine, followed by stirring at 70°C for 6 hours. The reaction mixture was then left for cooling to room temperature, diluted with tert-butyl methyl ether, and washed twice with an aqueous sodium chloride solution. The organic layer was dried over anhydrous magnesium sulfate and concentrated. The residue was subjected to silica gel thin layer chromatography to give 339 mg of 4-(2-butynyloxy)-6-(isopropylamino)pyrimidine (the present compound (106)).

25 ¹H-NMR: 1.22 (d, 6H), 1.87 (t, 3H), 3.76 (m, 1H), 4.88 (br, 1H), 4.91 (q, 2H), 5.68 (s, 1H), 8.22 (s, 1H)

Production Example 105

In 5 ml of tetrahydrofuran were dissolved 164 mg of 4-(2-bu-

tynyloxy)-6-(isopropylamino)pyrimidine and 106 mg of 1-bromo-2-butyne, to which 40 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 6 hours. Then, ice water was added to the reaction mixture, which was extracted with ethyl acetate. The organic layer was washed with an aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to preparative silica gel thin layer chromatography to give 155 mg of 4-(2-butyloxy)-6-(N-(2-butyryl)-N-isopropylamino)pyrimidine (the present compound (107)).

¹H-NMR: 1.24 (d, 6H), 1.76 (t, 3H), 1.88 (t, 3H), 3.99 (d, 2H), 4.80 (br, 1H), 4.92 (q, 2H), 5.93 (s, 1H), 8.35 (s, 1H)

Production Example 106

To 20 ml of dimethylsulfoxide were added 1.83 g of 4-chloro-6-(2-butyloxy)pyrimidine, 2.20 g of potassium carbonate, and 20 ml of ethylamine (2.0 M tetrahydrofuran solution), followed by stirring at 50°C for 8 hours. The reaction mixture was then left for cooling to room temperature, diluted with tert-butyl methyl ether, and washed twice with an aqueous sodium chloride solution. The organic layer was dried over anhydrous magnesium sulfate and concentrated. The residue was recrystallized from tert-butyl methyl ether to give 1.14 g of 4-(2-butyloxy)-6-(ethylamino)pyrimidine (the present compound (108)).

¹H-NMR: 1.25 (t, 3H), 1.87 (t, 3H), 3.23 (m, 2H), 4.90 (br, 1H), 4.92 (q, 2H), 5.69 (s, 1H), 8.24 (s, 1H)

Production Example 107

In 5 ml of tetrahydrofuran were dissolved 153 mg of 4-(2-butyloxy)-6-(ethylamino)pyrimidine and 117 mg of 1-bromo-2-butyne, to which 40 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 7 hours. Then, 2 ml of N,N-dimethylformamide was fur-

ther added, and the mixture was stirred at room temperature for 3 hours. Then, ice water was added to the reaction mixture, which was extracted with ethyl acetate. The organic layer was washed twice with an aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to preparative silica gel thin layer chromatography to give 176 mg of 4-(2-butynyloxy)-6-(N-ethyl-N-(2-butyn-
5 yl)amino)pyrimidine (the present compound (109)).

¹H-NMR: 1.20 (t, 3H), 1.79 (t, 3H), 1.88 (t, 3H), 3.56 (q, 2H), 4.21 (d, 2H), 4.92 (q, 2H), 5.87 (s, 1H), 8.35 (s, 1H)

10 Production Example 108

In 2 ml of N,N-dimethylformamide were dissolved 153 mg of 4-(2-butynyloxy)-6-(ethylamino)pyrimidine and 67 mg of allyl chloride, to which 40 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 3 hours. Then, ice water was added to the reaction mixture,
15 which was extracted with ethyl acetate. The organic layer was washed twice with an aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to preparative silica gel thin layer chromatography to give 157 mg of 4-(2-butynyloxy)-6-(N-ethyl-N-allylamino)pyrimidine (the present compound (110)).

20 ¹H-NMR: 1.16 (t, 3H), 1.87 (t, 3H), 3.48 (q, 2H), 4.06 (d, 2H), 4.91 (q, 2H), 5.05-5.2 (m, 2H), 5.79 (s, 1H), 5.7-5.9 (m, 1H), 8.31 (s, 1H)

 Production Example 109

In 2 ml of tetrahydrofuran was suspended 0.05 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.07 g
25 of 2-butyn-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 0.6 ml of a tetrahydrofuran solution containing 0.2 g of 4-chloro-6-(1-(3-fluorophenyl)ethyl)pyrimidine was slowly added dropwise at room temperature, follow-

ed by stirring for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.21 g of 4-(2-butynyloxy)-6-(1-(3-fluorophenyl)ethyl)pyrimidine (the present compound (116)).

¹H-NMR: 1.61 (d, 3H), 1.85 (t, 3H), 4.13 (q, 1H), 4.96 (q, 2H), 6.58 (s, 1H), 6.88-7.04 (m, 3H), 7.22-7.30 (m, 1H), 8.74 (s, 1H)

10 Production Example 110

To 4 ml of dimethylsulfoxide were added 365 mg of 4-chloro-6-(2-butynyloxy)pyrimidine, 442 mg of potassium carbonate, and 495 mg of 2,2,3,3,3-pentafluoropropylamine, followed by stirring at 80°C for 4 hours. Then, 596 mg of 2,2,3,3,3-pentafluoropropylamine was further added, and the mixture was stirred at 100°C for 6 hours. The reaction mixture was then left for cooling to room temperature, diluted with ethyl acetate, and washed twice with an aqueous sodium chloride solution. The organic layer was dried over anhydrous magnesium sulfate and concentrated. The residue was subjected to silica gel column chromatography to give 73 mg of 4-(2-butynyloxy)-6-(2,2,3,3,3-pentafluoropropylamino)pyrimidine (the present compound (111)).

¹H-NMR: 1.87 (t, 3H), 4.12 (dt, 2H), 4.93 (q, 2H), 5.13 (br, 1H), 5.86 (s, 1H), 8.33 (s, 1H)

Production Example 111

25 In 2 ml of N,N-dimethylformamide were dissolved 73 mg of 4-(2-butynyloxy)-6-(2,2,3,3,3-pentafluoropropylamino)pyrimidine and 47 mg of ethyl iodide, to which 12 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 3.5 hours. Then, ice water was added to

the reaction mixture, which was extracted with ethyl acetate. The organic layer was washed twice with an aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to preparative silica gel thin layer chromatography to give 43 mg of
5 4-(2-butynyloxy)-6-(N-ethyl-N-(2,2,3,3,3-pentafluoropropyl)amino)pyrimidine (the present compound (112)).

¹H-NMR: 1.21 (t, 3H), 1.88 (t, 3H), 3.52 (q, 2H), 4.33 (q, 2H), 4.93 (q, 2H), 5.93 (s, 1H), 8.34 (s, 1H)

Production Example 112

10 To 2 ml of dimethylsulfoxide were added 183 mg of 4-chloro-6-(2-butynyloxy)pyrimidine, 166 mg of potassium carbonate, and 152 mg of dipropylamine, followed by stirring at 80°C for 8 hours. The reaction mixture was then left for cooling to room temperature, diluted with ethyl acetate, and washed twice with an aqueous sodium chloride solution. The organic layer
15 was dried over anhydrous magnesium sulfate and concentrated. The residue was subjected to silica gel thin layer chromatography to give 179 mg of 4-(2-butynyloxy)-6-(dipropylamino)pyrimidine (the present compound (113)).

¹H-NMR: 0.91 (t, 6H), 1.60 (m, 4H), 1.88 (t, 3H), 3.35 (t, 4H), 4.91 (q, 2H), 5.73 (s, 1H), 8.29 (s, 1H)

20 Production Example 113

To 2 ml of dimethylsulfoxide were added 183 mg of 4-chloro-6-(2-butynyloxy)pyrimidine, 220 mg of potassium carbonate, and 495 mg of 2,2,2-trifluoroethylamine, followed by stirring at a bath temperature of 60°C for 9 hours. Then, 495 mg of 2,2,2-trifluoroethylamine was further added, and
25 the mixture was stirred at 80°C for 8 hours. Then, 495 mg of 2,2,2-trifluoroethylamine and 2 ml of dimethylsulfoxide were further added, and the mixture was stirred at 80°C for 8 hours. The reaction mixture was then left for cooling to room temperature, diluted with ethyl acetate, and washed

twice with an aqueous sodium chloride solution. The organic layer was dried over anhydrous magnesium sulfate and concentrated. The residue was subjected to silica gel thin layer chromatography to give 60 mg of 4-(2-butynyloxy)-6-(2,2,2-trifluoroethylamino)pyrimidine (the present compound
5 (114)).

¹H-NMR: 1.87 (t, 3H), 4.08 (dq, 2H), 4.93 (q, 2H), 5.02 (br, 1H), 5.85 (s, 1H), 8.39 (s, 1H)

Production Example 114

In 2 ml of N,N-dimethylformamide were dissolved 40 mg of 4-(2-butynyloxy)-6-(2,2,2-trifluoroethylamino)pyrimidine and 31 mg of ethyl iodide,
10 to which 8 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 1 hour. Then, 6 ml of tetrahydrofuran was added, and the mixture was further stirred at room temperature for 4.5 hours. Then, ice water was added to the reaction mixture, from which the tetra-
15 hydrofuran was distilled out under reduced pressure. The residue was extracted with tert-butyl methyl ether. The organic layer was washed twice with an aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel thin layer chromatography to give 37 mg of 4-(2-butynyloxy)-6-(N-ethyl-N-2,2,2-
20 trifluoroethylamino)pyrimidine (the present compound (115)).

¹H-NMR: 1.20 (t, 3H), 1.88 (t, 3H), 3.51 (q, 2H), 4.24 (q, 2H), 4.93 (q, 2H), 5.93 (s, 1H), 8.35 (s, 1H)

Production Example 115

In 9 ml of N,N-dimethylformamide were dissolved 202 mg of 4-chloro-6-(2,3-difluorophenyl)pyrimidine and 112 mg of 2-pentyn-1-ol, to which 54
25 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 5 hours. The reaction mixture was then poured into water and extracted with ethyl acetate. The organic layer was washed with a

saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 211 mg of 4-(2,3-difluorophenyl)-6-(2-pentynyloxy)pyrimidine (the present compound (117)).

5 ¹H-NMR: 1.19 (t, 3H), 2.28 (q, 2H), 5.06 (t, 2H), 7.15-7.34 (m, 2H), 7.30 (s, 1H), 7.86 (t, 1H), 8.88 (s, 1H)

Production Example 116

In 9 ml of N,N-dimethylformamide were dissolved 217 mg of 4-chloro-6-(2,3-difluorophenyl)pyrimidine and 141 mg of 2-hexyn-1-ol, to which 58
10 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 7 hours. The reaction mixture was then poured into water and extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected
15 to silica gel column chromatography to give 247 mg of 4-(2,3-difluorophenyl)-6-(2-hexynyloxy)pyrimidine (the present compound (118)).

¹H-NMR: 0.98 (t, 3H), 1.55 (sextet, 2H), 2.23 (quintet, 2H), 5.06 (t, 2H), 7.15-7.34 (m, 2H), 7.30 (s, 1H), 7.87 (t, 1H), 8.88 (s, 1H)

Production Example 117

20 In 6 ml of N,N-dimethylformamide were dissolved 199 mg of 4-chloro-6-(2,3-difluorophenyl)pyrimidine and 118 mg of 4,4-dimethyl-2-pentyn-1-ol, to which 43 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 3 hours. The reaction mixture was then poured into water and extracted with ethyl acetate. The organic layer was
25 washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 190 mg of 4-(4,4-dimethyl-2-pentynyloxy)-6-(2,3-difluorophenyl)pyrimidine (the present com-

pound (119)).

¹H-NMR: 1.22 (s, 9H), 5.05 (s, 2H), 7.14-7.35 (m, 2H), 7.30 (s, 1H), 7.88 (t, 1H), 8.86 (s, 1H)

Production Example 122

5 In 1 ml of tetrahydrofuran was suspended 0.02 g of sodium hydride (60% in oil), to which 0.3 ml of a tetrahydrofuran solution containing 0.02 g of 2-butyn-1-ol was slowly added dropwise with stirring at room temperature. The mixture was then stirred at room temperature for 20 minutes, to which 10 0.3 ml of a tetrahydrofuran solution containing 0.06 g of 4-chloro-6-(1-(2-fluorophenyl)ethyl)pyrimidine was slowly added dropwise at room temperature, followed by stirring for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. 15 The residue was subjected to silica gel column chromatography to give 0.06 g of 4-(2-butynyloxy)-6-(1-(2-fluorophenyl)ethyl)pyrimidine (the present compound (120)).

¹H-NMR: 1.66 (d, 3H), 1.86 (t, 3H), 4.45 (q, 1H), 4.95 (q, 2H), 6.60 (s, 1H), 6.98-7.36 (m, 4H), 8.74 (s, 1H)


20 Production Example 123

 In 2 ml of tetrahydrofuran was suspended 0.04 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.06 g of 2-butyn-1-ol was slowly added dropwise with stirring at room temperature. The mixture was then stirred at room temperature for 20 minutes, to which 25 0.6 ml of a tetrahydrofuran solution containing 0.2 g of 4-chloro-6-(2-chloro-5-methyl-6-fluorobenzyl)pyrimidine was slowly added dropwise at room temperature, followed by stirring for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted

three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.24 g of 4-(2-butynyloxy)-6-(2-chloro-5-methyl-6-fluorobenzyl)pyrimidine (the present compound (121)).

¹H-NMR: 1.86 (t, 3H), 2.35 (s, 3H), 4.26 (s, 2H), 4.94 (q, 2H), 6.41 (s, 1H), 6.95 (t, 1H), 7.16 (dd, 1H), 8.73 (s, 1H)

Production Example 124




In 2 ml of tetrahydrofuran was suspended 0.04 g of sodium hydride, to which 0.06g of 2-butyne-1-ol dissolved in 0.3 ml of tetrahydrofuran was added at room temperature. The mixture was then stirred for 15 minutes, to which 0.2 g of 4-(α -methoxyphenylmethyl)-6-methanesulfonylpyrimidine dissolved in 0.3 ml of tetrahydrofuran was added dropwise under ice cooling, followed by further stirring for 30 minutes. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.14 g of 4-(2-butynyloxy)-6-(α -methoxybenzyl)pyrimidine (the present compound (122)).

¹H-NMR: 1.86 (t, 3H), 3.41 (s, 3H), 4.97 (q, 2H), 5.21 (s, 1H), 7.02 (s, 1H), 7.28-7.42 (m, 5H), 8.69 (s, 1H)

Production Example 125


156



In 2 ml of ethanol was suspended 0.5 g of 6-(2-butynyloxy)-4-benzoylpyrimidine, to which 0.11 g of sodium borohydride was added, followed by stirring at 0°C for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with ethyl acetate. The organic layers were combined and washed with a saturated aqueous sodium chloride solution, and the combined organic layer was then dried over anhydrous magnesium sulfate and concentrated. The residue was subjected to silica gel column chromatography to give 0.4 g of 6-(2-butynyloxy)-4-(α-hydroxybenzyl)pyrimidine (the present compound (123)).

¹H-NMR: 1.84 (t, 3H), 4.58 (bs, 1H), 4.95 (q, 2H), 5.63 (s, 1H), 6.72 (s, 1H), 7.28-7.38 (m, 5H), 8.71 (s, 1H)

Production Example 126




In 4 ml of acetonitrile was dissolved 0.26 g of 6-(2-butynyloxy)-4-(α-hydroxybenzyl)pyrimidine, to which a solution of 0.17 g of 2,2-difluoro-1,3-dimethylimidazolidine in 1.5 ml of acetonitrile was added, followed by stirring at room temperature for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with ethyl acetate. The organic layers were combined and washed with a saturated aqueous sodium chloride solution, and the combined organic layer was dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.14 g of 4-(2-butynyloxy)-6-(α-fluorobenzyl)pyrimidine (the

present compound (124)).

¹H-NMR: 1.87 (t, 3H), 5.00 (q, 2H), 6.35 (d, 1H), 7.07 (s, 1H), 7.35-7.44 (m, 5H), 8.72 (s, 1H)

Production Example 127



In 3 ml of pyridine were added 0.3 g of 6-(2-butynyloxy)-4-benzoylpyrimidine and 0.15 g of O-methylhydroxylamine hydrochloride, followed by stirring at room temperature for 3 hours. The reaction mixture was then poured into 10% hydrochloric acid and extracted three times with ethyl acetate. The organic layers were combined and washed with a saturated aqueous sodium chloride solution, and the combined organic layer was dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give two isomers of (6-(2-butynyloxy)-4-pyrimidyl)phenylketone O-methyloxime (referred to as isomers A and B).

10

15

Isomer A (the present compound (125)): 0.25 g

Isomer B (the present compound (126)): 0.07 g

Isomer A

¹H-NMR: 1.89 (t, 3H), 3.98 (s, 3H), 5.04 (q, 2H), 6.93 (s, 1H), 7.32-7.48 (m, 5H), 8.90 (s, 1H)


20

Isomer B

¹H-NMR: 1.87 (t, 3H), 4.05 (s, 3H), 4.99 (q, 2H), 6.98 (s, 1H), 7.32-7.36 (m, 2H), 7.41-7.48 (m, 3H), 8.81 (s, 1H)

Production Example 128

158



In 3 ml of pyridine were added 0.3 g of 6-(2-butynyloxy)-4-benzoylpyrimidine and 0.17 g of O-ethylhydroxylamine hydrochloride, followed by stirring at room temperature for 3 hours. The reaction mixture was then poured into 10% hydrochloric acid and extracted three times with ethyl acetate. The organic layers were combined and washed with a saturated aqueous sodium chloride solution, and the combined organic layer was dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give two isomers of (6-(2-butynyloxy)-4-pyrimidyl)phenylketone O-ethyloxime (referred to as isomers A and B).

Isomer A (the present compound (127)): 0.20 g

Isomer B (the present compound (128)): 0.12 g

Geometrical Isomer A

¹H-NMR: 1.29 (t, 3H), 1.88 (t, 3H), 4.24 (q, 2H), 5.05 (q, 2H), 6.97 (s, 1H), 7.29-7.34 (m, 3H), 7.45-7.48 (m, 2H), 8.90 (s, 1H)

Geometrical Isomer B

¹H-NMR: 1.31 (t, 3H), 1.86 (t, 3H), 4.32 (q, 2H), 4.99 (q, 2H), 7.05 (s, 1H), 7.34-7.46 (m, 5H), 8.80 (s, 1H)

Production Example 129

To 4 ml of ethanol were added 0.5 g of 4-chloro-6-(2-butynyloxy)pyrimidine and 0.68 g of N-ethyl-N-benzylamine, followed by heating under reflux for 6 hours. The reaction mixture was then left for cooling to room temperature and concentrated under reduced pressure. A saturated aqueous ammonium chloride solution was poured onto the residue, which was extracted three times with ethyl acetate. The organic layers were

combined and washed with a saturated aqueous sodium chloride solution, and the combined organic layer was dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.42 g of 6-(2-butynyloxy)-4-(N-ethyl-N-benzyl-
5 amino)pyrimidine (the present compound (129)).


¹H-NMR: 1.14 (t, 3H), 1.84 (t, 3H), 3.29 (q, 2H), 4.71 (s, 2H), 4.91 (q, 2H), 5.78 (s, 1H), 7.17-7.32 (m, 5H), 8.34 (s, 1H)

Production Example 130

In 2 ml of tetrahydrofuran was suspended 0.04 g of sodium hydride
10 (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.06 g of 2-butyne-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 0.6 ml of a tetrahydrofuran solution containing 0.2 g of 4-chloro-6-(2-chloro-3,6-difluorobenzyl)pyrimidine was slowly added dropwise at room temperature,
15 followed by stirring for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.22 g of 4-(2-
20 butynyloxy)-6-(2-chloro-3,6-difluorobenzyl)pyrimidine (the present compound (130)).

¹H-NMR: 1.86 (t, 3H), 4.25 (s, 2H), 4.95 (q, 2H), 6.46 (s, 1H), 6.99-7.14 (m, 2H), 8.72 (s, 1H)

Production Example 131



In 3 ml of pyridine were added 0.3 g of 6-(2-butynyloxy)-4-benzoyl-

pyrimidine and 0.20 g of O-isopropylhydroxylamine hydrochloride, followed by stirring at room temperature for 3 hours. The reaction mixture was then poured into 10% hydrochloric acid and extracted three times with ethyl acetate. The organic layers were combined and washed with a saturated aqueous sodium chloride solution, and the combined organic layer was dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give two isomers of 6-(2-butynyloxy)-4-pyrimidylphenylketone O-isopropoxyime (referred to as isomers A and B).

10 Isomer A (the present compound (131)): 0.26 g

Isomer B (the present compound (132)): 0.15 g

Isomer A

¹H-NMR: 1.26 (d, 6H), 1.89 (t, 3H), 4.43-4.55 (m, 1H), 5.05 (q, 2H), 6.99 (s, 1H), 7.30-7.36 (m, 3H), 7.44-7.48 (m, 2H), 8.89 (s, 1H)

15 Isomer B

¹H-NMR: 1.29 (d, 6H), 1.87 (t, 3H), 4.50-4.61 (m, 1H), 4.99 (q, 2H), 7.12 (s, 1H), 7.37-7.41 (m, 5H), 8.79 (s, 1H)

Production Example 132

In 8 ml of ethanol were added 1 g of 4-chloro-6-(2-butynyloxy)pyrimidine and 1.17 g of benzylamine, followed by heating under reflux for 6 hours. The reaction mixture was then left for cooling to room temperature and concentrated under reduced pressure. A saturated aqueous ammonium chloride solution was poured onto the residue, which was extracted three times with ethyl acetate. The organic layers were combined and washed with a saturated aqueous sodium chloride solution, and the combined organic layer was dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 1.58 g of 6-(2-butynyloxy)-4-benzylaminopyrimidine (the present compound (133)).

¹H-NMR: 1.86 (t, 3H), 4.43 (d, 2H), 4.90 (q, 2H), 5.38 (bs, 1H), 5.70 (s, 1H), 7.28-7.37 (m, 5H), 8.25 (s, 1H)

Production Example 133

In 6 ml of N,N-dimethylformamide were dissolved 150 mg of 4-chloro-6-(2,3-difluorophenyl)pyrimidine and 56 mg of 3-butyn-2-ol, to which 32 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 12 hours. The reaction mixture was then poured into water and extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 120 mg of 4-(2,3-difluorophenyl)-6-(1-methyl-2-propynyloxy)pyrimidine (the present compound (134)).

¹H-NMR: 1.68 (d, 3H), 2.49 (t, 1H), 5.89 (dq, 1H), 7.17-7.42 (m, 2H), 7.82-7.92 (m, 1H), 7.89 (s, 1H), 8.89 (s, 1H)

15 Production Example 134

In 6 ml of acetonitrile was dissolved 255 mg of 4-(2,3-difluorophenyl)-6-(4-hydroxy-2-butynyloxy)pyrimidine, to which 148 mg of 2,2-difluoro-1,3-dimethylimidazolidine was added, followed by stirring at room temperature for 24 hours. The reaction mixture was then poured into water and extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 43 mg of 4-(2,3-difluorophenyl)-6-(4-fluoro-2-butynyloxy)pyrimidine (the present compound (135)).

-25 ¹H-NMR: 5.04 (dt, 2H), 5.17 (dd, 2H), 7.12-7.51 (m, 3H), 7.33 (s, 1H), 8.14 (t, 1H), 8.90 (s, 1H)


Production Example 135

In 5 ml of carbon tetrachloride was dissolved 241 mg of 4-(2,3-difluo-

ropheryl)-6-(2-propynyloxy)pyrimidine, to which 136 mg of potassium carbonate and 136 mg of tetra-n-butylammonium chloride were added, followed by stirring at room temperature for 16 hours. The reaction mixture was then poured into water and extracted with ethyl acetate. The organic layer
5 was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 159 mg of 4-(2,3-difluorophenyl)-6-(3-chloro-2-propynyloxy)pyrimidine (the present compound (136)).

10 ¹H-NMR: 5.09 (s, 2H), 7.16-7.34 (m, 3H, involving a singlet at 7.29), 7.88 (t, 1H), 8.90 (s, 1H)

Production Example 136




In 1.5 ml of tetrahydrofuran was suspended 0.05 g of sodium hydride
15 (60% in oil), to which 0.2 g of 6-(2-butynyloxy)-4-(α -hydroxybenzyl)pyrimidine was added under ice cooling, followed by stirring for 15 minutes. Then, 0.2 ml of a tetrahydrofuran solution containing 0.08 g of chloromethyl methyl ether was slowly added dropwise, followed by further stirring at the same temperature for 2 hours. The reaction mixture was then poured into a
20 saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.25 g of 6-(2-butynyloxy)-4-(α -methoxymethoxybenzyl)pyrimidine (the present compound
25 (137)).

¹H-NMR: 1.86 (t, 3H), 3.36 (s, 3H), 4.71 (dd, 2H), 4.98 (q, 2H), 5.67 (s,

1H), 7.08 (s, 1H), 7.27-7.43 (m, 5H), 8.70 (s, 1H)

Production Example 137



In 1.5 ml of tetrahydrofuran was suspended 0.05 g of sodium hydride
5 (60% in oil), to which 0.2 g of 6-(2-butynyloxy)-4-(α -hydroxybenzyl)pyrimidine was added under ice cooling, followed by stirring for 15 minutes. Then, 0.2 ml of a tetrahydrofuran solution containing 0.18 g of ethyl iodide was slowly added dropwise, followed by further stirring at the same temperature for 2 hours. The reaction mixture was then poured into a saturated aque-
10 ous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.07 g of 6-(2-butynyloxy)-4-(α -ethoxybenzyl)pyrimidine (the present compound (138)).


15 $^1\text{H-NMR}$: 1.27 (t, 3H), 1.86 (t, 3H), 3.55 (q, 2H), 4.97 (q, 2H), 5.32 (s, 1H), 7.07 (s, 1H), 7.27-7.43 (m, 5H), 8.68 (s, 1H)

Production Example 138

To 1.8 ml of tetrahydrofuran were added 0.15 ml of diisopropylethyl-
amine and 0.15 g of 6-(2-butynyloxy)-4-(α -hydroxybenzyl)pyrimidine under
20 ice cooling. Then, 0.2 ml of a tetrahydrofuran solution containing 0.06 g of acetyl chloride was slowly added dropwise, followed by further stirring at the same temperature for 3 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times
25 with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.18 g of the

164


present compound (139) of the following formula



¹H-NMR: 1.86 (t, 3H), 2.19 (s, 3H), 4.98 (q, 2H), 6.70 (s, 1H), 6.90 (s, 1H), 7.30-7.43 (m, 5H), 8.72 (s, 1H)

5 Production Example 139

To 1.8 ml of tetrahydrofuran were added 0.15 ml of diisopropylethylamine and 0.15 g of 6-(2-butynyloxy)-4-(α-hydroxybenzyl)pyrimidine under ice cooling. Then, 0.2 ml of a tetrahydrofuran solution containing 0.07 g of propionyl chloride was slowly added dropwise, and after completion of the dropwise addition, the mixture was stirred at the same temperature for 3 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.12 g of the present compound (140) of the following formula




¹H-NMR: 1.19 (t, 3H), 1.86 (t, 3H), 2.58 (q, 2H), 4.97 (q, 2H), 6.71 (s, 1H), 6.90 (s, 1H), 7.29-7.44 (m, 5H), 8.72 (s, 1H)

20 Production Example 140

To 1.8 ml of tetrahydrofuran were added 0.15 ml of diisopropylethylamine and 0.15 g of 6-(2-butynyloxy)-4-(α-hydroxybenzyl)pyrimidine under ice cooling. Then, 0.2 ml of a tetrahydrofuran solution containing 0.08 g of isobutyryl chloride was slowly added dropwise, and after completion of the

dropwise addition, the mixture was stirred at the same temperature for 3 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.21 g of the present compound (141) of the following formula



of 4-chloro-6-(4-fluorobenzyl)pyrimidine was slowly added dropwise, followed by further stirring at room temperature for 2 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.22 g of 4-(2-butynyloxy)-6-(4-fluorobenzyl)pyrimidine (the present compound (143)).

¹H-NMR: 1.85 (t, 3H), 3.99 (s, 2H), 4.95 (q, 2H), 6.51 (s, 1H), 6.97-7.03 (m, 2H), 7.19-7.23 (m, 2H), 8.72 (s, 1H)

Production Example 143

To 2 ml of N,N-dimethylformamide were added 183 mg of 4-chloro-6-(2-butynyloxy)pyrimidine, 166 mg of potassium carbonate, and 73 mg of diethylamine, followed by stirring at a bath temperature of 50°C for 4 hours. Then, 73 mg of diethylamine was further added, followed by stirring at a bath temperature of 50°C for 5.5 hours. Then, 146 mg of diethylamine was further added, followed by stirring at a bath temperature of 40°C for 6 hours. The reaction mixture was left for cooling to room temperature, diluted with ethyl acetate, and washed three times with an aqueous sodium chloride solution. The organic layers were dried over anhydrous magnesium sulfate and concentrated. The residue was subjected to silica gel thin layer chromatography to give 136 mg of 4-(2-butynyloxy)-6-(N,N-diethylamino)pyrimidine (the present compound (144)).

¹H-NMR: 1.17 (t, 6H), 1.88 (t, 3H), 3.47 (q, 4H), 4.91 (q, 2H), 5.75 (s, 1H), 8.30 (s, 1H)

Production Example 144

In 40 ml of tetrahydrofuran was suspended 4.61 g of potassium t-butoxide, to which 4.44 g of (2-fluorophenyl)acetonitrile was added under ice

cooling. Then, a solution of 5.00 g of 4-chloro-6-(2-butynyloxy)pyrimidine in 20 ml of tetrahydrofuran was added at 0°C, followed by stirring at room temperature for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times
5 with chloroform. The chloroform layers were combined and washed with a saturated aqueous sodium chloride solution. The organic layer was dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 6.40 g of 4-(α -cyano-2-fluorobenzyl)-6-(2-butynyloxy)pyrimidine (the present compound
10 (145)).

¹H-NMR: 1.87 (t, 3H), 4.98 (q, 2H), 5.44 (s, 1H), 6.83 (s, 1H), 7.11-7.49 (m, 4H), 8.78 (s, 1H)

Production Example 145

To 2.2 ml of chloroform were added 0.29 ml of diisopropylethylamine, 0.2 g of 4-(2-butynyloxy)-6-chloropyrimidine, and 0.17 g of 2-fluorothiophenol,
15 followed by stirring at room temperature for 7 hours. Then, a saturated aqueous ammonium chloride solution was poured into the reaction mixture, which was extracted three times with t-butyl methyl ether. The organic layers were combined, washed with water, dried over anhydrous magnesium
20 sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.07 g of 4-(2-butynyloxy)-6-(2-fluorophenylthio)pyrimidine (the present compound (146)).

¹H-NMR: 1.85 (t, 3H), 4.92 (q, 2H), 6.21 (s, 1H), 7.20-7.28 (m, 2H), 7.48-7.63 (m, 2H), 8.55 (s, 1H)


25 Production Example 146

In 1 ml of tetrahydrofuran was suspended 0.04 g of sodium hydride (60% in oil), to which 0.2 ml of a tetrahydrofuran solution containing 0.06 g of 3-pentyn-2-ol was slowly added dropwise under stirring at room tempera-

ture. The mixture was stirred at room temperature for 20 minutes, and 0.4 ml of a tetrahydrofuran solution containing 0.15 g of 4-chloro-6-(2,6-difluorobenzyl)pyrimidine was slowly added dropwise at room temperature, followed by stirring for 3 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with ethyl acetate. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.24 g of 4-(2,6-difluorobenzyl)-6-(1-methyl-2-butyloxy)pyrimidine (the present compound (147)).

¹H-NMR: 1.56 (d, 3H), 1.82 (t, 3H), 4.11 (s, 2H), 5.75-5.79 (m, 1H), 6.47 (s, 1H), 6.87-6.96 (m, 2H), 7.19-7.29 (m, 2H), 8.72 (s, 1H)


Production Example 147



In 3.2 ml of tetrahydrofuran was dissolved 0.4 g of 6-(2-butyloxy)-4-benzoylpyrimidine, to which 1.67 ml (1.14 mol/l diethyl ether solution) of methyl lithium was added dropwise at 0°C. After stirring at room temperature for 2 hours, the reaction mixture was poured into a saturated aqueous ammonium chloride solution and extracted three times with ethyl acetate. The organic layers were combined and washed with a saturated aqueous sodium chloride solution, and the combined organic layer was dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.38 g of 4-(2-butyloxy)-6-(α-hydroxy-α-methylbenzyl)pyrimidine (the present compound (148)).

¹H-NMR: 1.85 (t, 3H), 1.88 (s, 3H), 4.80 (s, 1H), 4.96 (q, 2H), 6.79 (s, 1H), 7.25 (t, 1H), 7.29 (t, 2H), 7.47 (d, 2H), 8.73 (s, 1H)

Production Example 148



To a solution of 0.2 g of 6-(2-butynyloxy)-4-(α -hydroxy-2-fluorobenzyl)pyrimidine in 3.6 ml of acetonitrile was added dropwise a solution of 0.12 g of 2,2-difluoro-1,3-dimethylimidazolidine in 1 ml of acetonitrile, followed by stirring at room temperature for 4 hours. The reaction mixture was then poured into water and extracted three times with ethyl acetate. The organic layers were combined and washed with a saturated aqueous sodium chloride solution, and the combined organic layer was dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.08 g of 4-(2-butynyloxy)-6-(α -fluoro-2-fluorobenzyl)pyrimidine (the present compound (149)).

$^1\text{H-NMR}$: 1.88 (t, 3H), 5.00 (q, 2H), 6.64 (d, 1H), 7.06-7.41 (m, 5H, involving a singlet at 7.10), 8.73 (s, 1H)

15 Production Example 149

In 10 ml of N,N-dimethylformamide were dissolved 458 mg of 4-chloro-6-(2-fluorophenyl)pyrimidine and 231 mg of 3-butyn-2-ol, to which 132 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 10 hours. The reaction mixture was then poured into water and extracted three times with ethyl acetate. The organic layers were combined, washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 506 mg of 4-(2-fluorophenyl)-6-(1-methyl-2-propynyloxy)pyrimidine (the present compound (150)).

$^1\text{H-NMR}$: 1.68 (d, 3H), 2.49 (d, 1H), 5.89 (dq, 1H), 7.13-7.23 (m, 1H),

7.25-7.34 (m, 2H, involving a singlet at 7.30), 7.39-7.49 (m, 1H), 8.07-8.16 (m, 1H), 8.89 (s, 1H)

Production Example 150

In 10 ml of N,N-dimethylformamide were dissolved 449 mg of 4-chloro-6-(2-fluorophenyl)pyrimidine and 272 mg of 3-pentyn-2-ol, to which 130 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 8 hours. The reaction mixture was then poured into water and extracted three times with ethyl acetate. The organic layers were combined, washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 510 mg of 4-(2-fluorophenyl)-6-(1-methyl-2-butyloxy)pyrimidine (the present compound (151)).

¹H-NMR: 1.63 (d, 3H), 2.32 (d, 3H), 5.87 (m, 1H), 7.03-7.21 (m, 1H), 7.25-7.32 (m, 2H, involving a singlet at 7.31), 7.39-7.49 (m, 1H), 8.07-8.16 (m, 1H), 8.88 (s, 1H)

Production Example 151

In 10 ml of N,N-dimethylformamide were dissolved 450 mg of 4-chloro-6-(2-fluorophenyl)pyrimidine and 227 mg of 3-buten-1-ol, to which 130 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 12 hours. The reaction mixture was then poured into water and extracted three times with ethyl acetate. The organic layers were combined, washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 302 mg of 4-(2-fluorophenyl)-6-(3-butyloxy)pyrimidine (the present compound (152)).

¹H-NMR: 2.06 (t, 1H), 2.72 (dt, 2H), 4.54 (t, 2H), 7.12-7.22 (m, 1H), 7.27 (s, 1H), 7.26-7.32 (m, 1H), 7.40-7.49 (m, 1H), 8.08-8.16 (m, 1H), 8.84 (s,


1H)

Production Example 152

In 10 ml of N,N-dimethylformamide were dissolved 457 mg of 4-chloro-6-(2-fluorophenyl)pyrimidine and 277 mg of 3-pentyn-1-ol, to which 132 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 8 hours. The reaction mixture was then poured into water and extracted three times with ethyl acetate. The organic layers were combined, washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 467 mg of 4-(2-fluorophenyl)-6-(3-pentynyloxy)pyrimidine (the present compound (153)).

¹H-NMR: 1.79 (t, 3H), 2.65 (m, 2H), 4.15 (t, 2H), 7.01-7.20 (m, 1H), 7.23-7.30 (m, 1H), 7.38-7.48 (m, 1H), 7.27 (s, 1H), 8.07-8.16 (m, 1H), 8.83 (s, 1H)

15 Production Example 153



To a solution of 0.33 g of 6-(2-butynyloxy)-4-(α-hydroxy-α-methylbenzyl)pyrimidine in 4 ml of acetonitrile was added dropwise a solution of 0.2 g of 2,2-difluoro-1,3-dimethylimidazolidine in 2 ml of acetonitrile, followed by stirring at room temperature for 6 hours. The reaction mixture was then poured into water and extracted three times with ethyl acetate. The organic layers were combined and washed with a saturated aqueous sodium chloride solution, and the combined organic layer was dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.20 g of 4-(2-butynyloxy)-6-(α-fluoro-α-methylbenzyl)pyrimidine (the present compound (154)) and 0.09 g of

4-(2-butynyloxy)-6-(1-phenylvinyl)pyrimidine (the present compound (155)).

The present compound (154):

¹H-NMR: 1.85 (t, 3H), 2.05 (d, 3H), 4.97 (q, 2H), 7.07 (d, 1H), 7.25-7.36 (m, 3H), 7.50-7.53 (m, 2H), 8.73 (s, 1H)

5 The present compound (155):


¹H-NMR: 1.86 (t, 3H), 4.98 (q, 2H), 5.66 (d, 1H), 6.03 (d, 1H), 6.65 (s, 1H), 7.29-7.37 (m, 5H), 8.80 (s, 1H)

Production Example 154

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.23 g of potassium carbonate and 0.19 g of 2-chloro-5-fluorophenol, followed by stirring at 60°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined and washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.30 g of 4-(2-butynyloxy)-6-(2-chloro-5-fluorophenoxy)pyrimidine (the present compound (156)).

20 ¹H-NMR: 1.88 (t, 3H), 5.00 (q, 2H), 6.33 (s, 1H), 6.94-7.00 (m, 2H), 7.41-7.46 (m, 1H), 8.44 (s, 1H)

Production Example 155



In 3.7 ml of tetrahydrofuran was dissolved 0.4 g of 6-(2-butynyloxy)-pyrimidin-4-yl 2-fluorophenyl ketone, to which 3.6 ml (1.14 mol/l diethyl ether solution) of methyl lithium was added dropwise at -78°C. After stirring at -78°C for 3 hours, the reaction mixture was poured into a saturated

aqueous ammonium chloride solution and extracted three times with ethyl acetate. The organic layers were combined and washed with a saturated aqueous sodium chloride solution, and the combined organic layer was dried over anhydrous magnesium sulfate and then concentrated. The residue
5 was subjected to silica gel column chromatography to give 0.34 g of 4-(2-butynyloxy)-6-(α -hydroxy- α -methyl-2-fluorobenzyl)pyrimidine (the present compound (157)).

¹H-NMR: 1.86 (t, 3H), 1.91 (s, 3H), 4.85 (s, 1H), 4.98 (q, 2H), 6.88 (s, 1H), 6.95 (dd, 1H), 7.13-7.30 (m, 2H), 7.70 (td, 1H), 8.71 (s, 1H)

10 Production Example 156

Under an atmosphere of a nitrogen gas, 400 mg of sodium hydride was added to 20 ml of N,N-dimethylformamide, followed by ice cooling, to which an N,N-dimethylformamide (4 ml) solution of 861 mg of cyclopentyl alcohol was added dropwise, and the mixture was stirred for 1 hour. Under
15 ice cooling, 1.49 g of 4,6-dichloropyrimidine was added, and the mixture was stirred at room temperature for 6 hours. The reaction mixture was then poured into water, which was extracted twice with t-butyl methyl ether and then washed three times with water. The organic layers were dried over anhydrous magnesium sulfate and concentrated to give 950 mg of the crude
20 product.

Under an atmosphere of a nitrogen gas, 240 mg of sodium hydride was added to 10 ml of N,N-dimethylformamide, followed by ice cooling, to which an N,N-dimethylformamide (2 ml) solution of 421 mg of 2-butyne-1-ol was added dropwise, and the mixture was stirred at room temperature for 30
25 minutes. After ice cooling, an N,N-dimethylformamide (2 ml) solution of the crude product obtained above was added dropwise. The mixture was stirred at room temperature for 1 hour and then at a bath temperature of 50°C for 4 hours. After left for cooling to room temperature, the reaction mixture was

poured into water. The mixture was extracted three times with t-butyl methyl ether and then washed three times with water. The organic layers were dried over anhydrous magnesium sulfate and concentrated. The residue was subjected to silica gel thin layer chromatography to give 396 mg of
5 4-(2-butynyloxy)-6-(cyclopentyloxy)pyrimidine (the present compound (158)).

¹H-NMR: 1.55-2.05 (m, 11H), 4.94 (q, 2H), 5.35 (m, 1H), 6.05 (s, 1H), 8.43 (s, 1H)

Production Example 157

Under an atmosphere of a nitrogen gas, 240 mg of sodium hydride
10 was added to 10 ml of N,N-dimethylformamide, followed by ice cooling, to which an N,N-dimethylformamide (2 ml) solution of 601 mg of cyclohexyl alcohol was added dropwise, and the mixture was stirred for 6 hours. Under ice cooling, 894 mg of 4,6-dichloropyrimidine was added, and the mixture was stirred at room temperature overnight. The reaction mixture was then
15 poured into water, which was extracted twice with t-butyl methyl ether and then washed three times with water. The organic layers were dried over anhydrous magnesium sulfate and concentrated to give 450 mg of the crude product.

Under an atmosphere of a nitrogen gas, 160 mg of sodium hydride
20 was added to 4 ml of N,N-dimethylformamide, followed by ice cooling, to which an N,N-dimethylformamide (2 ml) solution of 280 mg of 2-butyne-1-ol was added dropwise, and the mixture was stirred at room temperature for 1 hour. After ice cooling, an N,N-dimethylformamide (2 ml) solution of the crude product obtained above was added dropwise. After stirring at room
25 temperature for 6 hours, the reaction mixture was poured into water. The mixture was extracted two times with t-butyl methyl ether and then washed four times with water. The organic layers were dried over anhydrous magnesium sulfate and concentrated. The residue was subjected to silica gel

thin layer chromatography to give 50 mg of 4-(2-butynyloxy)-6-(cyclohexyloxy)pyrimidine (the present compound (159)).

¹H-NMR: 1.20-1.65 (m, 6H), 1.77 (m, 2H), 1.87 (t, 3H), 1.96 (m, 2H), 4.94 (q, 2H), 5.01 (m, 1H), 6.05 (s, 1H), 8.42 (s, 1H)

5 Production Example 158

To 2 ml of N,N-dimethylformamide were added 0.2 g of 4-chloro-6-(2-butynyloxy)pyrimidine, 0.23 g of potassium carbonate, and 0.20 g of 2,6-dichloro-4-fluorophenol, followed by stirring at 80°C for 7 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with chloroform. The chloroform layers were combined, washed with diluted hydrochloric acid and then with water, and dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 0.22 g of 4-(2-butynyloxy)-6-(2,6-dichloro-4-fluorophenoxy)pyrimidine (the present compound (160)).

¹H-NMR: 1.89 (t, 3H), 5.00 (q, 2H), 6.40 (s, 1H), 7.17 (d, 2H), 8.40 (s, 1H)

 Production Example 159

In 10 ml of N,N-dimethylformamide were dissolved 255 mg of 4-chloro-2-methyl-6-(2,3-difluorophenyl)pyrimidine and 97 mg of 3-butyn-2-ol, to which 56 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 9 hours. The reaction mixture was then poured into water and extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 250 mg of 2-methyl-4-(2,3-difluorophenyl)-6-(1-methyl-2-propynyloxy)pyrimidine (the present compound (161)).


¹H-NMR: 1.67 (d, 3H), 2.48 (d, 1H), 2.69 (s, 3H), 5.94 (dq, 1H), 7.06 (s, 1H), 7.13-7.30 (m, 2H), 7.79-7.86 (m, 1H)

Production Example 160

In 10 ml of N,N-dimethylformamide were dissolved 401 mg of 4-chloro-2-methyl-6-(2,3-difluorophenyl)pyrimidine and 152 mg of 2-butyne-1-ol, to which 87 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 11 hours. The reaction mixture was then poured into water and extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 395 mg of 2-methyl-4-(2,3-difluorophenyl)-6-(2-butynyloxy)pyrimidine (the present compound (162)).

¹H-NMR: 1.90 (t, 3H), 2.68 (s, 3H), 5.02 (q, 2H), 7.08 (s, 1H), 7.14-7.30 (m, 2H), 7.78-7.87 (m, 1H)

15 Production Example 161



In 5.5 ml of acetonitrile was dissolved 0.4 g of 6-(2-butynyloxy)-pyrimidin-4-yl 2-fluorophenyl ketone, to which a solution of 0.44 of 2,2-difluoro-1,3-dimethylimidazolidine in 2 ml of acetonitrile was added, followed by heating under reflux for 20 hours. The reaction mixture was then poured into water and extracted three times with ethyl acetate. The organic layers were combined and washed with a saturated aqueous sodium chloride solution, and the combined organic layer was dried over anhydrous magnesium sulfate and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 0.23 g of 4-(2-butynyloxy)-6-(α,α-difluoro-2-fluorobenzyl)pyrimidine (the present compound (163)).

¹H-NMR: 1.88 (t, 3H), 5.03 (q, 2H), 7.06 (dd, 1H), 7.10-7.30 (m, 2H, involving a singlet at 7.26), 7.46 (qd, 1H), 7.77 (td, 1H), 8.78 (s, 1H)

Production Example 162

In 1.5 ml of tetrahydrofuran was suspended 0.04 g of sodium hydride
5 (60% in oil), to which 0.5 ml of a tetrahydrofuran solution containing 0.06 g
of 2-butyne-1-ol was slowly added dropwise under stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, and 0.5
ml of a tetrahydrofuran solution containing 0.2 g of 4-chloro-6-(N-ethyl-N-(2,3-difluorophenyl)amino)pyrimidine was slowly added dropwise, followed
10 by stirring at room temperature for 5 hours. The reaction mixture was then
poured into a saturated aqueous ammonium chloride solution and extracted
three times with chloroform. The chloroform layers were combined, washed
with water, dried over anhydrous magnesium sulfate, and then concentrated.
The residue was subjected to silica gel column chromatography to give 0.16 g
15 of 6-(2-butyneoxy)-4-(N-ethyl-N-(2,3-difluorophenyl)amino)pyrimidine (the
present compound (164)).

¹H-NMR: 1.21 (t, 3H), 1.85 (t, 3H), 3.93 (q, 2H), 4.89 (q, 2H), 5.59 (s, 1H), 7.03-7.21 (m, 3H), 8.39 (s, 1H)

Production Example 163

20 In 3 ml of tetrahydrofuran was suspended 0.10 g of sodium hydride
(60% in oil), to which 0.5 ml of a tetrahydrofuran solution containing 0.15 g
of 2-butyne-1-ol was slowly added dropwise under stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, and 0.5
ml of a tetrahydrofuran solution containing 0.43 g of 4-chloro-6-(N-ethyl-N-
25 (3-fluorophenyl)amino)pyrimidine was slowly added dropwise, followed by
stirring at room temperature for 6 hours. The reaction mixture was then
poured into a saturated aqueous ammonium chloride solution and extracted
three times with ethyl acetate. The organic layers were combined, washed

with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.2 g of 6-(2-butynyloxy)-4-(N-ethyl-N-(3-fluorophenyl)amino)pyrimidine (the present compound (165)).

5 ¹H-NMR: 1.21 (t, 3H), 1.84 (t, 3H), 3.97 (q, 2H), 4.87 (q, 2H), 5.63 (s, 1H), 6.93-7.06 (m, 3H), 7.31-7.42 (m, 1H), 8.39 (s, 1H)

Production Example 164

In 1.2 ml of tetrahydrofuran was suspended 0.03 g of sodium hydride (60% in oil), to which 0.4 ml of a tetrahydrofuran solution containing 0.05 g of 2-butyne-1-ol was slowly added dropwise under stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, and 0.4 ml of a tetrahydrofuran solution containing 0.16 g of 4-chloro-6-(2-chlorocyclohexyloxy)pyrimidine was slowly added dropwise, followed by stirring at 0°C for 3 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with ethyl acetate. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.15 g of 6-(2-butyne-1-ol)-4-(2-chlorocyclohexyloxy)pyrimidine (the present compound (166)).

20 ¹H-NMR: 1.32-1.64 (m, 3H), 1.75-1.82 (m, 3H), 1.87 (t, 3H), 2.18-2.29 (m, 2H), 3.98-4.04 (m, 1H), 4.95 (q, 2H), 5.08-5.22 (m, 1H), 6.12 (s, 1H), 8.43 (s, 1H)

Production Example 165

In 5 ml of N,N-dimethylformamide were dissolved 135 mg of 4-chloro-6-(2,3-difluorophenyl)pyrimidine and 60 mg of 3-pentyn-2-ol, to which 29 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 5 hours. The reaction mixture was then poured into water and extracted with ethyl acetate. The organic layer was washed with a

saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 130 mg of 4-(2,3-difluorophenyl)-6-(1-methyl-2-butyloxy)pyrimidine (the present compound (167)).

5 ¹H-NMR: 1.64 (d, 3H), 1.86 (d, 3H), 5.76-5.90 (m, 1H), 7.14-7.33 (m, 3H, involving a singlet at 7.24), 7.81-7.91 (m, 1H), 8.90 (s, 1H)

Production Example 166

In 5 ml of tetrahydrofuran was suspended 0.15 g of sodium hydride (60% in oil), to which 1 ml of a tetrahydrofuran solution containing 0.20 g of
10 2-butylo-1-ol was slowly added dropwise under stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, and 1 ml of a tetrahydrofuran solution containing 0.55 g of 4-chloro-6-(2-methylcyclohexyloxy (cis : trans = 3 : 7))pyrimidine was slowly added dropwise, followed by stirring at room temperature for 3 hours. The reaction mixture was then
15 poured into a saturated aqueous ammonium chloride solution and extracted three times with ethyl acetate. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.36 g of 4-(2-butyloxy)-6-(2-methylcyclohexyloxy)pyrimidine (the present compound (168)) as a mixture of its cis and trans forms.
20

Cis Form:

¹H-NMR: 0.93 (d, 3H), 1.51-1.98 (m, 12H, involving a triplet at 1.87), 4.95 (q, 2H), 5.15-5.19 (m, 1H), 6.09 (s, 1H), 8.42 (s, 1H)

Trans Form:

25 ¹H-NMR: 0.93 (d, 3H), 1.10-1.37 (m, 4H), 1.63-1.83 (m, 4H), 1.87 (t, 3H), 2.08-2.14 (m, 1H), 4.69 (td, 1H), 4.94 (q, 2H), 6.06 (s, 1H), 8.42 (s, 1H)

Production Example 167

In 3.6 ml of tetrahydrofuran was suspended 0.09 g of sodium hydride

(60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.14 g of 2-butyn-1-ol was slowly added dropwise under stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, and 0.6 ml of a tetrahydrofuran solution containing 0.38 g of 4-chloro-6-(trans-2-methylcyclopentyloxy)pyrimidine was slowly added dropwise, followed by stirring at room temperature for 3 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with ethyl acetate. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.36 g of 4-(2-butynyloxy)-6-(trans-2-methylcyclopentyloxy)pyrimidine (the present compound (169)).

¹H-NMR: 1.04 (d, 3H), 1.20-1.27 (m, 1H), 1.66-2.17 (m, 9H, involving a triplet at 1.87), 4.90-4.95 (m, 3H), 6.06 (s, 1H), 8.43 (s, 1H)

15 Production Example 168

In 4 ml of tetrahydrofuran was suspended 0.10 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.15 g of 2-butyn-1-ol was slowly added dropwise under stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, and 0.6 ml of a tetrahydrofuran solution containing 0.43 g of 4-chloro-6-(cycloheptyloxy)pyrimidine was slowly added dropwise, followed by stirring at room temperature for 3 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with *t*-butyl methyl ether. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.52 g of 6-(2-butynyloxy)-4-(cycloheptyloxy)pyrimidine (the present compound (170)).

¹H-NMR: 1.48-1.80 (m, 10H), 1.88 (t, 3H), 1.98-2.06 (m, 2H), 4.93 (q, 2H), 5.10-5.22 (m, 1H), 6.04 (s, 1H), 8.42 (s, 1H)

Production Example 169

To a solution of 0.3 g of 4-(2-propynyloxy)-6-phenoxy pyrimidine dissolved in 3 ml of ethanol were added 1.87 ml of 10% sodium hydroxide and 0.51 g of iodine at 0°C. The mixture was stirred at room temperature for 5 hours, and the alcohol was distilled out under reduced pressure. A saturated sodium thiosulfate solution was added to the residue, which was extracted three times with ethyl acetate. The organic layers were combined, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.12 g of 4-(3-iodo-2-propynyloxy)-6-phenoxy pyrimidine (the present compound (171)).

¹H-NMR: 5.15 (s, 2H), 6.17 (s, 1H), 7.13 (d, 2H), 7.27 (t, 1H), 7.43 (t, 2H), 8.46 (s, 1H)

15 Production Example 170

In 1 ml of tetrahydrofuran was suspended 0.03 g of sodium hydride (60% in oil), to which 0.2 ml of a tetrahydrofuran solution containing 0.04 g of 2-butyne-1-ol was slowly added dropwise under stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, and 0.2 ml of a tetrahydrofuran solution containing 0.12 g of 4-chloro-6-(cis-2-methylcyclohexyloxy)pyrimidine was slowly added dropwise, followed by stirring at room temperature for 5 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with t-butyl methyl ether. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.10 g of 4-(2-butyloxy)-6-(cis-2-methylcyclohexyloxy)pyrimidine (the present compound (172)).

¹H-NMR: 0.93 (d, 3H), 1.51-1.98 (m, 12H, involving a triplet at 1.87), 4.95 (q, 2H), 5.15-5.19 (m, 1H), 6.09 (s, 1H), 8.42 (s, 1H)

Production Example 171

In 5 ml of N,N-dimethylformamide were dissolved 246 mg of 4-chloro-6-(3-fluorophenyl)pyrimidine and 119 mg of 3-pentyn-2-ol, to which 57 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 8 hours. The reaction mixture was then poured into water and extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 200 mg of 4-(3-fluorophenyl)-6-(1-methyl-2-butynyloxy)pyrimidine (the present compound (173)).

¹H-NMR: 1.63 (d, 3H), 1.84 (d, 3H), 5.81-5.91 (m, 1H), 7.09 (s, 1H), 7.14-7.21 (m, 1H), 7.40-7.51 (m, 1H), 7.73-7.83 (m, 2H), 8.85 (s, 1H)

15 Production Example 172

In 10 ml of N,N-dimethylformamide were dissolved 344 mg of 4-chloro-6-(4-fluorophenyl)pyrimidine and 166 mg of 3-pentyn-2-ol, to which 166 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 10 hours. The reaction mixture was then poured into water and extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 205 mg of 4-(4-fluorophenyl)-6-(1-methyl-2-butynyloxy)pyrimidine (the present compound (174)).

25 ¹H-NMR: 1.62 (d, 3H), 1.84 (d, 3H), 5.80-5.91 (m, 1H), 7.07 (s, 1H), 7.12-7.22 (m, 2H), 8.00-8.09 (m, 2H), 8.83 (s, 1H)

Production Example 173

In 1.6 ml of tetrahydrofuran was suspended 0.04 g of sodium hydride

(60% in oil), to which 0.4 ml of a tetrahydrofuran solution containing 0.06 g of 2-butyne-1-ol was slowly added dropwise under stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, and 0.4 ml of a tetrahydrofuran solution containing 0.18 g of 4-chloro-6-(trans-2-methylcyclohexyloxy)pyrimidine was slowly added dropwise, followed by stirring at room temperature for 6 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with t-butyl methyl ether. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.14 g of 6-(2-butyneoxy)-4-(trans-2-methylcyclohexyloxy)pyrimidine (the present compound (175)).

¹H-NMR: 0.93 (d, 3H), 1.10-1.37 (m, 4H), 1.63-1.83 (m, 4H), 1.87 (t, 3H), 2.08-2.14 (m, 1H), 4.69 (td, 1H), 4.94 (q, 2H), 6.06 (s, 1H), 8.42 (s, 1H)

15 Production Example 174

Under an atmosphere of a nitrogen gas, 192 mg of sodium hydride was added to 10 ml of tetrahydrofuran, followed by ice cooling, to which an tetrahydrofuran (4 ml) solution of 280 mg of 2-butyne-1-ol was added dropwise, and the mixture was stirred at room temperature for 1 hour. Under ice cooling, an tetrahydrofuran (4 ml) solution of 778 mg of 4-(2-butyneoxy)-6-(3,3-dimethyl-1-butyne)pyrimidine was added dropwise. The mixture was stirred at room temperature for 4 hours and then poured into water. The mixture was extracted with t-butyl methyl ether and then washed three times with water. And the organic layers were dried over sodium sulfate and concentrated. The residue was subjected to silica gel thin layer chromatography to give 560 mg of 4-(2-butyneoxy)-6-(3,3-dimethyl-1-ethynyl)pyrimidine (the present compound (176)).

¹H-NMR: 1.34 (s, 9H), 1.87 (t, 3H), 4.97 (q, 2H), 6.78 (s, 1H), 8.70 (s, 1H)

Production Example 175

To 0.5 ml of ethanol were added 0.3 g of 4,5-dichloro-6-(2-butynyl-oxy)pyrimidine and 0.48 g of N-ethylpropylamine, followed by heating under reflux for 8 hours. The reaction mixture was then left for cooling to room temperature and concentrated under reduced pressure. The residue was subjected to silica gel column chromatography to give 0.15 g of 5-choloro-4-(N-ethylpropylamino)-6-(2-butynyloxy)pyrimidine(the present compound (177)).

¹H-NMR: 0.91 (t, 3H), 1.22 (t, 3H), 1.67 (dt, 2H), 1.87 (t, 3H), 3.50 (t, 2H), 3.63 (q, 2H), 4.98 (q, 2H), 8.16 (s, 1H)

Production Example 176

In 5.4 ml of chloroform was dissolved 0.62 g of 4-chloro-6-(2-hydroxycyclohexyloxy)pyrimidine (mixture of cis-form and trans-form), to which 1.09 g of (dimethylamino)sulfur trifluoride was slowly added dropwise under stirring at room temperature, followed by further stirring at room temperature for 1 hour. The reaction mixture was then poured into water and extracted three times with t-butyl ethyl ether. The organic layers were combined, washed with a saturated aqueous sodium hydrogen carbonate, brine, and dried over anhydrous magnesium sulfate, and then concentrated. The residue was used for the next steps without purification.

In 2 ml of tetrahydrofuran was suspended 0.05 g of sodium hydride (60% in oil), to which 0.5ml of a tetrahydrofuran solution containing 0.08 g of 2-butyln-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 1 ml of a tetrahydrofuran solution containing the mixture of the crude product was slowly added dropwise, followed by further stirring at room temperature for 1 hour. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with t-butyl methyl

ether. The organic layers were combined, washed with brine, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.14 g of 4-(2-butynyloxy)-6-(2-fluorocyclohexyloxy)pyrimidine (the present compound
5 (178)).

¹H-NMR: 1.32-1.46 (m, 2H), 1.59-1.94 (m, 7H, involving a triplet at 1.87), 2.18-2.21 (m, 2H), 4.45-4.70 (m, 1H), 4.95 (q, 2H), 5.17-5.27 (m, 1H), 6.12 (s, 1H), 8.43 (s, 1H) with peaks due to the minor isomer at 6.18 (s), 8.48 (s)

10 Production Example 177

To 2 ml of N,N-dimethylformamide were added 0.1 g of 4-chloro-6-(2-butynyloxy)-5-fluoropyrimidine, 0.1 g of potassium carbonate, and 0.06 g of phenol, followed by stirring at 80°C for 2 hours. The reaction mixture was left for cooling to room temperature and poured into a saturated aqueous
15 ammonium chloride solution, which was extracted three times with t-butyl methyl ether. The organic layers were combined, washed with brine, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.12 g of 4-(2-butynyloxy)-5-fluoro-6-phenoxy pyrimidine (the present compound (179)).

20 ¹H-NMR: 1.88 (t, 3H), 5.07 (q, 2H), 7.17 (d, 2H), 7.27 (t, 1H), 7.43 (t, 2H), 8.15 (s, 1H)

 Production Example 178

To 2 ml of N,N-dimethylformamide were added 0.1 g of 4-chloro-6-(2-butynyloxy)-5-fluoropyrimidine, 0.1 g of potassium carbonate, and 0.08 g of
25 2,3-difluorophenol, followed by stirring at 80°C for 2 hours. The reaction mixture was left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with t-butyl methyl ether. The organic layers were combined, washed with brine,

dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.12 g of 4-(2-butynyloxy)-6-(2,3-difluorophenoxy)-5-fluoropyrimidine (the present compound (180)).

5 ¹H-NMR: 1.88 (t, 3H), 5.08 (q, 2H), 7.02-7.15 (m, 3H), 8.12 (s, 1H)

Production Example 179

In 4 ml of tetrahydrofuran was suspended 0.16 g of sodium hydride (60% in oil), to which 0.5ml of a tetrahydrofuran solution containing 0.23 g of 2-butyn-1-ol was slowly added dropwise with stirring at room temperature.

10 The mixture was stirred at room temperature for 20 minutes, to which 1.5 ml of a tetrahydrofuran solution containing 0.72 g of 4-chloro-6-(2,3-dimethylcyclohexyloxy)pyrimidine (mixture of isomers) was slowly added dropwise at 0°C, followed by further stirring at room temperature for 3 hours. The reaction mixture was then poured into a saturated aqueous ammonium

15 chloride solution and extracted three times with t-butyl methyl ether. The organic layers were combined, washed with brine, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.67 g of 4-(2-butynyloxy)-6-(2,3-dimethylcyclohexyloxy)pyrimidine (the present compound (181)).

20 ¹H-NMR: 0.83-2.26 (m, 17H, involving a triplet at 1.87), 4.67-4.77 (m, 1H), 4.94 (q, 2H), 6.07 (s, 1H), 8.42 (s, 1H) with peaks due to the minor isomers at 4.99-5.11 (m)

Production Example 180

In 2 ml of tetrahydrofuran was suspended 0.08 g of sodium hydride

25 (60% in oil), to which 0.5 ml of a tetrahydrofuran solution containing 0.14 g of 2-butyn-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 1.5 ml of a tetrahydrofuran solution containing 0.31 g of 4-chloro-6-(3-methyl-

cyclohexyloxy)pyrimidine (mixture of cis-form and trans-form) was slowly added dropwise at 0°C, followed by further stirring at room temperature for 3 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with t-butyl methyl
5 ether. The organic layers were combined, washed with brine, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.26 g of 4-(2-butynyloxy)-6-(3-methylcyclohexyloxy)pyrimidine (the present compound (182)).

¹H-NMR: 0.84-2.12 (m, 15H), 4.93-5.02 (m, 3H), 6.05 (s, 1H), 8.42 (s,
10 1H) with peaks due to the minor isomer at 5.31-5.34 (m), 6.07 (s)

Production Example 181

To 2 ml of N,N-dimethylformamide were added 0.1 g of 4-chloro-6-(2-butynyloxy)-5-fluoropyrimidine, 0.1 g of potassium carbonate, and 0.07 g of 2-fluorophenol, followed by stirring at 80°C for 3 hours. The reaction
15 mixture was left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with t-butyl methyl ether. The organic layers were combined, washed with brine, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.11 g of
20 4-(2-butynyloxy)-6-(2-fluorophenoxy)-5-fluoropyrimidine (the present compound (183)).

¹H-NMR: 1.88 (t, 3H), 5.07 (q, 2H), 7.17-7.28 (m, 4H), 8.13 (s, 1H)

Production Example 182

To 2 ml of N,N-dimethylformamide were added 0.1 g of 4-chloro-6-(2-butynyloxy)-5-fluoropyrimidine, 0.1 g of potassium carbonate, and 0.08 g of
25 2-chlorophenol, followed by stirring at 60°C for 1 hour. The reaction mixture was left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with

t-butyl methyl ether. The organic layers were combined, washed with brine, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.12 g of 4-(2-butynyloxy)-6-(2-chlorophenoxy)-5-fluoropyrimidine (the present compound (184)).

¹H-NMR: 1.88 (t, 3H), 5.07 (q, 2H), 7.22-7.37 (m, 3H), 7.50 (d, 1H), 8.12 (s, 1H)

Production Example 183

To 0.5 ml of ethanol were added 0.1 g of 4-(2-butynyloxy)-6-chloro-5-fluoropyrimidine and 0.11 g of N-ethylpropylamine, followed by heating under reflux for 10 hours. The reaction mixture was then left for cooling to room temperature and concentrated under reduced pressure. The residue was added water and extracted three times with t-butyl methyl ether. The organic layers were combined, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.15 g of 6-(2-butynyloxy)-4-(N-ethylpropylamino)-5-fluoropyrimidine (the present compound (185)).

¹H-NMR: 0.92 (t, 3H), 1.21 (t, 3H), 1.65 (dt, 2H), 1.87 (t, 3H), 3.45 (t, 2H), 3.57 (q, 2H), 4.97 (q, 2H), 8.01(s, 1H)

20 Production Example 184

In 2 ml of tetrahydrofuran was suspended 0.11 g of sodium hydride (60% in oil), to which 0.5 ml of a tetrahydrofuran solution containing 0.14 g of 2-butyne-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 1 ml of a tetrahydrofuran solution containing 0.5 g of 4,5-dichloro-6-(N-ethyl-N-phenylamino)pyrimidine was slowly added dropwise at room temperature, followed by further stirring at room temperature for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride

solution and extracted three times with t-butyl methyl ether. The organic layers were combined, washed with brine, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.38 g of 6-(2-butynyloxy)-5-chloro-4-(N-ethyl-N-phenylamino)pyrimidine (the present compound (186)).

¹H-NMR: 1.22 (t, 3H), 1.86 (t, 3H), 4.03 (q, 2H), 4.99 (q, 2H), 7.04-7.35 (m, 5H), 8.35 (s, 1H)

Production Example 185

In 1.5 ml of tetrahydrofuran was suspended 0.08 g of sodium hydride (60% in oil) to which 0.5 ml of a tetrahydrofuran solution containing 0.13 g of 2-butyn-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 1.5 ml of a tetrahydrofuran solution containing 0.37 g of 4-chloro-6-(cis-4-methylcyclohexyloxy)pyrimidine was slowly added dropwise at room temperature, followed by further stirring at room temperature for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with t-butyl methyl ether. The organic layers were combined, washed with brine, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.36 g of 4-(2-butynyloxy)-6-(cis-4-methylcyclohexyloxy)pyrimidine (the present compound (187)).

¹H-NMR: 0.91 (d, 3H), 1.27-1.66 (m, 7H), 1.86-2.01 (m, 5H involving a triplet at 1.88), 4.94 (q, 2H), 5.18-5.24 (m, 1H), 6.08 (s, 1H), 8.42 (s, 1H)

Production Example 186

In 1.5 ml of tetrahydrofuran was suspended 0.09 g of sodium hydride (60% in oil), to which 0.5ml of a tetrahydrofuran solution containing 0.13 g of 2-butyn-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 1.5 ml

of a tetrahydrofuran solution containing 0.37 g of 4-choloro-6-(trans-4-methylcyclohexyloxy)pyrimidine was slowly added dropwise at room temperature, followed by further stirring at room temperature for 2 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with t-butyl methyl ether. The organic layers were combined, washed with brine, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.43 g of 4-(2-butynyloxy)-6-(trans-4-methylcyclohexyloxy)pyrimidine (the present compound (188)).

¹H-NMR: 0.91 (d, 3H), 1.06-1.16 (m, 2H), 1.37-1.51 (m, 3H), 1.75-1.79 (m, 2H), 1.87 (t, 3H), 2.07-2.13 (m, 2H), 4.90-4.97 (m, 3H), 6.04 (s, 1H), 8.41 (s, 1H)

Production Example 187

In 1.5 ml of tetrahydrofuran was suspended 0.09 g of sodium hydride (60% in oil), to which 0.5ml of a tetrahydrofuran solution containing 0.13 g of 2-butyn-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 1.5 ml of a tetrahydrofuran solution containing 0.5 g of 4-choloro-6-(cis-2-trimethylsilanyloxy)cyclohexyloxy)pyrimidine was slowly added dropwise at room temperature, followed by further stirring at room temperature for 3 hours. The reaction mixture was then poured into 10 % hydrochloric acid at room temperature, followed by further stirring for 10 minutes. Then the mixture was extracted three times with t-butyl methyl ether. The organic layers were combined, washed with brine, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.25 g of 4-(2-butynyloxy)-6-(cis-2-hydroxycyclohexyloxy)pyrimidine (the present compound (189)).

¹H-NMR: 1.37-1.44 (m, 2H), 1.65-2.01 (m, 9H, involving a triplet at

1.87), 2.98 (bs, 1H), 3.96-3.99 (m, 1H), 4.95 (q, 2H), 5.18-5.23 (m, 1H), 6.12 (s, 1H), 8.40 (s, 1H)

Production Example 188

To 2 ml of N,N-dimethylformamide were added 0.1 g of 4-chloro-6-(2-butynyloxy)-5-fluoropyrimidine, 0.1 g of potassium carbonate, and 0.08 g of 2,6-difluorophenol, followed by stirring at 60°C for 4 hours. The reaction mixture was left for cooling to room temperature and poured into a saturated aqueous ammonium chloride solution, which was extracted three times with t-butyl methyl ether. The organic layers were combined, washed with brine, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.1 g of 4-(2-butynyloxy)-6-(2,6-difluorophenoxy)-5-fluoropyrimidine (the present compound (190)).

¹H-NMR: 1.88 (t, 3H), 5.08 (q, 2H), 7.00-7.07 (m, 2H), 7.18-7.27 (m, 1H), 8.12 (s, 1H)

Production Example 189


In 3 ml of tetrahydrofuran was suspended 0.1 g of sodium hydride (60% in oil), to which 0.5 ml of a tetrahydrofuran solution containing 0.15 g of 2-butyne-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 1 ml of a tetrahydrofuran solution containing 0.46 g of 4-chloro-6-(2-fluorobenzyl)-5-fluoropyrimidine was slowly added dropwise at 0°C, followed by further stirring at 0°C for 2 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with t-butyl methyl ether. The organic layers were combined, washed with brine, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.46 g of 4-(2-butynyloxy)-6-(2-fluorobenzyl)-5-

192

fluoropyrimidine (the present compound (191)).

¹H-NMR: 1.86 (t, 3H), 4.16 (s, 2H), 5.04 (q, 2H), 6.99-7.10 (m, 2H),
7.18-7.30 (m, 2H), 8.46 (s, 1H)

The present compounds described in the above production examples are shown with their compound numbers in the following tables.



Compd. No.	R ¹	R ⁴	R ²	R ³
1	2-butynyl	2-butynyloxy	H	H
2	2-propynyl	2-butynyloxy	H	H
3	2-pentynyl	2-pentynyloxy	H	H
4	2-butynyl	4-chloro-2-fluorophenoxy	H	H
5	2-pentynyl	2-propynyloxy	H	H
6	2-pentynyl	2-butynyloxy	H	H
7	4,4-dimethyl-2-pentynyl	2-butynyloxy	H	H
8	4,4-dimethyl-2-pentynyl	4,4-dimethyl-2-pentynyloxy	H	H
9	2-butynyl	phenoxy	H	H
10	2-butynyl	3,4-difluorophenoxy	H	H
11	2-propynyl	1-methyl-2-propynyloxy	H	H
12	2-propynyl	3-butynyloxy	H	H
13	2-propynyl	benzyloxy	H	H
14	2-propynyl	4-chlorophenoxy	H	H
15	2-propynyl	3-chlorophenoxy	H	H
16	2-butynyl	2-chloro-4-fluorophenoxy	H	H
17	2-propynyl	3-trifluoromethylphenoxy	H	H
18	2-propynyl	2-trifluoromethylphenoxy	H	H
19	2-propynyl	2-chlorophenoxy	H	H
20	2-propynyl	4-trifluoromethylphenoxy	H	H
21	2-propynyl	2,6-difluorophenoxy	H	H
22	2-propynyl	2,4-dichlorophenoxy	H	H
23	2-propynyl	3,4-dichlorophenoxy	H	H
24	2-propynyl	3,5-dichlorophenoxy	H	H
25	2-propynyl	2,5-dichlorophenoxy	H	H

Compd. No.	R ¹	R ⁴	R ²	R ³
26	2-propynyl	2,3-dichlorophenoxy	H	H
27	2-propynyl	2-methylphenoxy	H	H
28	2-propynyl	4-methylphenoxy	H	H
29	2-propynyl	3-methylphenoxy	H	H
30	2-propynyl	3-methoxyphenoxy	H	H
31	2-propynyl	4-methoxyphenoxy	H	H
32	2-propynyl	2-methoxyphenoxy	H	H
33	2-butyryl	2,6-difluorophenoxy	H	H
34	2-propynyl	2-fluorophenoxy	H	H
35	2-propynyl	4-fluorophenoxy	H	H
36	2-propynyl	3-fluorophenoxy	H	H
37	2-propynyl	phenyl	H	H
38	2-butyryl	phenyl	H	H
39	2-butyryl	2,3-difluorophenoxy	H	H
40	2-butyryl	3-cyanophenoxy	H	H
41	2-butyryl	4-cyanophenoxy	H	H
42	2-butyryl	2-cyanophenoxy	H	H
43	2-butyryl	2,5-difluorophenoxy	H	H
44	2-butyryl	2,4-difluorophenoxy	H	H
45	2-butyryl	2,4,6-trifluorophenoxy	H	H
46	2-butyryl	2,3,6-trifluorophenoxy	H	H
47	2-butyryl	2-chloro-4,6-difluorophenoxy	H	H
48	2-butyryl	4-fluoro-3-trifluoromethyl- phenoxy	H	H
49	2-butyryl	3-trifluoromethoxyphenoxy	H	H
50	2-butyryl	4-trifluoromethoxyphenoxy	H	H

Compd. No.	R ¹	R ⁴	R ²	R ³
51	2-propynyl	phenoxy	H	H
52	2-propynyl	2-propynyloxy	H	H
53	2-butynyl	phenylamino	H	H
54	2-propynyl	2-fluorophenyl	H	H
55	2-butynyl	2-fluorophenyl	H	H
56	2-propynyl	3-fluorophenyl	H	H
57	2-butynyl	3-fluorophenyl	H	H
58	2-propynyl	4-fluorophenyl	H	H
59	2-butynyl	4-fluorophenyl	H	H
60	2-pentynyl	2,3-difluorophenoxy	H	H
61	2-butynyl	3-fluorophenoxy	H	H
62	2-butynyl	4-fluorophenoxy	H	H
63	2-butynyl	2-fluorophenoxy	H	H
64	2-butynyl	2,3-(methylenedioxy)phenoxy	H	H
65	2-butynyl	2-fluoro-4-nitrophenoxy	H	H
66	2-butynyl	N-(2,3-difluorophenyl)-N-methylamino	H	H
67	2-butynyl	2,3-dimethylphenoxy	H	H
68	2-butynyl	2,6-difluorobenzyloxy	H	H
69	2-butynyl	3-phenylphenoxy	H	H
70	2-butynyl	3-phenoxyphenoxy	H	H
71	2-butynyl	3-acetylphenoxy	H	H
72	2-butynyl	α -cyano-2,3-difluorobenzyl	H	H
73	2-butynyl	2,3-difluorophenylamino	H	H
74	2-butynyl	α -cyanobenzyl	H	H
75	2-butynyl	benzoyl	H	H

Compd. No.	R ¹	R ⁴	R ²	R ³
76	2-butynyl	2,3-difluorobenzyloxy	H	H
77	2-propynyl	phenyl	CH ₃	H
78	2-butynyl	phenyl	CH ₃	H
79	2-butynyl	N-(2,3-difluorophenyl)-N-methoxymethylamino	H	H
80	2-butynyl	2,6-difluorobenzyl	H	H
81	2-butynyl	N-phenyl-N-ethylamino	H	H
82	2-propynyl	2,3-difluorophenyl	H	H
83	2-butynyl	2,3-difluorophenyl	H	H
84	2-butynyl	α -cyano-2-chloro-6-fluorobenzyl	H	H
85	2-butynyl	N-(2,3-difluorophenyl)-N-cyanomethylamino	H	H
86	2-pentynyl	2-fluorophenyl	H	H
87	2-butynyl	2,6-difluorophenyl	H	H
88	2-butynyl	2-fluorobenzyl	H	H
89	2-butynyl	2-chlorobenzyl	H	H
90	2-butynyl	2,3,5,6-tetrafluorophenoxy	H	H
91	2-butynyl	benzyl	H	H
92	2-butynyl	2-methylbenzyl	H	H
93	2-butynyl	N-methyl-N-phenylamino	H	H
94	2-butynyl	1-phenylethyl	H	H
95	2-butynyl	2-trifluorobenzyl	H	H
96	2-butynyl	2,3-difluorobenzyl	H	H
97	2-butynyl	phenylthio	H	H
98	2-butynyl	N-propyl-N-phenylamino	H	H
99	2-butynyl	2,4-difluorobenzyl	H	H
100	2-butynyl	3-fluorobenzyl	H	H

Compd. No.	R ¹	R ⁴	R ²	R ³
101	2-butynyl	2-chloro-6-fluorobenzyl	H	H
102	2-butynyl	3-chloro-2-fluorobenzyl	H	H
103	2-butynyl	2-bromobenzyl	H	H
104	2-butynyl	N-ethyl-N-methylamino	H	H
105	2-butynyl	N-ethyl-N-isopropylamino	H	H
106	2-butynyl	isopropylamino	H	H
107	2-butynyl	N-(2-butynyl)-N-isopropylamino	H	H
108	2-butynyl	ethylamino	H	H
109	2-butynyl	N-ethyl-N-(2-butynyl)amino	H	H
110	2-butynyl	N-ethyl-N-(2-propenyl)amino	H	H
111	2-butynyl	2,2,3,3,3-pentafluoropropyl amino	H	H
112	2-butynyl	N-(2,2,3,3,3-pentafluoropropyl)-N-ethylamino	H	H
113	2-butynyl	di-n-propylamino	H	H
114	2-butynyl	2,2,2-trifluoroethylamino	H	H
115	2-butynyl	N-ethyl-N-(2,2,2-trifluoroethyl)amino	H	H
116	2-butynyl	1-(3-fluorophenyl)ethyl	H	H
117	2-pentynyl	2,3-difluorophenyl	H	H
118	2-heptynyl	2,3-difluorophenyl	H	H
119	4,4-dimethyl-2-pentynyl	2,3-difluorophenyl	H	H
120	2-butynyl	1-(2-fluorophenyl)ethyl	H	H
121	2-butynyl	2-chloro-5-methyl-6-fluorobenzyl	H	H
122	2-butynyl	α -methoxybenzyl	H	H
123	2-butynyl	α -hydroxybenzyl	H	H
124	2-butynyl	α -fluorobenzyl	H	H
125	2-butynyl	α -methoxyiminobenzyl (A)	H	H

Compd. No.	R ¹	R ⁴	R ²	R ³
126	2-butynyl	α -methoxyiminobenzyl (B)	H	H
127	2-butynyl	α -ethoxyiminobenzyl (A)	H	H
128	2-butynyl	α -ethoxyiminobenzyl (B)	H	H
129	2-butynyl	N-benzyl-N-ethylamino	H	H
130	2-butynyl	2-chloro-3,6-difluorobenzyl	H	H
131	2-butynyl	α -isopropoxyiminobenzyl (A)	H	H
132	2-butynyl	α -isopropoxyiminobenzyl (B)	H	H
133	2-butynyl	benzylamino	H	H
134	1-methyl-propynyl	2,3-difluorophenyl	H	H
135	4-fluoro-2-butynyl	2,3-difluorophenyl	H	H
136	3-chloro-2-propynyl	2,3-difluorophenyl	H	H
137	2-butynyl	α -methoxymethoxybenzyl	H	H
138	2-butynyl	α -ethoxybenzyl	H	H
139	2-butynyl	α -acetoxybenzyl	H	H
140	2-butynyl	α -propionyloxybenzyl	H	H
141	2-butynyl	α -isobutyryloxybenzyl	H	H
142	2-butynyl	2-chlorophenylthio	H	H
143	2-butynyl	4-fluorobenzyl	H	H
144	2-butynyl	diethylamino	H	H
145	2-butynyl	α -cyano-2-fluorobenzyl	H	H
146	2-butynyl	2-fluorophenylthio	H	H
147	1-methyl-2-butynyl	2,6-difluorobenzyl	H	H
148	2-butynyl	α -hydroxy- α -methylbenzyl	H	H
149	2-butynyl	α -fluoro-2-fluorobenzyl	H	H
150	1-methyl-2-propynyl	2-fluorophenyl	H	H

Compd. No.	R ¹	R ⁴	R ²	R ³
151	1-methyl-2-butyne	2-fluorophenyl	H	H
152	3-butyne	2-fluorophenyl	H	H
153	3-pentyne	2-fluorophenyl	H	H
154	2-butyne	α -fluoro- α -methylbenzyl	H	H
155	2-butyne	1-phenylvinyl	H	H
156	2-butyne	2-chloro-5-fluorophenoxy	H	H
157	2-butyne	α -hydroxy- α -methyl-2-fluorobenzyl	H	H
158	2-butyne	cyclopentyloxy	H	H
159	2-butyne	cyclohexyloxy	H	H
160	2-butyne	2,6-dichloro-4-fluorophenoxy	H	H
161	1-methyl-2-propyne	2,3-difluorophenyl	CH ₃	H
162	2-butyne	2,3-difluorophenyl	CH ₃	H
163	2-butyne	α,α -difluoro-2-fluorobenzyl	H	H
164	2-butyne	N-ethyl-N-2,3-difluorophenyl-amino	H	H
165	2-butyne	N-ethyl-N-3-fluorophenyl-amino	H	H
166	2-butyne	2-chlorocyclohexyloxy	H	H
167	1-methyl-2-butyne	2,3-difluorophenyl	H	H
168	2-butyne	2-methylcyclohexyloxy	H	H
169	2-butyne	2-methylcyclopentyloxy	H	H
170	2-butyne	cycloheptyloxy	H	H
171	3-iodo-2-propyne	phenoxy	H	H
172	2-butyne	cis-2-methylcyclohexyloxy	H	H
173	1-methyl-2-butyne	3-fluorophenyl	H	H
174	1-methyl-2-butyne	4-fluorophenyl	H	H
175	2-butyne	trans-2-methylcyclohexyloxy	H	H


Compd. No.	R ¹	R ⁴	R ²	R ³
176	2-butynyl	3,3-dimethyl-1-butynyl	H	H
177	2-butynyl	N-ethyl-N-n-propyl-amino	H	Cl
178	2-butynyl	2-fluorocyclohexyloxy	H	H
179	2-butynyl	phenoxy	H	F
180	2-butynyl	2,3-difluorophenoxy	H	F
181	2-butynyl	2,3-dimethylcyclohexyloxy	H	H
182	2-butynyl	3-methylcyclohexyloxy	H	H
183	2-butynyl	2-fluorophenoxy	H	F
184	2-butynyl	2-chlorophenoxy	H	F
185	2-butynyl	N-ethyl-N-n-propyl-amino	H	F
186	2-butynyl	N-ethyl-N-phenyl-amino	H	Cl
187	2-butynyl	cis-4-methylcyclohexyloxy	H	H
188	2-butynyl	trans-4-methylcyclohexyloxy	H	H
189	2-butynyl	2-hydroxycyclohexyloxy	H	H
190	2-butynyl	2,6-difluorophenoxy	H	F
191	2-butynyl	2-fluorobenzyl	H	F

The following will describe the reference production examples for the intermediates used in the production of the present compounds.

Reference Production Example 1

In 12 ml of tetrahydrofuran was suspended 0.61 g of sodium hydride
5 (60% in oil), to which 4 ml of a tetrahydrofuran solution containing 0.57 g of
2-propyn-1-ol was slowly added dropwise with stirring at room temperature.
The mixture was stirred at room temperature for 20 minutes and then cooled
to 0°C, to which 4 ml of a tetrahydrofuran solution containing 1.5 g of 4,6-di-
chloropyrimidine was slowly added dropwise, followed by further stirring for
10 2.5 hours. The reaction mixture was then poured into a saturated aqueous
ammonium chloride solution and extracted three times with chloroform.
The chloroform layers were combined, washed with water, dried over anhy-
drous magnesium sulfate, and then concentrated. The residue was subject-
ed to silica gel column chromatography to give 1.61 g of 4-chloro-6-(2-pro-
15 pynyloxy)pyrimidine.

4-Chloro-6-(2-propynyloxy)pyrimidine




¹H-NMR: 2.58 (t, 1H), 5.06 (d, 2H), 6.86 (s, 1H), 8.63 (s, 1H)

Reference Production Example 2

20 In 24 ml of tetrahydrofuran was suspended 1.05 g of sodium hydride
(60% in oil), to which 8 ml of a tetrahydrofuran solution containing 1.42 g of
2-butyln-1-ol was slowly added dropwise with stirring at room temperature.
The mixture was stirred at room temperature for 20 minutes and then cooled
to 0°C, to which 8 ml of a tetrahydrofuran solution containing 3 g of 4,6-di-
25 chloropyrimidine was slowly added dropwise, followed by further stirring at
0°C for 4 hours. The reaction mixture was then poured into a saturated

aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 3.16 g of 4-chloro-6-(2-butynyloxy)pyrimidine, m.p.: 43.5°C.


4-Chloro-6-(2-butynyloxy)pyrimidine



Reference Production Example 3

In 3 ml of tetrahydrofuran was suspended 0.12 g of sodium hydride (60% in oil), to which 0.5 ml of a tetrahydrofuran solution containing 0.22 g of benzyl alcohol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes and then cooled to 0°C, to which 0.5 ml of a tetrahydrofuran solution containing 0.3 g of 4,6-dichloropyrimidine was slowly added dropwise, followed by stirring at 0°C for 1.5 hours and then further stirring at room temperature for 4.5 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.43 g of 4-chloro-6-benzyl-oxyprymidine.

4-Chloro-6-benzyl-oxyprymidine




¹H-NMR: 5.38 (s, 2H), 6.73 (s, 1H), 7.29-7.39 (m, 5H), 8.52 (s, 1H)

Reference Production Example 4

A reaction vessel was charged with 0.17 g of tetrakis(triphenylphosphine) palladium, 0.91 g of phenylboronic acid, and 3.53 g of barium hydroxide, to which 44 ml of 1,2-dimethoxyethane, 8 ml of water, and 1.11 g of 4,6-dichloropyrimidine were added, followed by stirring at 80°C under an atmosphere of a nitrogen gas for 6 hours. The reaction mixture was then left for cooling to room temperature, and water was added to the reaction mixture, which was extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.51 g of 4-chloro-6-phenylpyrimidine, m.p.: 101.3°C.

4-Chloro-6-phenylpyrimidine




Reference Production Example 5

A reaction vessel was charged with 0.258 g of tetrakis(triphenylphosphine) palladium, 1.143 g of 2-fluorophenylboronic acid, and 3.153 g of tripotassium phosphate n-hydrate, to which 36 ml of 1,2-dimethoxyethane, 9 ml of water, and 1.106 g of 4,6-dichloropyrimidine were added, followed by stirring at 80°C under an atmosphere of a nitrogen gas for 9 hours. The reaction mixture was then left for cooling to room temperature, and water was added to the reaction mixture, which was extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resi-

due was subjected to silica gel column chromatography to give 0.519 g of 4-chloro-6-(2-fluorophenyl)pyrimidine.

4-Chloro-6-(2-fluorophenyl)pyrimidine




5 $^1\text{H-NMR}$: 7.17-7.37 (m, 2H), 7.46-7.56 (m, 1H), 7.90 (s, 1H), 8.19 (dt, 1H), 9.07 (s, 1H)

Reference Production Example 6

A reaction vessel was charged with 0.303 g of tetrakis(triphenylphosphine) palladium, 1.344 g of 3-fluorophenylboronic acid, and 3.707 g of tripotassium phosphate n-hydrate, to which 36 ml of 1,2-dimethoxyethane, 9 ml of water, and 1.301 g of 4,6-dichloropyrimidine were added, followed by stirring at 80°C under an atmosphere of a nitrogen gas for 7 hours. The reaction mixture was then left for cooling to room temperature, and water was added to the reaction mixture, which was extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.45 g of 4-chloro-6-(3-fluorophenyl)pyrimidine.

4-Chloro-6-(3-fluorophenyl)pyrimidine




20

$^1\text{H-NMR}$: 7.19-7.29 (m, 1H), 7.45-7.55 (m, 1H), 7.73 (s, 1H), 7.79-7.89 (m, 2H), 9.04 (s, 1H)

Reference Production Example 7

A reaction vessel was charged with 0.255 g of tetrakis(triphenylphosphine) palladium, 1.132 g of 4-fluorophenylboronic acid, and 3.122 g of tripotassium phosphate n-hydrate, to which 36 ml of 1,2-dimethoxyethane, 9 ml of water, and 1.095 g of 4,6-dichloropyrimidine were added, followed by stirring at 80°C under an atmosphere of a nitrogen gas for 8 hours. The reaction mixture was then left for cooling to room temperature, and water was added to the reaction mixture, which was extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.399 g of 4-chloro-6-(4-fluorophenyl)pyrimidine.

4-Chloro-6-(4-fluorophenyl)pyrimidine




¹H-NMR: 7.20 (t, 2H), 7.71 (s, 1H), 8.07-8.12 (m, 2H), 9.01 (s, 1H)

Reference Production Example 8

To 14 ml of ethanol were added 2.2 g of 4,6-dichloropyrimidine and 2.3 g of 2,3-difluoroaniline, followed by heating under reflux for 6 hours. The reaction mixture was then left for cooling to room temperature and concentrated under reduced pressure. The residue was recrystallized from ethyl acetate to give 3.5 g of 4-chloro-6-(2,3-difluoroanilino)pyrimidine.

4-Chloro-6-(2,3-difluoroanilino)pyrimidine




¹H-NMR (DMSO-d₆): 6.89 (s, 1H), 7.20-7.24 (m, 2H), 7.60-7.81 (m, 1H), 8.47 (s, 1H), 9.95 (bs, 1H)

Reference Production Example 9

5 In 2 ml of tetrahydrofuran was suspended 0.07 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.3 g of 4-chloro-6-(2,3-difluorophenylamino)pyrimidine was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 0.6 ml of a tetrahydrofuran solution containing
10 0.2 g of iodomethane was slowly added dropwise, followed by further stirring for 8 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subject-
15 ed to silica gel column chromatography to give 0.16 g of 4-chloro-6-(N-methyl-N-(2,3-difluorophenyl)amino)pyrimidine.

4-Chloro-6-(N-methyl-N-(2,3-difluorophenyl)amino)pyrimidine




¹H-NMR: 3.46 (s, 3H), 6.28 (s, 1H), 7.05-7.26 (m, 3H), 8.49 (s, 1H)

20 Reference Production Example 10

To 2 ml of chloroform were added 0.3 g of 4-chloro-6-(2,3-difluorophenylamino)pyrimidine and 0.65 ml of diisopropylethylamine, to which 0.6 ml of a

chloroform solution containing 0.15 g of chloromethyl methyl ether was slowly added dropwise. After stirring for 8 hours, the reaction mixture was poured into a saturated aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed
5 with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.12 g of 4-(N-methoxymethyl-N-2,3-difluorophenylamino)-6-chloropyrimidine.

4-(N-Methoxymethyl-N-2,3-difluorophenylamino)-6-chloropyrimidine




10 ¹H-NMR: 3.44 (s, 3H), 5.31 (s, 2H), 6.36 (s, 1H), 7.14-7.29 (m, 3H), 8.52 (s, 1H)

Reference Production Example 11

In 2 ml of tetrahydrofuran was suspended 0.07 g of sodium hydride (60% in oil), to which 0.6 ml of a tetrahydrofuran solution containing 0.3 g of
15 4-chloro-6-(2,3-difluoroanilino)pyrimidine was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 0.6 ml of a tetrahydrofuran solution containing 0.18 g of bromoacetonitrile was slowly added dropwise, followed by further stirring for 8 hours. The reaction mixture was then poured into a saturated
20 aqueous ammonium chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.24 g of 4-chloro-6-(N-cyanomethyl-N-(2,3-difluorophenyl)amino)pyrimidine.

4-Chloro-6-(N-cyanomethyl-N-(2,3-difluorophenyl)amino)pyrimidine




Reference Production Example 12

¹H-NMR: 4.85 (bs, 2H), 6.29 (s, 1H), 7.21-7.39 (m, 3H), 8.64 (s, 1H)

5 To 10 ml of an ethanol solution of 1.5 g of 4,6-dichloropyrimidine was slowly added 0.78 g of sodium thiomethoxide at 0°C, followed by stirring at 0°C for 7 hours. A saturated aqueous ammonium chloride solution was added to the reaction mixture and concentrated under reduced pressure. The residue was extracted three times with chloroform. The chloroform
10 layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 1.1 g of 4-chloro-6-methylthiopyrimidine.

4-Chloro-6-methylthiopyrimidine




15 ¹H-NMR: 2.58 (s, 3H), 7.21 (s, 1H), 8.72 (s, 1H)

Reference Production Example 13

In 10 ml of tetrahydrofuran was suspended 0.41 g of sodium hydride (60% in oil), to which 2 ml of a tetrahydrofuran solution containing 0.58 g of 2-butyne-1-ol was slowly added dropwise with stirring at room temperature.
20 The mixture was stirred at room temperature for 20 minutes, to which 2 ml of a tetrahydrofuran solution containing 1.1 g of 4-chloro-6-methylthiopyrimidine was slowly added dropwise, followed by further stirring for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium

chloride solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 1.3 g of 4-(2-butynyloxy)-6-methylthio-
5 pyrimidine.

4-(2-butynyloxy)-6-methylthiopyrimidine




¹H-NMR: 1.87 (t, 3H), 2.52 (s, 3H), 4.59 (q, 2H), 6.60 (s, 1H), 8.57 (s,
1H)

10 Reference Production Example 14

To 14 ml of chloroform were added 4-(2-butynyloxy)-6-methylthiopyrimidine and 3.5 g of m-chloroperbenzoic acid (>65%), followed by stirring at 0°C for 10 hours. The reaction mixture was then poured into a saturated aqueous sodium thiosulfate solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was
15 subjected to silica gel column chromatography to give 1.4 g of 4-(2-butynyloxy)-6-methanesulfonylpyrimidine.

4-(2-Butynyloxy)-6-methanesulfonylpyrimidine



20


¹H-NMR: 1.88 (t, 3H), 3.23 (s, 3H), 5.07 (q, 2H), 7.46 (s, 1H), 8.92 (s,
1H)

Reference Production Example 15

A reaction vessel was charged with 9.84 g of 2-methyl-4,6-dihydroxy-

pyrimidine, 29.46 g of phosphorus oxychloride, and 20.18 g of diisopropyl-ethylamine under ice cooling, followed by stirring for 30 minutes and further stirring at 80°C for 3 hours. The reaction mixture was then left for cooling to room temperature and poured into a saturated aqueous sodium bicarbonate, which was extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium bicarbonate and a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 9.5 g of 2-methyl-4,6-dichloropyrimidine.

10 2-Methyl-4,6-dichloropyrimidine




¹H-NMR: 2.71 (s, 3H), 7.26 (s, 1H)

Reference Production Example 16

A reaction vessel was charged with 2.19 g of tetrakis(triphenylphosphine) palladium, 847 mg of phenylboronic acid, and 2.99 g of barium hydroxide, to which 40 ml of 1,2-dimethoxyethane, 7 ml of water, and 1.01 g of 2-methyl-4,6-dichloropyrimidine were added, followed by stirring at 80°C under an atmosphere of a nitrogen gas for 6 hours. The reaction mixture was then left for cooling to room temperature, and water was added to the reaction mixture, which was extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.65 g of 2-methyl-4-chloro-6-phenylpyrimidine.

25 2-Methyl-4-chloro-6-phenylpyrimidine




¹H-NMR: 1.58 (s, 3H), 7.47-7.58 (m, 4H, involving a singlet at 7.56), 8.02-8.11 (m, 2H)

Reference Production Example 17

5 In 30 ml of tetrahydrofuran was suspended 5.24 g of potassium t-butoxide, to which 2.63 g of phenylacetonitrile and 3.0 g of 4-chloro-6-methylthiopyrimidine, followed by stirring at 0°C for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with ethyl acetate. The organic layers were combined and washed with a saturated aqueous sodium chloride solution, and
10 the combined organic layer was dried over anhydrous magnesium sulfate and then concentrated. The residue was subjected to silica gel column chromatography to give 3.67 g of 4-(α-cyanobenzyl)-6-methylthiopyrimidine.

4-(α-Cyanobenzyl)-6-methylthiopyrimidine



15


¹H-NMR: 2.55 (s, 3H), 5.13 (s, 1H), 7.25 (s, 1H), 7.35-7.45 (m, 5H), 8.89 (s, 1H)

Reference Production Example 18

To 2 ml of ethanol was suspended 0.3 g of 6-methylthio-4-benzoylpyrimidine, to which 0.07 g of sodium borohydride was added, followed by
20 stirring at 0°C for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with ethyl acetate. The organic layers were combined and washed with a

saturated aqueous sodium chloride solution, and the combined organic layer was dried over anhydrous magnesium sulfate and then concentrated to give 6-methylthio-4-(α -hydroxybenzyl)pyrimidine.

6-Methylthio-4-(α -hydroxybenzyl)pyrimidine



5


$^1\text{H-NMR}$: 2.52 (s, 3H), 4.64 (bs, 1H), 5.60 (s, 1H), 7.16 (s, 1H), 7.28-7.35 (m, 5H), 8.81 (s, 1H)

Reference Production Example 19

In 3 ml of tetrahydrofuran was suspended 0.08 g of sodium hydride (60% in oil), to which 6-methylthio-4-(α -hydroxylbenzyl)pyrimidine dissolved in 0.3 ml of tetrahydrofuran was added. The mixture was stirred at 0°C for 15 minutes, to which 0.28 g of iodomethane was added dropwise at room temperature, followed by stirring at the same temperature for 30 minutes. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with ethyl acetate. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.17 g of 6-methylthio-4-(α -methoxybenzyl)pyrimidine.

20

6-Methylthio-4-(α -methoxybenzyl)pyrimidine




$^1\text{H-NMR}$: 2.54 (s, 3H), 3.40 (s, 3H), 5.19 (s, 1H), 7.25-7.41 (m, 5H), 7.45 (s, 1H), 8.82 (s, 1H)

Reference Production Example 20

To 2 ml of chloroform were added 0.17 g of 6-methylthio-4-(α -methoxybenzyl)pyrimidine and 0.43 g of m-chloroperbenzoic acid (>65%), followed by stirring at 0°C for 10 hours. The reaction mixture was then poured into a saturated aqueous sodium thiosulfate solution and extracted three times with chloroform. The chloroform layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.2 g of 4-(α -methoxybenzyl)-6-methanesulfonylpyrimidine.

4-(α -Methoxybenzyl)-6-methanesulfonylpyrimidine



10


¹H-NMR: 3.25 (s, 3H), 3.45 (s, 3H), 5.39 (s, 1H), 7.31-7.44 (m, 5H), 8.31 (s, 1H), 9.21 (s, 1H)

Reference Production Example 21

In 10 ml of tetrahydrofuran was suspended 1.3 g of zinc (powder), to which dibromoethane (2 drops) was added. The mixture was heated under reflux for 5 minutes, to which trimethylsilane chloride was added. The mixture was further heated under reflux for 5 minutes, to which a solution of 1.7 g of benzyl bromide dissolved in 20 ml of tetrahydrofuran was slowly added with heating under reflux, followed by stirring for 20 minutes. (The solution thus obtained is referred to as solution A). In 10 ml of tetrahydrofuran were suspended 1.5 g of 4,6-dichloropyrimidine and 0.1 g of dichlorobistriphenylphosphine palladium, to which the above solution A was added, followed by heating under reflux for 3 hours and further stirring at room temperature for 12 hours. The reaction mixture was then poured into water and extracted three times with ethyl acetate. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and

then concentrated. The residue was subjected to silica gel column chromatography to give 0.83 g of 4-chloro-6-benzylpyrimidine.

4-Chloro-6-benzylpyrimidine




5 $^1\text{H-NMR}$: 4.11 (s, 2H), 7.13 (s, 1H), 7.24-7.38 (m, 5H), 8.91 (s, 1H)

Reference Production Example 22

In 10 ml of tetrahydrofuran was suspended 1.3 g of zinc (powder), to which dibromoethane (2 drops) was added. The mixture was heated under reflux for 5 minutes, to which trimethylsilane chloride was added. The mixture was further heated under reflux for 5 minutes, to which a solution of 1.9 g of 2-fluorobenzyl bromide dissolved in 20 ml of tetrahydrofuran was slowly added with heating under reflux, followed by stirring for 20 minutes. (The solution thus obtained is referred to as solution B). In 10 ml of tetrahydrofuran were suspended 1.5 g of 4,6-dichloropyrimidine and 0.1 g of di-
15 chlorobistriphenylphosphine palladium, to which the above solution B was added, followed by heating under reflux for 3 hours and further stirring at room temperature for 12 hours. The reaction mixture was then poured into water and extracted three times with ethyl acetate. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate,
20 and then concentrated. The residue was subjected to silica gel column chromatography to give 0.63 g of 4-chloro-6-(2-fluorobenzyl)pyrimidine.

4-Chloro-6-(2-fluorobenzyl)pyrimidine




25 $^1\text{H-NMR}$: 4.05 (s, 2H), 6.70-7.08 (m, 3H), 7.17-7.23 (m, 2H), 8.81 (s, 1H)

Reference Production Example 23

A reaction vessel was charged with [1,1'-bis(diphenylphosphino)ferrocene dichloropalladium] methylene chloride complex, 1.007 g of 2,6-difluorophenylboronic acid, and 2.707 g of tripotassium phosphate n-hydrate, to which 16 ml of 1,2-dimethoxyethane, 4 ml of water, and 0.95 g of 4,6-dichloropyrimidine were added, followed by stirring at 80°C under an atmosphere of a nitrogen gas for 5 hours. The reaction mixture was then left for cooling to room temperature, and water was added to the reaction mixture, which was extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.241 g of 4-chloro-6-(2,6-difluorophenyl)pyrimidine.

4-Chloro-6-(2,6-difluorophenyl)pyrimidine




¹H-NMR: 7.06 (t, 2H), 7.49-7.53 (m, 1H), 7.58 (s, 1H), 9.13 (s, 1H)

Reference Production Example 24

In 10 ml of tetrahydrofuran was suspended 1.3 g of zinc (powder), to which dibromoethane (2 drops) was added. The mixture was heated under reflux for 5 minutes, to which trimethylsilane chloride was added. The mixture was further heated under reflux for 5 minutes, to which a solution of 2.1 g of 2-chlorobenzyl bromide dissolved in 20 ml of tetrahydrofuran was slowly added with heating under reflux, followed by stirring for 20 minutes. (The solution thus obtained is referred to as solution C). In 10 ml of tetrahydrofuran were suspended 1.5 g of 4,6-dichloropyrimidine and 0.1 g of di-

chlorobistriphenylphosphine palladium, to which the above solution C was added, followed by heating under reflux for 3 hours and further stirring at room temperature for 12 hours. The reaction mixture was then poured into water and extracted three times with ethyl acetate. The organic layers
5 were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.84 g of 4-chloro-6-(2-chlorobenzyl)pyrimidine.

4-Chloro-6-(2-chlorobenzyl)pyrimidine




10 $^1\text{H-NMR}$: 4.25 (s, 2H), 7.10 (s, 1H), 7.26-7.33 (m, 3H), 7.41-7.44 (m, 1H), 8.91 (s, 1H)

Reference Production Example 25

In 10 ml of tetrahydrofuran was suspended 1.3 g of zinc (powder), to which dibromoethane (2 drops) was added. The mixture was heated under
15 reflux for 5 minutes, to which trimethylsilane chloride was added. The mixture was further heated under reflux for 5 minutes, to which a solution of 1.9 g of 2-methylbenzyl bromide dissolved in 20 ml of tetrahydrofuran was slowly added with heating under reflux, followed by stirring for 20 minutes. (The solution thus obtained is referred to as solution D). In 10 ml of tetra-
20 hydrofuran were suspended 1.5 g of 4,6-dichloropyrimidine and 0.1 g of dichlorobistriphenylphosphine palladium, to which the above solution D was added, followed by heating under reflux for 3 hours and further stirring at room temperature for 12 hours. The reaction mixture was then poured into water and extracted three times with ethyl acetate. The organic layers
25 were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column

chromatography to give 0.55 g of 4-chloro-6-(2-methylbenzyl)pyrimidine.

4-Chloro-6-(2-methylbenzyl)pyrimidine




¹H-NMR: 2.24 (s, 3H), 4.13 (s, 2H), 6.98 (s, 1H), 7.18-7.23 (m, 4H),

5 8.91 (s, 1H)

Reference Production Example 26

In 10 ml of tetrahydrofuran was suspended 1.3 g of zinc (powder), to which dibromoethane (2 drops) was added. The mixture was heated under reflux for 5 minutes, to which trimethylsilane chloride was added. The mixture was further heated under reflux for 5 minutes, to which a solution of 2.5 g of 2,6-difluorobenzyl bromide dissolved in 20 ml of tetrahydrofuran was slowly added with heating under reflux, followed by stirring for 20 minutes. (The solution thus obtained is referred to as solution E). In 10 ml of tetrahydrofuran were suspended 1.5 g of 4,6-dichloropyrimidine and 0.1 g of dichlorobis(triphenyl)phosphine palladium, to which the above solution D was added, followed by heating under reflux for 3 hours and further stirring at room temperature for 12 hours. The reaction mixture was then poured into water and extracted three times with ethyl acetate. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.69 g of 4-chloro-6-(2,6-difluorobenzyl)pyrimidine.

4-Chloro-6-(2,6-difluorobenzyl)pyrimidine




¹H-NMR: 4.18 (s, 2H), 6.92-6.99 (m, 2H), 7.17 (s, 1H), 7.24-7.34 (m,

25 1H), 8.90 (s, 1H)

Reference Production Example 27

A reaction vessel was charged with 2.03 g of tetrakis(triphenylphosphine) palladium, 5.55 g of 2,3-difluorophenylboronic acid, and 14.9 g of tripotassium phosphate n-hydrate, to which 120 ml of 1,2-dimethoxyethane, 30 ml of water, and 5.20 g of 4,6-dichloropyrimidine were added, followed by stirring at 80°C under an atmosphere of a nitrogen gas for 12 hours. The reaction mixture was then left for cooling to room temperature, and water was added to the reaction mixture, which was extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 3.32 g of 4-chloro-6-(2,3-difluorophenyl)pyrimidine.

4-Chloro-6-(2,3-difluorophenyl)pyrimidine




¹H-NMR: 7.19-7.51 (m, 2H), 7.85-7.98 (m, 2H, involving a singlet at 7.89), 9.08 (s, 1H)

Reference Production Example 28

In 10 ml of tetrahydrofuran was suspended 1.3 g of zinc (powder), to which dibromoethane (2 drops) was added. The mixture was heated under reflux for 5 minutes, to which trimethylsilane chloride was added. The mixture was further heated under reflux for 5 minutes, to which a solution of 2.4 g of 2-trifluoromethylbenzyl bromide dissolved in 20 ml of tetrahydrofuran was slowly added with heating under reflux, followed by stirring for 20 minutes. (The solution thus obtained is referred to as solution F). In 10 ml of tetrahydrofuran were suspended 1.5 g of 4,6-dichloropyrimidine and

0.1 g of dichlorobistriphenylphosphine palladium, to which the above solution F was added, followed by heating under reflux for 3 hours and further stirring at room temperature for 12 hours. The reaction mixture was then poured into water and extracted three times with ethyl acetate. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.92 g of 4-chloro-6-(2-trifluoromethylbenzyl)pyrimidine.

4-Chloro-6-(2-trifluoromethylbenzyl)pyrimidine



10


¹H-NMR: 4.32 (s, 2H), 7.02 (s, 1H), 7.38-7.45 (m, 2H), 7.55 (t, 1H), 7.72 (d, 1H), 8.92 (s, 1H)

Reference Production Example 29

In 10 ml of tetrahydrofuran was suspended 1.3 g of zinc (powder), to which dibromoethane (2 drops) was added. The mixture was heated under reflux for 5 minutes, to which trimethylsilane chloride was added. The mixture was further heated under reflux for 5 minutes, to which a solution of 2.1 g of 2,3-difluorobenzyl bromide dissolved in 20 ml of tetrahydrofuran was slowly added with heating under reflux, followed by stirring for 20 minutes. (The solution thus obtained is referred to as solution G). In 10 ml of tetrahydrofuran were suspended 1.5 g of 4,6-dichloropyrimidine and 0.1 g of dichlorobistriphenylphosphine palladium, to which the above solution G was added, followed by heating under reflux for 3 hours and further stirring at room temperature for 12 hours. The reaction mixture was then poured into water and extracted three times with ethyl acetate. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate,

and then concentrated. The residue was subjected to silica gel column chromatography to give 0.93 g of 4-chloro-6-(2,3-difluorobenzyl)pyrimidine.

4-Chloro-6-(2,3-difluorobenzyl)pyrimidine




5 ¹H-NMR: 4.16 (s, 2H), 6.95-7.16 (m, 3H), 7.20 (s, 1H), 8.92 (s, 1H)

Reference Production Example 30

In 10 ml of tetrahydrofuran was suspended 1.3 g of zinc (powder), to which dibromoethane (2 drops) was added. The mixture was heated under reflux for 5 minutes, to which trimethylsilane chloride was added. The mixture was further heated under reflux for 5 minutes, to which a solution of 2.1 g of 2,4-difluorobenzyl bromide dissolved in 20 ml of tetrahydrofuran was slowly added with heating under reflux, followed by stirring for 20 minutes. (The solution thus obtained is referred to as solution H). In 10 ml of tetrahydrofuran were suspended 1.5 g of 4,6-dichloropyrimidine and 0.1 g of dichlorobistriphenylphosphine palladium, to which the above solution H was added, followed by heating under reflux for 3 hours and further stirring at room temperature for 12 hours. The reaction mixture was then poured into water and extracted three times with ethyl acetate. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.95 g of 4-chloro-6-(2,4-difluorobenzyl)pyrimidine.

4-Chloro-6-(2,4-difluorobenzyl)pyrimidine




¹H-NMR: 4.10 (s, 2H), 6.82-6.92 (m, 2H), 7.18 (s, 1H), 7.23-7.28 (m,

1H), 8.90 (s, 1H)

Reference Production Example 31

In 10 ml of tetrahydrofuran was suspended 1.3 g of zinc (powder), to which dibromoethane (2 drops) was added. The mixture was heated under reflux for 5 minutes, to which trimethylsilane chloride was added. The mixture was further heated under reflux for 5 minutes, to which a solution of 1.5 g of 3-fluorobenzyl bromide dissolved in 20 ml of tetrahydrofuran was slowly added with heating under reflux, followed by stirring for 20 minutes. (The solution thus obtained is referred to as solution I). In 10 ml of tetrahydrofuran were suspended 1.5 g of 4,6-dichloropyrimidine and 0.1 g of dichlorobistriphenylphosphine palladium, to which the above solution I was added, followed by heating under reflux for 3 hours and further stirring at room temperature for 12 hours. The reaction mixture was then poured into water and extracted three times with ethyl acetate. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.62 g of 4-chloro-6-(3-fluorobenzyl)pyrimidine.

4-Chloro-6-(3-fluorobenzyl)pyrimidine




¹H-NMR: 4.09 (s, 2H), 6.96-7.05 (m, 3H), 7.15 (s, 1H), 7.26-7.33 (m, 1H), 8.92 (s, 1H)

Reference Production Example 32

In 10 ml of tetrahydrofuran was suspended 1.3 g of zinc (powder), to which dibromoethane (2 drops) was added. The mixture was heated under reflux for 5 minutes, to which trimethylsilane chloride was added. The mixture was further heated under reflux for 5 minutes, to which a solution of

2.2 g of 2-chloro-6-fluorobenzyl bromide dissolved in 20 ml of tetrahydrofuran was slowly added with heating under reflux, followed by stirring for 20 minutes. (The solution thus obtained is referred to as solution J). In 10 ml of tetrahydrofuran were suspended 1.5 g of 4,6-dichloropyrimidine and 0.1 g of dichlorobistriphenylphosphine palladium, to which the above solution J was added, followed by heating under reflux for 3 hours and further stirring at room temperature for 12 hours. The reaction mixture was then poured into water and extracted three times with ethyl acetate. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.77 g of 4-chloro-6-(2-chloro-6-fluorobenzyl)pyrimidine.

4-Chloro-6-(2-chloro-6-fluorobenzyl)pyrimidine




¹H-NMR: 4.30 (s, 2H), 7.05-7.08 (m, 2H), 7.24-7.29 (m, 2H), 8.90 (s, 1H)

Reference Production Example 33

In 10 ml of tetrahydrofuran was suspended 1.3 g of zinc (powder), to which dibromoethane (2 drops) was added. The mixture was heated under reflux for 5 minutes, to which trimethylsilane chloride was added. The mixture was further heated under reflux for 5 minutes, to which a solution of 2.2 g of 3-chloro-2-fluorobenzyl bromide dissolved in 20 ml of tetrahydrofuran was slowly added with heating under reflux, followed by stirring for 20 minutes. (The solution thus obtained is referred to as solution K). In 10 ml of tetrahydrofuran were suspended 1.5 g of 4,6-dichloropyrimidine and 0.1 g of dichlorobistriphenylphosphine palladium, to which the above solu-

tion K was added, followed by heating under reflux for 3 hours and further stirring at room temperature for 12 hours. The reaction mixture was then poured into water and extracted three times with ethyl acetate. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.89 g of 4-chloro-6-(3-chloro-2-fluorobenzyl)pyrimidine.

4-Chloro-6-(3-chloro-2-fluorobenzyl)pyrimidine




¹H-NMR: 4.14 (s, 2H), 7.04-7.43 (m, 4H), 8.90 (s, 1H)

Reference Production Example 34

In 10 ml of tetrahydrofuran was suspended 1.3 g of zinc (powder), to which dibromoethane (2 drops) was added. The mixture was heated under reflux for 5 minutes, to which trimethylsilane chloride was added. The mixture was further heated under reflux for 5 minutes, to which a solution of 2.0 g of 2-bromobenzyl bromide dissolved in 20 ml of tetrahydrofuran was slowly added with heating under reflux, followed by stirring for 20 minutes. (The solution thus obtained is referred to as solution L). In 10 ml of tetrahydrofuran were suspended 1.5 g of 4,6-dichloropyrimidine and 0.1 g of dichlorobis(triphenyl)phosphine palladium, to which the above solution L was added, followed by heating under reflux for 3 hours and further stirring at room temperature for 12 hours. The reaction mixture was then poured into water and extracted three times with ethyl acetate. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.69 g of 4-chloro-6-(2-bromobenzyl)pyrimidine.

4-Chloro-6-(2-bromobenzyl)pyrimidine




¹H-NMR: 4.27 (s, 2H), 7.10 (s, 1H), 7.16-7.22 (m, 1H), 7.31-7.34 (m, 2H), 7.61 (d, 1H), 8.90 (s, 1H)

5 Reference Production Example 35

In 10 ml of tetrahydrofuran was suspended 1.3 g of zinc (powder), to which dibromoethane (2 drops) was added. The mixture was heated under reflux for 5 minutes, to which trimethylsilane chloride was added. The mixture was further heated under reflux for 5 minutes, to which a solution of 2.0 g of 1-(1-bromoethyl)-3-fluorobenzene dissolved in 20 ml of tetrahydrofuran was slowly added with heating under reflux, followed by stirring for 20 minutes. (The solution thus obtained is referred to as solution M). In 10 ml of tetrahydrofuran were suspended 1.5 g of 4,6-dichloropyrimidine and 0.2 g of tetrakis(triphenylphosphine) palladium, to which the above solution M was added, followed by heating under reflux for 3 hours and further stirring at room temperature for 12 hours. The reaction mixture was then poured into water and extracted three times with ethyl acetate. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.43 g of 4-chloro-6-(1-(3-fluorophenyl)ethyl)pyrimidine.

4-Chloro-6-(1-(3-fluorophenyl)ethyl)pyrimidine




¹H-NMR: 1.68 (d, 3H), 4.21 (q, 1H), 6.90-7.11 (m, 3H), 7.15 (s, 1H), 7.28-7.36 (m, 1H), 8.93 (s, 1H)

Reference Production Example 36

In 10 ml of tetrahydrofuran was suspended 1.3 g of zinc (powder), to which dibromoethane (2 drops) was added. The mixture was heated under reflux for 5 minutes, to which trimethylsilane chloride was added. The mixture was further heated under reflux for 5 minutes, to which a solution of 2.0 g of 1-(1-bromoethyl)-2-fluorobenzene dissolved in 20 ml of tetrahydrofuran was slowly added with heating under reflux, followed by stirring for 20 minutes. (The solution thus obtained is referred to as solution N). In 10 ml of tetrahydrofuran were suspended 1.5 g of 4,6-dichloropyrimidine and 0.2 g of tetrakis(triphenyl)phosphine palladium, to which the above solution N was added, followed by heating under reflux for 3 hours and further stirring at room temperature for 12 hours. The reaction mixture was then poured into water and extracted three times with ethyl acetate. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.67 g of 4-chloro-6-(1-(2-fluorophenyl)ethyl)pyrimidine.

20 4-Chloro-6-(1-(2-fluorophenyl)ethyl)pyrimidine




¹H-NMR: 1.68 (d, 3H), 4.50 (q, 1H), 7.04-7.37 (m, 5H, involving a singlet at 7.20), 8.92 (s, 1H)

Reference Production Example 37

25 In 10 ml of tetrahydrofuran was suspended 1.3 g of zinc (powder), to which dibromoethane (2 drops) was added. The mixture was heated under

reflux for 5 minutes, to which trimethylsilane chloride was added. The mixture was further heated under reflux for 5 minutes, to which a solution of 2.4 g of 2-chloro-5-methyl-6-fluorobenzyl bromide dissolved in 20 ml of tetrahydrofuran was slowly added with heating under reflux, followed by stirring for 20 minutes. (The solution thus obtained is referred to as solution O). In 10 ml of tetrahydrofuran were suspended 1.5 g of 4,6-dichloropyrimidine and 0.1 g of dichlorobis(triphenyl)phosphine palladium, to which the above solution L was added, followed by heating under reflux for 3 hours and further stirring at room temperature for 12 hours. The reaction mixture was then poured into water and extracted three times with ethyl acetate. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.53 g of 4-chloro-6-(2-chloro-5-methyl-6-fluorobenzyl)pyrimidine.

4-Chloro-6-(2-chloro-5-methyl-6-fluorobenzyl)pyrimidine




¹H-NMR: 2.38 (s, 3H), 4.34 (s, 2H), 7.02 (t, 1H), 7.07 (s, 1H), 7.24 (dd, 1H), 7.27 (s, 1H), 8.91 (s, 1H)

Reference Production Example 38

In 10 ml of tetrahydrofuran was suspended 1.3 g of zinc (powder), to which dibromoethane (2 drops) was added. The mixture was heated under reflux for 5 minutes, to which trimethylsilane chloride was added. The mixture was further heated under reflux for 5 minutes, to which a solution of 2.4 g of 2-chloro-3,6-difluorobenzyl bromide dissolved in 20 ml of tetrahydrofuran was slowly added with heating under reflux, followed by stirring for 20

minutes. (The solution thus obtained is referred to as solution P). In 10 ml of tetrahydrofuran were suspended 1.5 g of 4,6-dichloropyrimidine and 0.1 g of dichlorobistriphenylphosphine palladium, to which the above solution P was added, followed by heating under reflux for 3 hours and further stirring at room temperature for 12 hours. The reaction mixture was then poured into water and extracted three times with ethyl acetate. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.64 g of 4-chloro-6-(2-chloro-3,6-difluorobenzyl)pyrimidine.


4-Chloro-6-(2-chloro-3,6-difluorobenzyl)pyrimidine



¹H-NMR: 4.32 (s, 2H), 7.02-7.19 (m, 3H, involving a singlet at 7.13), 8.90 (s, 1H)

Reference Production Example 39

In 10 ml of N,N-dimethylformamide were dissolved 608 mg of 4-chloro-6-(2-fluorophenyl)pyrimidine and 527 mg of 2-butyne-1,4-diol, to which 245 mg of sodium hydride (60% in oil) was added, followed by stirring at room temperature for 9 hours. The reaction mixture was then poured into water and extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then concentrated. The resulting residue was subjected to silica gel column chromatography to give 465 mg of 4-(2-fluorophenyl)-6-(4-hydroxy-2-butyloxy)pyrimidine.




¹H-NMR: 1.68 (t, 1H), 4.34 (dt, 2H), 5.13 (t, 2H), 7.18 (dt, 1H), 7.23-7.51 (m, 2H), 7.33 (s, 1H), 8.12 (dt, 1H), 8.88 (s, 1H)

Reference Production Example 40

5 In 10 ml of tetrahydrofuran was suspended 1.3 g of zinc (powder), to which dibromoethane (2 drops) was added. The mixture was heated under reflux for 5 minutes, to which trimethylsilane chloride was added. The mixture was further heated under reflux for 5 minutes, to which a solution of 1.9 g of 4-fluorobenzyl bromide dissolved in 20 ml of tetrahydrofuran was
10 slowly added with heating under reflux, followed by stirring for 20 minutes. (The solution thus obtained is referred to as solution Q). In 10 ml of tetrahydrofuran were suspended 1.5 g of 4,6-dichloropyrimidine and 0.1 g of dichlorobis(triphenylphosphine) palladium, to which the above solution Q was added, followed by heating under reflux for 3 hours and further stirring at
15 room temperature for 12 hours. The reaction mixture was then poured into water and extracted three times with ethyl acetate. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.72 g of 4-chloro-6-(4-fluorobenzyl)pyrimidine.

20 4-Chloro-6-(4-fluorobenzyl)pyrimidine




¹H-NMR: 4.07 (s, 2H), 6.94-7.24 (m, 5H), 8.91 (s, 1H)

Reference Production Example 41

A reaction vessel was charged with [1,1'-bis(diphenylphosphino)ferrocene dichloropalladium] methylene chloride complex, 984 mg of 2,6-difluorophenylboronic acid, and 1.32 g of sodium carbonate, to which 15 ml of toluene, 4 ml of ethanol, 4 ml of water, and 997 mg of 2-methyl-4,6-dichloro-
5 pyrimidine were added, followed by stirring at 80°C under an atmosphere of a nitrogen gas for 6 hours. The reaction mixture was then left for cooling to room temperature, and water was added to the reaction mixture, which was extracted with ethyl acetate. The organic layer was washed with a saturated aqueous sodium chloride solution, dried over anhydrous magnesium
10 sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 656 mg of 4-chloro-2-methyl-6-(2,3-difluorophenyl)pyrimidine.

4-Chloro-2-methyl-6-(2,3-difluorophenyl)pyrimidine




15 ¹H-NMR: 2.79 (s, 3H), 7.15-7.48 (m, 2H), 7.67 (s, 1H), 7.85-7.95 (m, 1H)

Reference Production Example 42

In 4.2 ml of tetrahydrofuran was suspended 0.12 g of sodium hydride (60% in oil), to which 0.8 ml of a tetrahydrofuran solution containing 0.5 g of
20 4-chloro-6-(2,3-difluorophenylamino)pyrimidine was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 0.8 ml of a tetrahydrofuran solution containing 0.48 g of iodoethane was slowly added dropwise at 0°C, followed by further stirring for 8 hours. The reaction mixture was then poured into a
25 saturated aqueous ammonium chloride solution and extracted three times

with t-butyl methyl ether. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.23 g of 4-chloro-6-(N-ethyl-N-(2,3-difluorophenyl)amino)pyrimidine.

5 4-Chloro-6-(N-ethyl-N-(2,3-difluorophenyl)amino)pyrimidine




¹H-NMR: 1.25 (t, 3H), 3.96 (q, 2H), 6.72 (s, 1H), 7.00-7.30 (m, 2H), 7.64 (dd, 1H), 8.55 (s, 1H)

Reference Production Example 43

10 In 6.7 ml of tetrahydrofuran was suspended 0.16 g of sodium hydride (60% in oil), to which 0.8 ml of a tetrahydrofuran solution containing 0.5 g of 4-chloro-6-(3-fluorophenylamino)pyrimidine was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes, to which 0.8 ml of a tetrahydrofuran solution containing
15 0.42 g of iodoethane was slowly added at 0°C, followed by further stirring for 8 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with t-butyl methyl ether. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was
20 subjected to silica gel column chromatography to give 0.43 g of 4-chloro-6-(N-ethyl-N-(3-fluorophenyl)amino)pyrimidine.

4-Chloro-6-(N-ethyl-N-(3-fluorophenyl)amino)pyrimidine

231




¹H-NMR: 1.22 (t, 3H), 3.99 (q, 2H), 6.17 (s, 1H), 6.93-7.15 (m, 3H), 7.43-7.52 (m, 1H), 8.46 (s, 1H)

Reference Production Example 44

5 In 4.7 ml of tetrahydrofuran was suspended 0.12 g of sodium hydride (60% in oil), to which 1 ml of a tetrahydrofuran solution containing 0.45 g of 2-chlorocyclohexanol was slowly added dropwise with stirring at 0°C. The mixture was stirred at 0°C for 10 minutes, to which 1 ml of a tetrahydrofuran solution containing 0.5 g of 4,6-dichloropyrimidine was added at 0°C,
10 followed by further stirring for 20 minutes. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with ethyl acetate. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.38 g
15 of 4-chloro-6-(2-chlorocyclohexyloxy)pyrimidine.

4-Chloro-6-(2-chlorocyclohexyloxy)pyrimidine



¹H-NMR: 1.44-1.49 (m, 2H), 1.73-1.82 (m, 2H), 2.15-2.30 (m, 2H), 3.91-4.08 (m, 1H), 5.19-5.31 (m, 1H), 6.79 (s, 1H), 8.56 (s, 1H)


20 Reference Production Example 45

In 6.7 ml of tetrahydrofuran was suspended 0.18 g of sodium hydride (60% in oil), to which 1 ml of a tetrahydrofuran solution containing 0.42 g of 2-methylcyclohexanol (cis : trans = 3 : 7) was slowly added dropwise with

stirring at 0°C. The mixture was stirred at 0°C for 10 minutes, to which 1 ml of a tetrahydrofuran solution containing 0.5 g of 4,6-dichloropyrimidine was added at 0°C, followed by further stirring at room temperature for 2 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with t-butyl methyl ether. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.55 g of 4-chloro-6-(2-methylcyclohexyloxy)pyrimidine as a mixture of cis and trans forms.

10

4-Chloro-6-(2-methylcyclohexyloxy)pyrimidine



Cis Form:

¹H-NMR: 0.92 (d, 3H), 1.33-2.00 (m, 9H), 5.26-5.31 (m, 1H), 6.76 (s, 1H), 8.53 (s, 1H)

15

Trans Form:

¹H-NMR: 0.93 (d, 3H), 1.11-1.38 (m, 4H), 1.65-1.83 (m, 4H), 2.10-2.14 (m, 1H), 4.78-4.85 (m, 1H), 6.72 (s, 1H), 8.53 (s, 1H)


Reference Production Example 46

In 5 ml of tetrahydrofuran was suspended 0.11 g of sodium hydride (60% in oil), to which 1 ml of a tetrahydrofuran solution containing 0.22 g of trans-2-methylcyclopentanol was slowly added dropwise with stirring at 0°C. The mixture was stirred at 0°C for 10 minutes, to which 1 ml of a tetrahydrofuran solution containing 0.3 g of 4,6-dichloropyrimidine was added at 0°C, followed by further stirring for 2 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with t-butyl methyl ether. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then con-

25

centrated. The residue was subjected to silica gel column chromatography to give 0.38 g of 4-chloro-6-(trans-2-methylcyclopentyloxy)pyrimidine.

4-Chloro-6-(trans-2-methylcyclopentyloxy)pyrimidine




5 ¹H-NMR: 1.05 (d, 3H), 1.21-1.32 (m, 1H), 1.68-1.81 (m, 3H), 1.93-2.00 (m, 1H), 2.01-2.18 (m, 1H), 4.99-5.04 (m, 1H), 6.72 (s, 1H), 8.55 (s, 1H)

Reference Production Example 47

In 4 ml of tetrahydrofuran was suspended 0.11 g of sodium hydride (60% in oil), to which 1 ml of a tetrahydrofuran solution containing 0.25 g of cycloheptanol was slowly added dropwise with stirring at 0°C. The mixture was stirred at 0°C for 10 minutes, to which 1 ml of a tetrahydrofuran solution containing 0.3 g of 4,6-dichloropyrimidine was added at 0°C, followed by further stirring for 2 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with t-butyl methyl ether. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.43 g of 4-chloro-6-cycloheptyloxypyrimidine.

4-Chloro-6-cycloheptyloxypyrimidine



20


¹H-NMR: 1.50-1.83 (m, 10H), 1.99-2.05 (m, 2H), 5.27-5.35 (m, 1H), 6.70 (s, 1H), 8.54 (s, 1H)

Reference Production Example 48

First, 18.9 g of 28% sodium methoxide in methanol was rapidly

mixed with 4.7 g of formamide, and the mixture was heated under reflux, to which 30 ml of a methanol solution containing 5 g of ethyl methylmalonate was slowly added dropwise over 3 hours. The mixture was further heated under reflux with stirring for 10 hours and then left for cooling, and the suspension was concentrated under reduced pressure. Then, 10 ml of water was added to the residue, which was acidified with concentrated hydrochloric acid. The resulting precipitate was filtered by suction and dried under reduced pressure to give 2.8 g of 4,6-dihydroxy-5-methylpyrimidine.

4,6-Dihydroxy-5-methylpyrimidine




¹H-NMR (DMSO-d₆): 1.73 (s, 3H), 7.90 (s, 1H)

Reference Production Example 49

A reaction vessel was charged at 0°C with 2.8 g of 4,6-dihydroxy-5-methylpyrimidine, 5.00 g of phosphorus oxychloride, and 3.3 g of diisopropylethylamine, followed by stirring at 80°C for 4 hours. The mixture was then left for cooling and poured into ice water. The mixture was further stirred for 30 minutes and then extracted with ethyl acetate. The organic layers were combined and washed with a saturated aqueous sodium hydrogencarbonate solution and then with a saturated aqueous sodium chloride solution, and the organic layers were then dried over anhydrous magnesium sulfate and concentrated. The residue was subjected to silica gel column chromatography to give 2.4 g of 4,6-dichloro-5-methylpyrimidine.

4,6-Dichloro-5-methylpyrimidine




¹H-NMR: 2.50 (s, 3H), 8.63 (s, 1H)

Reference Production Example 50

In 12 ml of tetrahydrofuran was suspended 0.32 g of sodium hydride (60% in oil), to which 2 ml of a tetrahydrofuran solution containing 0.43 g of 2-butyn-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes and then cooled to 0°C, to which 2 ml of a tetrahydrofuran solution containing 1 g of 4,6-dichloro-5-methylpyrimidine was slowly added, followed by further stirring at 0°C for 2 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with t-butyl methyl ether. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 1.1 g of 4-chloro-6-(2-butynyloxy)-5-methylpyrimidine.

15 4-Chloro-6-(2-butynyloxy)-5-methylpyrimidine




¹H-NMR: 1.88 (t, 3H), 2.26 (s, 3H), 5.00 (q, 2H), 8.44 (s, 1H)

Reference Production Example 51

In 4 ml of tetrahydrofuran was suspended 0.12 g of sodium hydride (60% in oil), to which 1 ml of a tetrahydrofuran solution containing 0.28 g of cis-2-methylcyclohexanol was slowly added dropwise with stirring at 0°C. The mixture was stirred at 0°C for 10 minutes, to which 1 ml of a tetrahydrofuran solution containing 0.3 g of 4,6-dichloropyrimidine was added at 0°C, followed by further stirring at room temperature for 3 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with t-butyl methyl ether. The

organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.12 g of 4-chloro-6-(cis-2-methylcyclohexyloxy)pyrimidine.

5 4-Chloro-6-(cis-2-methylcyclohexyloxy)pyrimidine




¹H-NMR: 0.92 (d, 3H), 1.33-2.00 (m, 9H), 5.26-5.31 (m, 1H), 6.76 (s, 1H), 8.53 (s, 1H)

Reference Production Example 52

10 In 4 ml of tetrahydrofuran was suspended 0.11 g of sodium hydride (60% in oil), to which 1 ml of a tetrahydrofuran solution containing 0.25 g of trans-2-methylcyclohexanol was slowly added dropwise with stirring at 0°C. The mixture was stirred at 0°C for 10 minutes, to which 1 ml of a tetrahydrofuran solution containing 0.3 g of 4,6-dichloropyrimidine was added at
15 0°C, followed by further stirring at room temperature for 3 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with ethyl acetate. The organic layers were combined, washed with water, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel
20 column chromatography to give 0.18 g of 4-chloro-6-(trans-2-methylcyclohexyloxy)pyrimidine.

4-Chloro-6-(trans-2-methylcyclohexyloxy)pyrimidine




¹H-NMR: 0.93 (d, 3H), 1.11-1.38 (m, 4H), 1.65-1.83 (m, 4H), 2.10-2.14
25 (m, 1H), 4.78-4.85 (m, 1H), 6.72 (s, 1H), 8.53 (s, 1H)

Reference Production Example 53

Under an atmosphere of a nitrogen gas, 2.98 g of 4,6-dichloro-pyrimidine, 1.81 g of 3,3-dimethyl-1-butyne, 351 mg of dichlorobis(triphenylphosphine)palladium, 381 mg of copper iodide, 4.05 g of triethylamine and 525 mg of triphenylphosphine was added to 20 ml of acetonitrile and the suspension was stirred for 6 hours at 45°C. After cooling the reaction mixture was diluted with t-butyl methyl ether and washed three times with water. The organic layers were dried over sodium sulfate and concentrated. The residue was subjected to silica gel thin layer chromatography to give 1.86 g of 4-chloro-6-(3,3-dimethyl-1-butynyl)pyrimidine.

10 4-Chloro-6-(3,3-dimethyl-1-butynyl)pyrimidine




¹H-NMR: 1.35 (s, 9H), 7.37 (d, 1H), 8.90 (s, 1H)

Reference Production Example 54

In 18 ml of tetrahydrofuran was suspended 0.56 g of sodium hydride (60% in oil), to which 2 ml of a tetrahydrofuran solution containing 0.8 g of 2-butyne-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes and then cooled to 0°C, to which 3 ml of a tetrahydrofuran solution containing 2 g of 4,5,6-trichloropyrimidine was slowly added dropwise, followed by further stirring at 0°C for 2 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with ethyl acetate. The organic layers were combined, washed with brine, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 2.23 g of 4,5-dichloro-6-(2-butynyloxy)pyrimidine.

4,5-Dichloro-6-(2-butynyloxy)pyrimidine




¹H-NMR: 1.88 (t, 3H), 5.08 (q, 2H), 8.48 (s, 1H)

Reference Production Example 55

5 In 30 ml of tetrahydrofuran was suspended 1.78 g of sodium hydride (60% in oil), to which 3 ml of a tetrahydrofuran solution containing 2.35 g of 1,2-cyclohexanediol (mixture of cis-form and trans-form) was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 10 minutes and then cooled to 0°C, to which 7 ml of a
10 tetrahydrofuran solution containing 3 g of 4,6-dichloropyrimidine was slowly added dropwise, followed by further stirring at 0°C for 3 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with ethyl acetate. The organic layers were combined, washed with brine, dried over anhydrous magnesium sulfate,
15 and then concentrated. The residue was subjected to silica gel column chromatography to give 1.6 g of 4-chloro-6-(2-hydroxycyclohexyloxy)pyrimidine.

4-Chloro-6-(2-hydroxycyclohexyloxy)pyrimidine




20 ¹H-NMR: 1.37-1.45 (m, 2H), 1.59-2.04 (m, 6H), 2.48 (bs, 1H), 3.99-4.02 (m, 1H), 5.30-5.35 (m, 1H), 6.80 (s, 1H), 8.54 (s, 1H) with peaks due to the minor trans-form at 2.55 (bs), 3.64-3.75 (m), 4.93-5.00 (m), 6.78 (s)

Reference Production Example 56

In 2 ml of tetrahydrofuran was suspended 0.05 g of sodium hydride

(60% in oil), to which 0.5 ml of a tetrahydrofuran solution containing 0.12 g of 2-butyne-1-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes and then cooled to 0°C, to which 0.5 ml of a tetrahydrofuran solution containing 0.25 g of 4,6-dichloro-5-fluoropyrimidine was slowly added dropwise, followed by further stirring at 0°C for 30 minutes. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with t-butyl methyl ether. The organic layers were combined, washed with brine, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.30 g of 4-chloro-5-fluoro-6-(2-butynyloxy)pyrimidine.

4-Chloro-5-fluoro-6-(2-butynyloxy)pyrimidine




¹H-NMR: 1.88 (t, 3H), 5.08 (q, 2H), 8.37 (s, 1H)

Reference Production Example 57

In 5 ml of tetrahydrofuran was suspended 0.19 g of sodium hydride (60% in oil), to which 0.5 ml of a tetrahydrofuran solution containing 0.48 g of 2,3-dimethylcyclohexanol (mixture of isomers) was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 10 minutes and then cooled to 0°C, to which 1.5 ml of a tetrahydrofuran solution containing 0.5 g of 4,6-dichloropyrimidine was slowly added dropwise, followed by further stirring at room temperature for 3 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with t-butyl methyl ether. The organic layers were combined, washed with brine, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to

silica gel column chromatography to give 0.72 g of 4-chloro-6-(2,3-dimethylcyclohexyloxy)pyrimidine.

4-Chloro-6-(2,3-dimethylcyclohexyloxy)pyrimidine




5 ¹H-NMR: 0.84-2.14 (m, 14H), 4.74-4.89 (m, 1H), 6.72 (s, 1H), 8.53 (s, 1H) with peaks due to the minor isomers at 5.10-5.21 (m)

Reference Production Example 58

In 5 ml of tetrahydrofuran was suspended 0.14 g of sodium hydride (60% in oil), to which 0.5 ml of a tetrahydrofuran solution containing 0.34 g
10 of 3-methylcyclohexanol (mixture of cis-form and trans-form) was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 10 minutes and then cooled to 0°C, to which 1.5 ml of a tetrahydrofuran solution containing 0.4 g of 4,6-dichloropyrimidine was slowly added dropwise, followed by further stirring at room temperature
15 for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with t-butyl methyl ether. The organic layers were combined, washed with brine, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.31 g of 4-chloro-
20 6-(3-methylcyclohexyloxy)pyrimidine.

4-Chloro-6-(3-methylcyclohexyloxy)pyrimidine




¹H-NMR: 0.85-2.12 (m, 12H), 5.02-5.12 (m, 1H), 6.70 (s, 1H), 8.54 (s, 1H) with peaks due to the minor isomer at 5.42-5.51 (m), 6.74 (s)

Reference Production Example 59

To 2 ml of ethanol were added 0.3 g of 4,5,6-trichloropyrimidine and 0.5 g of N-ethylaniline, followed by heating under reflux for 8 hours. The reaction mixture was then left for cooling to room temperature and concentrated under reduced pressure. The residue was added water and extracted three times with t-butyl methyl ether. The organic layers were combined, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.5 g of 4,5-dichloro-6-(N-ethyl-N-phenylamino)pyrimidine.

4,5-Dichloro-6-(N-ethyl-N-phenylamino)pyrimidine




¹H-NMR: 1.23 (t, 3H), 4.04 (q, 2H), 7.07 (d, 2H), 7.25 (t, 1H), 7.34 (t, 2H), 8.40 (s, 1H)

Reference Production Example 60

In 4 ml of tetrahydrofuran was suspended 0.1 g of sodium hydride (60% in oil), to which 0.5 ml of a tetrahydrofuran solution containing 0.23 g of cis-4-methylcyclohexanol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 10 minutes and then cooled to 0°C, to which 1 ml of a tetrahydrofuran solution containing 0.3 g of 4,6-dichloropyrimidine was slowly added dropwise, followed by further stirring at room temperature for 4 hours. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with t-butyl methyl ether. The organic layers were combined, washed with brine, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chroma-

tography to give 0.37 g of 4-chloro-6-(cis-4-methylcyclohexyloxy)pyrimidine.

4-Chloro-6-(cis-4-methylcyclohexyloxy)pyrimidine




¹H-NMR: 0.94 (d, 3H), 1.26-1.67 (m, 7H), 1.95-2.02 (m, 2H), 5.30-5.39
5 (m, 1H), 6.76 (s, 1H), 8.54 (s, 1H)

Reference Production Example 61

In 4 ml of tetrahydrofuran was suspended 0.1 g of sodium hydride (60% in oil), to which 0.5 ml of a tetrahydrofuran solution containing 0.23 g of trans-4-methylcyclohexanol was slowly added dropwise with stirring at
10 room temperature. The mixture was stirred at room temperature for 10 minutes and then cooled to 0°C, to which 1 ml of a tetrahydrofuran solution containing 0.3 g of 4,6-dichloropyrimidine was slowly added dropwise, followed by further stirring at room temperature for 4 hours. The reaction
15 mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with t-butyl methyl ether. The organic layers were combined, washed with brine, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.37 g of 4-chloro-6-(trans-4-methylcyclohexyloxy)pyrimidine.

20 4-Chloro-6-(trans-4-methylcyclohexyloxy)pyrimidine




¹H-NMR: 0.94 (d, 3H), 1.03-1.17 (m, 2H), 1.39-1.52 (m, 3H), 1.76-1.81
(m, 2H), 2.07-2.13 (m, 2H), 4.98-5.08 (m, 1H), 6.69 (s, 1H), 8.53 (s, 1H)

Reference Production Example 62

Reference Production Example 61

In 4 ml of tetrahydrofuran was suspended 0.2 g of sodium hydride (60% in oil), to which 1 ml of a tetrahydrofuran solution containing 0.39 g of cis-1,2-cyclohexanediol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 10 minutes and then cooled to 0°C, to which 2 ml of a tetrahydrofuran solution containing 0.5 g of 4,6-dichloropyrimidine was slowly added dropwise, followed by further stirring at 0°C for 30 minutes. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with ethyl acetate. The organic layers were combined, washed with brine, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.42 g of 4-chloro-6-(cis-2-hydroxycyclohexyloxy)pyrimidine.

4-Chloro-6-(cis-2-hydroxycyclohexyloxy)pyrimidine




¹H-NMR: 1.37-1.45 (m, 2H), 1.59-2.04 (m, 6H), 2.48 (bs, 1H), 3.99-4.02 (m, 1H), 5.30-5.35 (m, 1H), 6.80 (s, 1H), 8.54 (s, 1H)

Reference Production Example 63

To 3.2 ml of tetrahydrofuran were added 0.37 g of 4-chloro-6-(cis-2-hydroxycyclohexyloxy)pyrimidine and 0.45 ml of triethylamine, to which 0.25 ml of chlorotrimethylsilane was slowly added dropwise with stirring at 0°C. After stirring at room temperature for 3 hours, the reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with ethyl acetate. The organic layers were combined, washed with brine, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography.

phy to give 0.5 g of 4-chloro-6-(cis-2-trimethylsilanyloxycyclohexyloxy)pyrimidine.

4-Chloro-6-(cis-2-trimethylsilanyloxycyclohexyloxy)pyrimidine




5 ¹H-NMR: 0.14 (s, 9H), 1.12-1.36 (m, 2H), 1.48-1.73 (m, 5H), 1.88-1.95 (m, 1H), 3.94-3.98 (m, 1H), 5.12-5.16 (m, 1H), 6.69 (s, 1H), 8.46 (s, 1H)

Reference Production Example 64

In 2 ml of tetrahydrofuran was suspended 0.63 g of zinc (powder), to which dibromoethane (1 drop) was added. The mixture was heated under
10 reflux for 5 minutes, to which chlorotrimethylsilane (1 drop) was added. The mixture was further heated under reflux for 5 minutes, to which a solution of 0.91 g of 2-fluorobenzylbromide dissolved in 4 ml of tetrahydrofuran was slowly added with heating under reflux, followed by stirring for 20 minutes. (The solution thus obtained is referred to as solution R). In 4 ml
15 of tetrahydrofuran were suspended 0.8 g of 4,6-dichloropyrimidine and 0.02 g of dichlorobistriphenylphosphine palladium, to which the above solution R was added, followed by heating under reflux for 3 hours and further stirring at room temperature for 12 hours. The reaction mixture was then poured into water and extracted three times with t-butyl methyl ether. The organic
20 layers were combined, washed with brine, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.46 g of 4-chloro-5-fluoro-6-(2-fluorobenzyl)pyrimidine.

4-Chloro-5-fluoro-6-(2-fluorobenzyl)pyrimidine

245




¹H-NMR: 4.23 (s, 2H), 7.01-7.12 (m, 2H), 7.21-7.30 (m, 2H), 8.66 (s, 1H)

Reference Production Example 65

- 5 In 1 ml of tetrahydrofuran was suspended 0.04 g of sodium hydride (60% in oil), to which 0.5 ml of a tetrahydrofuran solution containing 0.06 g of 3-pentyn-2-ol was slowly added dropwise with stirring at room temperature. The mixture was stirred at room temperature for 20 minutes and then cooled to 0°C, to which 0.5 ml of a tetrahydrofuran solution containing
- 10 0.1 g of 4,6-dichloropyrimidine was slowly added dropwise, followed by further stirring for 30 minutes. The reaction mixture was then poured into a saturated aqueous ammonium chloride solution and extracted three times with t-butyl methyl ether. The organic layers were combined, washed with brine, dried over anhydrous magnesium sulfate, and then concentrated.
- 15 The residue was subjected to silica gel column chromatography to give 0.12 g of 4-chloro-6-(1-methyl-2-butynyloxy)pyrimidine.

4-Chloro-6-(1-methyl-2-butynyloxy)pyrimidine




- ¹H-NMR: 1.60 (d, 3H), 1.84 (d, 3H), 5.74-5.82 (m, 1H), 6.78 (s, 1H),
- 20 8.60 (s, 1H)

Reference Production Example 66

In 1 ml of tetrahydrofuran was suspended 0.03 g of sodium hydride (60% in oil), to which 0.5 ml of a tetrahydrofuran solution containing 0.06 g of 2-pentyn-1-ol was slowly added dropwise with stirring at room tempera-

ture. The mixture was stirred at room temperature for 20 minutes and then cooled to 0°C, to which 0.5 ml of a tetrahydrofuran solution containing 0.1 g of 4,6-dichloropyrimidine was slowly added dropwise, followed by further stirring for 30 minutes. The reaction mixture was then poured into
5 a saturated aqueous ammonium chloride solution and extracted three times with t-butyl methyl ether. The organic layers were combined, washed with brine, dried over anhydrous magnesium sulfate, and then concentrated. The residue was subjected to silica gel column chromatography to give 0.12 g of 4-chloro-6-(2-pentynyloxy)pyrimidine.

10 4-Chloro-6-(2-pentynyloxy)pyrimidine



¹H-NMR: 1.17 (t, 3H), 2.25 (br. q, 2H), 5.01 (t, 2H), 6.82 (s, 1H), 8.60 (s, 1H)

The following will describe formulation examples, in which parts are
15 by weight and the present compounds are designated by their compound numbers shown in the above tables.

Formulation Example 1 Emulsifiable concentrates

Nine parts of each of the present compounds (1) to (191) is dissolved in 37.5 parts of xylene and 37.5 parts of dimethylformamide, and 10 parts of
20 polyoxyethylene styryl phenyl ether and 6 parts of calcium dodecylbenzene-sulfonate are added thereto, followed by well mixing with stirring, to give an emulsifiable concentrate for each compound.

Formulation Example 2 Wettable powders

Nine parts of each of the present compounds (1) to (191) is added to a
25 mixture of 4 parts of sodium lauryl sulfate, 4 parts of calcium lignin sulfonate, 20 parts of synthetic hydrated silicon oxide fine powder, and 65 parts of

diatomaceous earth, followed well mixing with stirring, to give a wettable powder for each compound.

Formulation Example 3 Granules

Three parts of each of the present compounds (1) to (191), 5 parts of
5 synthetic hydrated silicon oxide fine powder, 5 parts of sodium dodecylbenzenesulfonate, 30 parts of bentonite, and 57 parts of clay are well mixed with stirring, and an appropriate amount of water is added to the mixture of these ingredients, followed by further stirring, granulation with a granulator, and drying by ventilation, to give a granule for each compound.

10 Formulation Example 4 Dusts

First, 4.5 parts of each of the present compounds (1) to (191), 1 part of synthetic hydrated silicon oxide fine powder, 1 part of Doriresu B (available from Sankyo Co., Ltd.) as a flocculant, and 7 parts of clay are well mixed in a mortar and then mixed with stirring in a juicer. To the resulting mixture is
15 added 86.5 parts by cut clay, followed by well mixing with stirring, to give a dust for each compound.

Formulation Example 5

Ten parts of each of the present compounds (1) to (191), 35 parts of white carbon containing 50 parts of polyoxyethylene alkyl ether sulfate ammonium salt, and 55 parts of water are mixed and pulverized by wet grinding method to give a formulation for each compound.
20

Formulation Example 6 Oil sprays

First, 0.5% by weight of each of the present compounds (1) to (191) is dissolved in 10% by weight of dichloromethane, and this solution is mixed
25 with 89.5% by weight of ISOPAR M (isoparaffin: registered trade name of Exxon Chemical Japan Ltd.) to give a 0.5% oil spray for each compound.

Formulation Example 7: Oil-based Aerosols

First, 0.1% by weight of each of the present compounds (1) to (191)

and 49.9% by weight of NEO-CHIOZOL (available from Chuo Kasei Co., Ltd.) are placed in an aerosol bomb, which is fitted with an aerosol valve and then filled with 25% by weight of dimethyl ether and 25% by weight of LPG, followed by shaking and fitting with an actuator, to give an oil-based aerosol.

5 Formulation Example 8 Water-based aerosols

 An aerosol vessel is filled with a solution prepared by mixing 0.6% by weight of each of the present compounds (1) to (191), 0.01% by weight of BHT, 5% by weight of xylene, 3.3 parts of deodorized kerosine, and 1% by weight of an emulsifier {ATMOS 300 (registered trade name of Atlas Chemical Co.)};
10 and with 50% by weight of distilled water. The vessel is fitted with a valve part, through which 40% by weight of a propellant (LPG) is filled into the vessel under increased pressure, to give a water-based aerosol for each compound.

 The following test examples are provided for demonstrating that the
15 present compounds are useful as the active ingredients of pesticides. The present compounds are designated by their compound numbers shown in the above tables.

 Test Example 1 Insecticidal Test against cotton aphids (*Aphis gossypii*)

20 A formulation of a test compound obtained in Formulation Example 5 was diluted with water so that the active ingredient concentration came to 500 ppm to prepare a test spray solution.

 The seeds of cucumber were planted in polyethylene cups and grown until their first foliage leaves developed, on which about 20 cotton aphids
25 (*Aphis gossypii*) were made parasitic. After one day, the test spray solution was sprayed at the rate of 20 ml/cup onto the cucumber plants. On the 6th day after the application, the number of cotton aphids was examined and the control value was determined by the following formula:

$$\text{Control value (\%)} = \{1 - (\text{Cb} \times \text{Tai}) / (\text{Cai} \times \text{Tb})\} \times 100$$

wherein the variables in the formula have the following meanings:

5 Cb: the number of insects before the treatment in the non-treated area;

 Cai: the number of insects at the time of observation in the non-treated area;

 Tb: the number of insects before the treatment in the treated area;
10 and

 Tai: the number of insects at the time of observation in the treated area.

 As a result, the present compounds (1), (2), (3), (5), (9), (10), (13), (15) to (19), (21), (26), (27), (29), (33) to (39), (42) to (47), (51), (54) to (64), (66),
15 (67), (72), (73), (75), (77) to (83), (85) to (92), (94) to (110), (112), (113), (115) to (118), (120), (121), (122), (124), (125), (129), (130), (134), (135), (137), (138), (142) to (151), (154) to (170), (172), (173), (175) to (180), (182), (183), (185), (187) to (191) had the control value of 90% or higher.

 Test Example 2 Insecticidal Test against cotton aphids (*Aphis gossypii*)
20

 A formulation of a test compound obtained in Formulation Example 5 was diluted with water so that the active ingredient concentration came to 500 ppm to prepare a test chemical solution.

 Cucumber seedlings (at the stage of developing the first foliage leaf)
25 cultivated in polyethylene cups each having 5 holes of 5 mm in diameter at the bottom were treated with the test chemical solution, which had been prepared as described above and was absorbed in a volume of 55 ml from the bottom of the cups. The cucumber plants were then left in a greenhouse at

25°C for 6 days, and about 20 cotton aphids were made parasitic thereon. On the 6th day after the application, the number of cotton aphids was examined, and the control value was determined in the same manner as described in Test Example 1.

5 As a result, the present compounds (9), (21), (33), (36), (39), (42), (44), (45), (51), (63), (80), (83), (88), (96), (101), (104), (105), (106), (108), (110), (112), (115), (120), (121), (124), (137), (138), (144), (148), (149), (157), (158) and (185) had the control value of 90% or higher.

Test Example 3 Insecticidal test against western flower thrips
10 (*Frankliniella occidentalis*)

A formulation of a test compound obtained in Formulation Example 5 was diluted with water so that the active ingredient concentration came to 500 ppm to prepare a test spray solution.

The seeds of cucumber were planted in polyethylene cups and grown
15 until their first foliage leaves developed, and the above test spray solution was sprayed at the rate of 20 ml/cup onto these cucumber plants. After the chemical solution sprayed onto the cucumber plants was dried, the first foliage leaves were cut off and placed on a water-containing filter paper (70 mm in diameter) in a polyethylene cup (110 mm in diameter). Thirty larvae of
20 western flower thrips (*Frankliniella occidentalis*) were set free in the polyethylene cup, and the lid was put on the polyethylene cup. After 7 days, the number of surviving insects was examined.

As a result, for the present compounds (1), (2), (3), (5), (6), (7), (13), (15), (16), (17), (19), (20), (21), (29), (34) to (39), (43), (44), (45), (51), (54), (55),
25 (58), (59), (63), (77), (80), (82), (83), (88), (134), (149), (150), (169), (176), (180), (182), (183), (190) and (191), the number of surviving insects on the leaves treated with each of these compounds was zero.

Test Example 4 Insecticidal test against silverleaf whiteflies (*Be-*

misia argentifoli)

A formulation of a test compound obtained in Formulation Example 5 was diluted with water so that the active ingredient concentration came to 500 ppm to prepare a test spray solution.

5 The seeds of cabbage were planted in polyethylene cups and grown until their first foliage leaves developed. The first foliage leaves were left and the other leaves were cut off. Some adults of silverleaf whiteflies were set free on the cabbage plants and allowed to lay eggs for about 24 hours. The cabbage plants with about 80 to 100 eggs thus laid were left in a green-
10 house for 8 days, and the above test spray solution was sprayed at the rate of 20 ml/cup onto the cabbage plants with larvae being hatched from the laid eggs. On the 7th day after the application, the number of surviving larvae was counted.

As a result, for the present compounds (1) to (6), (9), (10), (13) to (16),
15 (19), (21), (26) to (29), (33) to (39), (42) to (47), (49), (51), (52), (54) to (63), (66), (67), (68), (73), (76) to (83), (85) to (92), (94) to (105), (110), (112), (113), (115) to (118), (120), (121), (124), (130), (131), (134) to (138), (142), (143), (144), (146), (147), (149) to (156), (158), (159), (160), (164) to (184), (187), (188), (190) and (191), the number of surviving larvae on the cabbage leaves
20 treated with each of these compounds was not greater than 10.

Test Example 5 Insecticidal Test against brown planthoppers (*Nilaparvata lugens*)

A formulation of a test compound obtained in Formulation Example 5 was diluted with water so that the active ingredient concentration came to
25 500 ppm to prepare a test spray solution.

Fifty grams of molding Bonsoru 2 (available from Sumitomo Chemical Co., Ltd.) was put into a polyethylene cup, and 10 to 15 seeds of rice were planted in the polyethylene cup. The rice plants were grown until

the second foliage leaves developed and then cut into the same height of 5 cm. The test spray solution, which had been prepared as described above, was sprayed at the rate of 20 ml/cup onto these rice plants. After the chemical solution sprayed onto the rice plants was dried, thirty first-instar larvae of brown planthoppers (*Nilaparvata lugens*) were set free on the rice plants, which were then left in a greenhouse at 25°C. On the 6th day after the release of brown planthopper larvae, the number of brown planthoppers parasitic on the rice plants was examined.

As a result, in the treatment with each of the present compounds (5), (11), (21), (36), (38), (39), (55), (57), (59), (60), (61), (80), (83), (86), (87), (88), (91) to (94), (96), (97), (101), (104), (106), (109), (117), (118), (130), (147), (150), (151), (155), (167), (169), (172), (173), (174), (176), (178), (179), (181) to (185), (187), (188) and (191), the number of parasitic insects on the 6th day after the treatment was not greater than 3.

Test Example 6 Insecticidal Test against brown planthoppers (*Nilaparvata lugens*)

A formulation of a test compound obtained in Formulation Example 5 was diluted with water so that the active ingredient concentration came to 45.5 ppm to prepare a test spray solution.

Fifty grams of molding Bonsoru 2 (available from Sumitomo Chemical Co., Ltd.) was put into a polyethylene cup having a hole of 5 mm in diameter, and 10 to 15 seeds of rice were planted in the polyethylene cup. The rice plants were grown until the second foliage leaves developed and then treated with the test chemical solution, which had been prepared as described above and was absorbed in a volume of 55 ml from the bottom of the cup. The rice plants were left in a greenhouse at 25°C for 6 days and then cut into the same height of 5 cm. Thirty first-instar larvae of brown planthoppers (*Nilaparvata lugens*) were set free on the rice plants, which

were then left in the greenhouse at 25°C. On the 6th day after the release of brown planthopper larvae, the number of brown planthoppers parasitic on the rice plants was examined.


As a result, in the treatment with each of the present compounds (5),
5 (55), (57), (86), (87), (101), (110), (118), (144), (151), (167), (169), (172), (173),
(176), (181) and (185), the number of parasitic insects on the 6th day after the treatment was not greater than 3.

Industrial Availability

The present compounds have excellent pesticidal effect and they are
10 therefore used as the active ingredients of pesticidal compositions.

CLAIMS

1. A pyrimidine compound of formula (1):



- 5 wherein R¹ is C₃-C₇ alkynyl optionally substituted with halogen;
 R² and R³ are independently hydrogen, halogen or C₁-C₄ alkyl; and
 R⁴ is C₃-C₇ alkynyloxy optionally substituted with halogen; C₃-C₈
 cycloalkoxy optionally substituted with halogen, hydroxy, C₁-C₄ alkyl or C₁-C₄
 alkoxy; ethynyl substituted with C₃-C₆ alkyl or
 10 a group of formula -A¹R⁵ wherein A¹ is a single bond, oxygen, sulfur, carbonyl,
 $\diagup \text{C}=\text{CH}_2$ or $\diagup \text{C}=\text{NOR}^8$ wherein R⁸ is C₁-C₄ alkyl; when A¹ is a single bond
 oxygen, or sulfur, then R⁵ is optionally substituted phenyl or optionally sub-
 stituted C₇-C₉ aralkyl; or when A¹ is carbonyl $\diagup \text{C}=\text{CH}_2$ or $\diagup \text{C}=\text{NOR}^8$
 wherein R⁸ is C₁-C₄ alkyl, then R⁵ is optionally substituted phenyl; or
 15 a group of formula -NR⁶R⁷ wherein R⁶ is C₁-C₇ alkyl, C₁-C₃ haloalkyl,
 C₂-C₄ (alkoxymethyl), C₂-C₄ (haloalkoxymethyl), C₃-C₆ alkenyl, C₃-C₆ haloal-
 kenyl, C₃-C₇ alkynyl, cyanomethyl, optionally substituted phenyl, or option-
 ally substituted with C₇-C₉ aralkyl; and R⁷ is hydrogen, C₁-C₇ alkyl, C₁-C₃
 haloalkyl, C₂-C₄ (alkoxymethyl), C₂-C₄ (haloalkoxymethyl), C₃-C₆ alkenyl, C₃-
 20 C₆ haloalkenyl, C₃-C₇ alkynyl, cyanomethyl, optionally substituted phenyl, or
 optionally substituted C₇-C₉ aralkyl;

wherein the substituent in the optionally substituted phenyl and in
 the optionally substituted C₇-C₉ aralkyl is at least one selected from halogen,

hydroxy, cyano, nitro, phenyl, phenoxy, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, C₁-C₃ alkylthio, C₁-C₄ haloalkylthio, C₃-C₇ alkynyl-oxy, C₂-C₆ (alkoxyalkoxy), C₂-C₆ (alkylcarbonyl), and C₂-C₆ (alkylcarbonyl-oxy).

- 5 2. The pyrimidine compound according to claim 1, wherein R¹ is 2-propynyl optionally substituted with halogen, 2-butynyl optionally substituted with halogen, or 2-methyl-2-butynyl optionally substituted with halogen.
- 10 3. The pyrimidine compound according to claim 1, wherein R² and R³ are both hydrogen.
4. The pyrimidine compound according to claim 1, wherein R² is hydrogen and R³ is chlorine or fluorine.
- 15 5. The pyrimidine compound according to claim 1, wherein R¹ is C₃-C₇ alkynyl optionally substituted with halogen; R² and R³ are both hydrogen; and R⁴ is 2-halophenyl, 3-halophenyl, 2,3-dihalophenyl or 2,6-dihalophenyl.
- 20 6. The pyrimidine compound according to claim 1, wherein R¹ is C₃-C₇ alkynyl optionally substituted with halogen; R² and R³ are both hydrogen; and R⁴ is 2-halophenoxy, 3-halophenoxy, 2,3-dihalophenoxy or 2,6-dihalophenoxy.
7. The pyrimidine compound according to claim 1, wherein R¹ is C₃-C₇ alkynyl optionally substituted with halogen; R² and R³ are both hydrogen; and R⁴ is 2-halobenzyl, 3-halobenzyl, 2,3-dihalobenzyl or 2,6-dihalobenzyl.
- 25 8. The pyrimidine compound according to claim 1, wherein R¹ is C₃-C₇ alkynyl optionally substituted with halogen; R² and R³ are both hydrogen; R⁴ is a group of formula -N(C₂H₅)R⁷; and R⁷ is C₃-C₆ alkynyl optionally substituted with halogen, C₃-C₅ alkenyl optionally substituted

with halogen or C₂-C₃ alkyl.

9. A pyrimidine compound according to claim 1, which is 4-(2-pentynyloxy)-6-(2-propynyloxy)pyrimidine.

10. A pyrimidine compound according to claim 1, which is 4-(2-butynyloxy)-6-(2,6-difluorophenoxy)pyrimidine.

11. A pyrimidine compound according to claim 1, which is 4-(2-butynyloxy)-6-(2,3-difluorophenoxy)pyrimidine.

12. A pyrimidine compound according to claim 1, which is 4-(2-fluorophenyl)-6-(2-butynyloxy)pyrimidine.

13. A pyrimidine compound according to claim 1, which is 4-(3-fluorophenyl)-6-(2-butynyloxy)pyrimidine.

14. A pyrimidine compound according to claim 1, which is 4-(2,3-difluorophenoxy)-6-(2-pentynyloxy)pyrimidine.

15. A pyrimidine compound according to claim 1, which is 4-(3-fluorophenoxy)-6-(2-butynyloxy)pyrimidine.

16. A pyrimidine compound according to claim 1, which is 4-(2-fluorophenoxy)-6-(2-butynyloxy)pyrimidine.

17. A pyrimidine compound according to claim 1, which is 4-(N-methyl-N-(2,3-difluorophenyl)amino)-6-(2-butynyloxy)pyrimidine.

18. A pyrimidine compound according to claim 1, which is 4-(2,6-difluorobenzyl)-6-(2-butynyloxy)pyrimidine.

19. A pyrimidine compound according to claim 1, which is 4-(N-ethyl-N-phenylamino)-6-(2-butynyloxy)pyrimidine.

20. A pyrimidine compound according to claim 1, which is 4-(2,3-difluorophenyl)-6-(2-butynyloxy)pyrimidine.

21. A pyrimidine compound according to claim 1, which is 4-(2-fluorophenyl)-6-(2-pentynyloxy)pyrimidine.

22. A pyrimidine compound according to claim 1, which is 4-(2-

fluorobenzyl)-6-(2-butynyloxy)pyrimidine.

23. A pyrimidine compound according to claim 1, which is 4-(2-chlorobenzyl)-6-(2-butynyloxy)pyrimidine.

24. A pyrimidine compound according to claim 1, which is 4-(2-
5 butynyloxy)-6-(α -methylbenzyl)pyrimidine.

25. A pyrimidine compound according to claim 1, which is 4-(2-butynyloxy)-6-(2,3-difluorobenzyl)pyrimidine.

26. A pyrimidine compound according to claim 1, which is 4-(2-butynyloxy)-6-(2-chloro-6-fluorobenzyl)pyrimidine.

10 27. A pyrimidine compound according to claim 1, which is 4-(2-butynyloxy)-6-(N-ethyl-N-n-propylamino)pyrimidine.

28. A pyrimidine compound according to claim 1, which is 4-(2-butynyloxy)-6-(N-ethyl-N-iso-propylamino)pyrimidine.

29. A pyrimidine compound according to claim 1, which is 4-(2-
15 butynyloxy)-6-(N-ethyl-N-2,2,2-trifluoroethylamino)pyrimidine.

30. A pyrimidine compound according to claim 1, which is 4-(2,3-difluorophenyl)-6-(2-hexynyloxy)pyrimidine.

31. A pyrimidine compound according to claim 1, which is 4-(2-butynyloxy)-6-(1-(2-fluorophenyl)ethyl)pyrimidine.

20 32. A pyrimidine compound according to claim 1, which is 4-(2-butynyloxy)-6-(2-chloro-3,6-difluorobenzyl)pyrimidine.

33. A pyrimidine compound according to claim 1, which is 4-(2-fluorophenyl)-6-(1-methyl-2-butynyloxy)pyrimidine.

34. A pyrimidine compound according to claim 1, which is 4-(2-
25 butynyloxy)-6-(cyclopentyloxy)pyrimidine.

35. A pyrimidine compound according to claim 1, which is 4-(2-butynyloxy)-6-(cyclohexyloxy)pyrimidine.

36. A pyrimidine compound according to claim 1, which is 4-(2-

butynyloxy)-6-(N-ethyl-N-(2,3-difluorophenyl)amino)pyrimidine.

37. A pyrimidine compound according to claim 1, which is 4-(2,3-difluorophenyl)-6-(1-methyl-2-butynyloxy)pyrimidine.

38. A pyrimidine compound according to claim 1, which is 4-(2-butynyloxy)-6-(trans-2-methylcyclopentyloxy)pyrimidine.

39. A pyrimidine compound according to claim 1, which is 4-(2-butynyloxy)-6-(cis-2-methylcyclohexyloxy)pyrimidine.

40. A pyrimidine compound according to claim 1, which is 4-(3-fluorophenyl)-6-(1-methyl-2-butynyloxy)pyrimidine.

41. A pyrimidine compound according to claim 1, which is 5-fluoro-4-(2-butynyloxy)-6-(2-fluorophenoxy)pyrimidine.

42. A pyrimidine compound according to claim 1, which is 5-fluoro-4-(2-butynyloxy)-6-(2-chlorophenoxy)pyrimidine.

43. A pesticidal composition comprising a carrier and a pyrimidine compound according to claim 1 as an active ingredient.

44. A method for pest control comprising applying an effective amount of a pyrimidine compound according to claim 1 to a pest or a habitat of the pest.

45. A pyrimidine compound selected from the group consisting of 4-chloro-6-(2-propynyloxy)pyrimidine, 4-chloro-6-(2-butynyloxy)pyrimidine, 4-chloro-6-(1-methyl-2-butynyloxy)pyrimidine, 4-chloro-6-(2-pentynyloxy)pyrimidine, 4-chloro-6-(2-fluorophenyl)pyrimidine, 4-chloro-6-(N-methyl-N-(2,3-difluorophenyl) amino)pyrimidine, 4-chloro-6-(2,6-difluorobenzyl)pyrimidine, 4-chloro-6-(N-ethyl-N-phenylamino)pyrimidine, 4-chloro-6-(2,3-difluorophenyl)pyrimidine, 4-chloro-6-(2-fluorobenzyl)pyrimidine, 4-chloro-6-(2-chlorobenzyl)pyrimidine, 4-chloro-6-(α -methylbenzyl)pyrimidine, 4-chloro-6-(2,3-difluorobenzyl)pyrimidine, 4-chloro-6-(2-chloro-6-fluorobenzyl)pyrimidine, 4-chloro-6-(1-(2-fluorophenyl)ethyl)pyrimidine, 4-chloro-6-(2-chloro-

3,6-difluorobenzyl)pyrimidine, 4-chloro-6-(cyclopentyloxy)pyrimidine, 4-chloro-6-(cyclohexyloxy)pyrimidine, 4-chloro-6-(N-ethyl-N-(2,3-difluorophenyl)amino)pyrimidine, 4-chloro-6-(trans-2-methylcyclopentyloxy)pyrimidine and 4-chloro-6-(cis-2-methylcyclohexyloxy)pyrimidine.

5 46. The pyrimidine compound according to claim 45, which is 4-chloro-6-(2-propynyloxy)pyrimidine.

 47. The pyrimidine compound according to claim 45, which is 4-chloro-6-(2-butyloxy)pyrimidine.

 48. The pyrimidine compound according to claim 45, which is 4-
10 chloro-6-(1-methyl-2-butyloxy)pyrimidine.

 49. The pyrimidine compound according to claim 45, which is 4-chloro-6-(2-pentyloxy)pyrimidine.

 50. The pyrimidine compound according to claim 45, which is 4-chloro-6-(2-fluorophenyl)pyrimidine.

15 51. The pyrimidine compound according to claim 45, which is 4-chloro-6-(N-methyl-N-(2,3-difluorophenyl) amino)pyrimidine.

 52. The pyrimidine compound according to claim 45, which is 4-chloro-6-(2,6-difluorobenzyl)pyrimidine.

 53. The pyrimidine compound according to claim 45, which is 4-
20 chloro-6-(N-ethyl-N-phenylamino)pyrimidine.

 54. The pyrimidine compound according to claim 45, which is 4-chloro-6-(2,3-difluorophenyl)pyrimidine.

 55. The pyrimidine compound according to claim 45, which is 4-chloro-6-(2-fluorobenzyl)pyrimidine.

25 56. The pyrimidine compound according to claim 45, which is 4-chloro-6-(2-chlorobenzyl)pyrimidine.

 57. The pyrimidine compound according to claim 45, which is 4-chloro-6-(α -methylbenzyl)pyrimidine.

58. The pyrimidine compound according to claim 45, which is 4-chloro-6-(2,3-difluorobenzyl)pyrimidine.

59. The pyrimidine compound according to claim 45, which is 4-chloro-6-(2-chloro-6-fluorobenzyl)pyrimidine.

5 60. The pyrimidine compound according to claim 45, which is 4-chloro-6-(1-(2-fluorophenyl)ethyl)pyrimidine.

61. The pyrimidine compound according to claim 45, which is 4-chloro-6-(2-chloro-3,6-difluorobenzyl)pyrimidine.

62. The pyrimidine compound according to claim 45, which is 4-
10 chloro-6-(cyclopentyloxy)pyrimidine.

63. The pyrimidine compound according to claim 45, which is 4-chloro-6-(cyclohexyloxy)pyrimidine.

64. The pyrimidine compound according to claim 45, which is 4-chloro-6-(N-ethyl-N-(2,3-difluorophenyl)amino)pyrimidine.

15 65. The pyrimidine compound according to claim 45, which is 4-chloro-6-(trans-2-methylcyclopentyloxy)pyrimidine.

66. The pyrimidine compound according to claim 45, which is 4-chloro-6-(cis-2-methylcyclohexyloxy)pyrimidine.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau




(43) International Publication Date
28 March 2002 (28.03.2002)

PCT

(10) International Publication Number
WO 02/024663 A3

- (51) International Patent Classification⁷: C07D 239/52, 239/34, 239/46, 239/30, 239/42, A01N 43/54
- (74) Agents: AOYAMA, Tamotsu et al.; Aoyama & Partners, IMP Building, 3-7, Shiromi 1-chome, Chuo-ku, Osaka-shi, Osaka 540-0001 (JP).
- (21) International Application Number: PCT/JP01/07766
- (22) International Filing Date:
7 September 2001 (07.09.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
2000-283113 19 September 2000 (19.09.2000) JP
2001-142975 14 May 2001 (14.05.2001) JP
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): SUMITOMO CHEMICAL COMPANY, LIMITED [JP/JP]; 5-33, Kitahama 4-chome, Chuo-ku, Osaka-shi, Osaka 541-0041 (JP).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): MIZUNO, Hajime [JP/JP]; 4-9-17-323, Sakuragaoka, Minoo-shi, Osaka 562-0046 (JP). SAKAMOTO, Noriyasu [JP/JP]; 4-1-4, Kitasakurazuka, Toyonaka-shi, Osaka 560-0022 (JP). KINOSHITA, Yoshiharu [JP/JP]; 4-9-17-329, Sakuragaoka, Minoo-shi, Osaka 562-0046 (JP).
- Published:
— with international search report
- (88) Date of publication of the international search report:
9 October 2003
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: PYRIMIDINE COMPOUNDS AND THEIR USE



(1)

(57) Abstract: Pyrimidine compounds of formula (1), wherein R¹ is C₃-C₇ alkynyl optionally substituted with halogen; R² and R³ are independently hydrogen or the like; and R⁴ is C₃-C₇ alkynyloxy optionally substituted with halogen, optionally substituted phenyl, or the like have an excellent pesticidal effect.

WO 02/024663 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/JP 01/07766

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D239/52 C07D239/34 C07D239/46 C07D239/30 C07D239/42
A01N43/54

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DE 40 29 654 A (HOECHST) 2 April 1992 (1992-04-02) claims; table 1	1,43,44
A	CHEMICAL ABSTRACTS, vol. 131, no. 9, 1999 Columbus, Ohio, US; abstract no. 116250j, XP002191469 abstract & JP 11 193277 A (NIPPON SODA) 21 July 1999 (1999-07-21)	1,43,44
A	GB 2 078 222 A (SPOFA) 6 January 1982 (1982-01-06) page 1 -page 3	1
	--- -/-	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

26 February 2002

Date of mailing of the international search report

13/03/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Francois, J

INTERNATIONAL SEARCH REPORT

Internati. application No
PCT/JP 01/07766

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 972 770 A (AMERICAN CYANAMID) 19 January 2000 (2000-01-19) figures IV,X,XXIV	45
A	page 1	1,43
A	C.A. RASMUSSEN: "FACTORS INFLUENCING THE OCCURENCE OF AN SN(ANRORC)MECHANISM IN THE AMINO-DEHALOGENATION OF 4-SUBSTIT. 6-HALOGENOPYRIMIDINES" RECUEIL DES TRAVAUX CHIMIQUES DES PAYS-BAS., vol. 98, no. 1, January 1979 (1979-01), pages 5-8, XP002191468 ELSEVIER SCIENCE PUBLISHERS. AMSTERDAM., NL ISSN: 0165-0513 page 5; example 3E	45,53

INTERNATIONAL SEARCH REPORT

Internat. application No
PCT/JP 01/07766

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
DE 4029654	A	02-04-1992	DE 4029654 A1	02-04-1992
			WO 9205159 A1	02-04-1992
			ZA 9107429 A	27-05-1992
<hr/>				
JP 11193277	A	21-07-1999	NONE	
<hr/>				
GB 2078222	A	06-01-1982	CS 216348 B1	29-10-1982
			BE 889204 A1	01-10-1981
			CA 1138455 A1	28-12-1982
			DE 3122836 A1	01-04-1982
			DK 258281 A	14-12-1981
			FI 811610 A	14-12-1981
			FR 2484416 A1	18-12-1981
			JP 57016871 A	28-01-1982
			NL 8102670 A	04-01-1982
			SE 8103702 A	14-12-1981
			US 4384118 A	17-05-1983
<hr/>				
EP 972770	A	19-01-2000	AT 205835 T	15-10-2001
			DE 69900297 D1	25-10-2001
			EP 0972770 A1	19-01-2000
			JP 2000080084 A	21-03-2000
<hr/>				